

GALLONS NET

# RODINE

LIQUID EXTRACT OF RED SQUILL

FOR RODENTICIDE USE ONLY


Minimum toxicity LD 50 for male rats 500 mg. kg.

ACTIVE INGREDIENT:

\* Extractives of red squill

INERT INGREDIENTS

\* Toxicity LD 50 for male rats of red squill extractives mg. kg.



**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
Keep away from domestic animals and foodstuff.

Meat and fish baits will attract dogs and cats, and when eaten will make these pets violently ill. Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.

## DIRECTIONS FOR USE

Rodine should be mixed with ground fresh beef, fish, cereal or ground grain in the proportion of one part of Rodine with nine parts of bait material. Place tablespoonful quantities of the bait at about 10 foot intervals where rats feed, water, or travel. Bait placements along walls, in corners, and in concealed locations where rats can feed unobserved are desirable. Before placing baits, exposed food should be removed if possible. Refuse receptacles should be covered. Place baits in late afternoon or evening where rats have been seen. If uncertain which food rats prefer, place baits without Rodine as a test. Follow in two to three days with the Rodine-bait combination. Two or more varieties of bait may be used in one procedure. After three days, collect uneaten baits and destroy.