



NET WT. POUNDS

DETHMOR

'DEATH TO MICE AND RATS

ACTIVE INGREDIENT

WARFARIN

(3-(α -acetylbenzyl)-4-hydroxy-2-methyl-5-pyridinyl)-2-methylimidazole

INERT INGREDIENTS

*Only WARFARIN tested and certified for purity by the U.S. Government.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

DETHMOR contains as its active ingredient an anticoagulant chemical which if taken by humans, domestic animals, wildlife or pets may reduce the clotting ability of the blood and serious hemorrhage may result. In case baits are eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

Do not reuse container. Destroy when empty.

NOTE FOR PHYSICIANS

When a human has been known to have ingested DETHMOR, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol[®].

DIRECTIONS FOR THE MANUFACTURE AND USE OF RAT AND MOUSE BAITS

Mix thoroughly one part of DETHMOR with 19 parts, by weight, of suitable semi-permanent bait material such as fresh corn meal, rolled oats, nut crumbs, poultry mash, etc. A small amount, up to 5% of a bland oil, such as mineral or corn oil or any other edible oil, may be used to prevent excessive dusting where the mixture has a tendency to dust.

When a perishable bait such as fresh ground meat, fish, vegetable, or similar material is used, bait placement should be made in the late afternoon and replaced daily with fresh baits.

DIRECTIONS FOR USE OF RAT BAIT:

Place 4 to 6 ounces of bait in dry container next to burrows, in runways, or where rats are accustomed to feed. Bait should be placed in locations

not accessible to children, pets, wildlife and domestic animals or in tamper-proof bait boxes. Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped. Replace stale, damp or moldy baits with fresh bait.

DIRECTIONS FOR USE OF MOUSE BAIT:

Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations should be placed at intervals of 8 to 12 feet. Assure an uninterrupted supply of bait for a period not less than 15 days.

IMPORTANT NOTICE:

Baiting Should Continue in All Cases Until Complete Lack of Feeding Is Noted.