

DIRECTIONS FOR USE (continued)

TO PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, begonias and coleus for control of aphids, exposed thrips, adult whiteflies, spider mites, and fungus gnats. Use with caution. To ascertain possible phytotoxicity, spray only a few plants, wait 24 hours, and check for possible injury.

FOR PROTECTION OF WOOLENS: Apply only to garments, blankets, carpets, and other woolen fabrics before placing in storage for protection from larvae of webbing clothing moths and black carpet beetles.

Dry clean or wash garments and blankets before treating. Clean carpets, draperies and other fabrics by thoroughly brushing before treating.

Hold the container 14 inches to 18 inches (30 to 46 cm) from the fabric and apply the spray while moving evenly in a forward and back motion across the fabric while covering one foot (30 cm) per second. Spray the entire fabric surface until it is slightly moist or damp to the touch. When possible treat both sides of the fabric. Do not soak or wet fabric. For maximum protection place treated fabrics after drying in plastic or paper bags. Also treat cracks and crevices of closets, chests and trunks where these fabrics are being stored. Treat the cracks and crevices at the rate of two seconds per linear foot (30 cm).

When applied as directed on fabric and in storage areas this treatment will offer protection for 6 months. To insure protection inspect the fabric in the storage area once every three months. Repeat application only when necessary. Avoid heavy applications. When possible make application outdoors. If application is made indoors do not remain in treated areas. Ventilate the area after treatment is complete.

OUTDOORS

TO KILL FLYING INSECTS: Spray areas infested with mosquitoes, gnats (midges), houseflies, stable flies, deer flies, and flying moths (millers). Follow precautions listed above. Hold container 18 to 24 inches (46 to 61 cm) from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard (28.3 cu m) of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

TO PROTECT ORNAMENTAL PLANTS: To control aphids, exposed thrips, adult whiteflies, leafhoppers, gypsy moths, plant bugs, spittlebugs and Japanese beetles on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyon, viburnum, euonymus, oaks, maples, locust, forsythia and althea, and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 18 inches (46 cm) from the infested plants. Spray in short spray bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, iris, zinnia, snapdragons, geraniums, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatshedera, ivy, tobira, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above.

Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus may also be treated to control

DIRECTIONS FOR USE (continued)

TO PROTECT ORNAMENTAL PLANTS (continued):

the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

To Kill Tent Caterpillars: Application should be made when caterpillars are young and tents are first noticed (usually late spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

To Kill Gypsy Moth Caterpillars: Application should be made when caterpillars are young and are first noticed (usually late spring). Spray plants evenly to the point of run-off, applying to both sides of leaves and other areas of plants where insects are suspected. For dense evergreens and shrubs, be sure spray reaches interior leaves and branches.

TO CONTROL NUISANCE PESTS:

CRICKETS, SILVERFISH, EARWIGS, SPIDERS, SOWBUGS, CENTIPEDES, AND MILLIPEDES: Same as directed above for indoor use.

ANTS: Spray trails, nests and points of entry.

Spray on ants where possible. Repeat as necessary.

FIRE ANTS: Kills and temporarily reduces annoyance of these insects. To reduce annoyance from these ants and keep them from entering homes, spray patios, barbecue areas, driveways, door sills, window sills, other entrance areas, foundations of buildings, ant trails, around pools and areas where the ants are noticed. Spray until the area is wet contacting as many insects as possible. This treatment will not kill the colonies but will reduce the immediate population for up to four hours. Repeat as necessary. Avoid contacting plants, shrubbery and large areas of lawns.

FLEAS AND TICKS: Same as directed for indoor use.

SPIDERS (Including Black Widow and Brown Recluse): Direct short bursts of spray on spiders and repeat as necessary.

WASPS, HORNETS, YELLOW JACKETS AND BEES: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

1982

327 / 85585
11 / 4

Penick Corporation
Pesticides Technical Support Group
1050 Wall Street West
Lyndhurst, NJ 07071

Attention: Mrs. Emily Renal

Subject: SSP-1332 Aqueous Pressurized Spray Insecticide
0.25% for House and Garden
EPA Registration No. 432-452
Your Letter of July 27, 1982


The amendment referred to above, submitted in connection with registration under FIFRA sec. 3(c)(7)(A), is acceptable provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


Franklin D. R. Gee
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (13-767)

Enclosure

IRB/RD:SCROEDER/GEE:DCR-23723:WANG-1677C:sc:Raven:479-2013:12/16/82

PHS
6



PENICK CORPORATION

4/2/82
(CENTER PANEL)

SBP-1382^(R) INSECTICIDE AQUEOUS PRESSURIZED SPRAY 0.25%

FOR HOUSE AND GARDEN

Authorized ^{DA} for use in edible product areas of official establishments operating under The Meat, Poultry, Shell Egg Grading, and Egg Products Inspection Program.

ACTIVE INGREDIENTS:

*^oResmethrin {5-(phenylmethyl)-3-furanyl)methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate 0.25% w/w

INERT INGREDIENTS: 99.75%
100.00% w/w

*Cis/trans isomers ratio max. 30% (+)cis and min. 70% (+)trans.

+U.S. Patents Nos. 3,465,007 and 3,542,928.

*Penick's SBP-1382^(R) brand of resmethrin insecticide.

Penick Corp., 1050 Wall Street West, Lyndhurst, N.J. 07071.

PRECAUCION AL CONSUMIDOR: SI USTED NO LEES INGLES, NO USE ESTE PRODUCTO HASTA QUE LA ETIQUETA LE HAYA SIDO EXPLICADA AMPLIAMENTE.

TRANSLATION: (TO THE USER: IF YOU CANNOT READ ENGLISH, DO NOT USE THIS PRODUCT UNTIL THE LABEL HAS BEEN FULLY EXPLAINED TO YOU.)

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

SHAKE WELL BEFORE USING

EPA Reg. No. 432-452

EPA Est. No. 432-NJ-1

NET ~~WEIGHT~~:
4.12oz

AGREEMENT

Penick Corporation
1050 Wall Street West
Lyndhurst, New Jersey 07071

432-452

(SIDE PANEL)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, with potable water. *In the home,* all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If in eyes or on skin: Flush with plenty of water. Get medical attention if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54.5°C) may cause bursting.

Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazards. Do not use on surfaces that can be harmed or stained by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISPOSAL

DO NOT REUSE EMPTY CONTAINER. WRAP CONTAINER AND PUT IN TRASH COLLECTION.

For use in and around hotels, motels, homes, greenhouses and lathouses, food handling establishments, packaging and processing plants, seed houses, commercial, industrial office, institutional and storage buildings, milk handling areas of milk, cheese and ice cream plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

DIRECTIONS FOR USE (continued)

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

INDOORS

FOR HOUSEFLIES, MOSQUITOES, GNATS, FLYING MOTHS (Millers), FRUIT FLIES AND VINEGAR FLIES: Close all doors and windows. Point nozzle upward; direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

FOR CRAWLING INSECTS: COCKROACHES (waterbugs), CARWIGS, GRAIN MITES, SOWBUGS (pillbugs), SILVERFISH (fire brats), SPIDERS, CENTIPEDES, MILLIPEDES, FLEAS, ANTS, BOOKLICE AND CRICKETS: Spray thoroughly and frequently into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, door frames, bookcases and storage areas. Spray directly on insects where possible. Repeat as necessary.

TO KILL ANTS - INCLUDING FIRE ANTS: Spray doors, around window frames, and other points of entry. Spray ant trails, nests, and hills contacting as many ants directly as possible.

TO KILL BEDBUGS: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds, and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

TO KILL FLEAS AND TICKS ON DOGS, CATS AND PREMISES: Apply directly to animals along back, under forelegs, and on other areas except scrotum and anus. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes or mouth of animal. Repeat within twenty-four hours of first treatment, if necessary. To reduce reinfestation, thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 16 inches (30 to 46 cm). Also spray the entire inside surface of dog houses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible.