

CONFIDENTIAL UNTIL
ACCEPTED BY EPA

PROPOSED AMENDED LABEL 6/7/85

SBP-1382® INSECTICIDE AQUEOUS PRESSURIZED SPRAY 0.35% FOR HOUSE AND GARDEN

Authorized by USDA for use in edible product areas of official establishments operating under The Meat, Poultry, Shell Egg Grading, and Egg Products Inspection Program.

ACTIVE INGREDIENTS:

Resmethrin [5-(phenylmethyl)-3-furanyl]methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate 0.35% w/w

INERT INGREDIENTS: 99.65%

100.00% w/w

*Cis/trans isomers ratio max. 30% (±) cis and min. 70% (±) trans.
(U.S. Patent Nos. 3,488,897 and 3,542,828.)
Penick's SBP-1382® brand of resmethrin insecticide.
Penick Corp., 1056 Wall Street West, Lyndhurst, N.J. 07071

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada completamente.
(To the user: If you cannot read English, do not use this product until the label has been fully explained to you.)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled with possible contact in the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.
Remove plants, birds and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT
If in eyes or on skin: Flush with plenty of water. Get medical attention if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS
Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 110°F (54.5°C) may cause bursting. Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazards. Do not use on surfaces that can be harmed or stained by water.

SPECIAL LABEL

SHAKE WELL BEFORE USING

EPA Reg. No. 432-454

EPA Est. No. 432-NJ-1

NET WT.

PENICK Penick Corporation

1050 Wall Street West, Lyndhurst, N.J. 07071 USA

ACCEPTED
JUL 17 1985
EPA REG. NO. 432-454

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage: Keep in locked storage area.
Container Disposal: Replace cap and discard container in trash.

DIRECTIONS FOR USE continued on reverse side.

↑ thoroughly with an effective cleaning compound and rinse

BEST AVAILABLE COPY

hotels, motels, homes, greenhouses and other establishments, packaging and processing areas, commercial, industrial, office, and public buildings, milk handling areas of milk plants, milk storage rooms, egg processing plants, restaurants, bottling plants, canneries, dairies, boats, ship cabins and hospitals. Insects should be removed prior to treatment. Also use on trailers, trucks, boxcars and ship cargo holds, dog kennels and horse stables. In horse stables, use only if horses will be used for food.

Use: Remove protective cap. Aim spray nozzle. Push button to spray.

INDOORS

MOSQUITOES, GNATS, FLYING MOTHS AND VINEGAR FLIES: Close all doors and windows; direct the spray mist to all parts of the room with mist, then leave the treated room for at least 15 minutes. Ventilate the room as usual.

PESTS: COCKROACHES (Waterbugs), TERmites, SOWBUGS (Pillbugs), SILVERFISH BEETLES, CENTIPEDES, MILLIPEDES, FLEAS, AND CRICKETS: Spray thoroughly and in places such as cracks, crevices, moist areas, pipes and sinks, under refrigerators, and bookcases and storage areas. Spray as possible. Repeat as necessary.

SPRING FIRE ANTS: Spray doors, around hot points of entry. Spray ant trails, nests, and many ants directly as possible.

AND LICE: Apply as a spot treatment on devices around baselards, floorboards, tiles, first, thoroughly clean and sand springs. Then spot treat only areas with pests, concentrating on tufts, folds, etc. as necessary. For persistent infestation, consult professional applicator.

TICKS ON DOGS, CATS AND PREMISES: Spray along back, under forelegs, and on chest and anus. Part hair to contact skin. Hold nozzle about 1 foot (30 cm) away from skin. Thoroughly moist. Do not spray on face of animal. Repeat within 24 hours of first application. To reduce reinfestation, thoroughly spray dog and other resting areas until the

surface is slightly moist. Treat these areas at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30 to 46 cm). Also spray the entire inside surface of dog houses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible.

TO PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, begonias, geraniums and coleus for control of aphids, exposed thrips, adult whiteflies, spider mites, and fungus gnats. Use with caution. To ascertain possible phytotoxicity, spray only a few plants, wait 24 hours, and check for possible injury.

OUTDOORS

TO KILL FLYING INSECTS: Spray areas infested with mosquitoes, gnats (midges), house flies, stable flies, deer flies and flying moths (millers). Follow precautions listed above. Hold container 18 to 24 inches (46 to 61 cm) from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard (0.76 cu. m) of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

TO PROTECT ORNAMENTAL PLANTS: To control aphids, exposed thrips, adult whiteflies, leafhoppers, gypsy moths, plant bugs, spittlebugs and Japanese beetles on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, loyca, viburnum, euonymus, oaks, maples, locust, forsythia and althea, and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 18 inches (46 cm) from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot (30 cm) of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, iris, zinnia, snapdragons, ageratum, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatsy, ivy, lobelia, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above.

Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus may also be treated to control the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

TO KILL TENT CATERPILLARS: Application should be made when caterpillars are young and tents are first noticed (usually late

spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

TO KILL GYPSY MOTI; CATERPILLARS: Application should be made when caterpillars are young and first noticed (usually late spring). Spray plants evenly to the point of run-off, applying to both sides of leaves and other areas of plants where insects are suspected. For dense evergreens and shrubs, be sure spray reaches interior leaves and branches.

TO CONTROL NUISANCE PESTS: CRICKETS, SILVERFISH, EARWIGS, SOWBUGS, CENTIPEDES AND MILLIPEDES: Same as directed above for indoor use.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FIRE ANTS: Kills and temporarily reduces annoyance of these insects. To reduce annoyance from these ants and keep them from entering homes, spray patios, barbecue areas, driveways, door sills, window sills, other entrance areas, foundations of buildings, ant trails, around pools and areas where the ants are noticed. Spray until the area is wet contacting as many insects as possible. This treatment will not kill the colonies, but will reduce the immediate population for up to four hours. Repeat as necessary. Avoid contacting plants, shrubbery and large areas of lawns.

FLEAS AND TICKS: Same as directed for indoor use.

SPIDERS: (Including Black Widow and Brown Recluse): Direct short bursts of spray on spiders and repeat as necessary.

WASPS, HORNET, YELLOW JACKETS AND BEES: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

BEST AVAILABLE COPY

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such, or the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

CONFIDENTIAL UNTIL
ACCEPTED BY EPA

PROPOSED AMENDED LABEL 6/7/85