



AgrEvo

PH 03

432-454

9-3-98

1672

A company of Hoechst and Schering, Berlin

SBP-1382[®] INSECTICIDE AQUEOUS PRESSURIZED SPRAY 0.35% FOR HOUSE AND GARDEN

ACCEPTED
SEP 1998
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 432-454

* AUTHORIZED BY USDA FOR USE IN EDIBLE PRODUCT AREAS OF OFFICIAL ESTABLISHMENTS OPERATING UNDER THE MEAT, POULTRY, SHELL EGG GRADING, AND EGG PRODUCTS INSPECTION PROGRAM.

ACTIVE INGREDIENTS:

*Resmethrin [5-(phenylmethyl)-3-furanyl] methyl 2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate.....	0.35% w/w
OTHER INGREDIENTS:	99.65%
	100.00% w/w

*Cis/trans isomers ratio max. 30% (±)cis and min. 70% (±)trans.
†AgrEvo Environmental Health, Inc.'s SBP-1382® brand of resmethrin insecticide.
SBP-1382 is a registered trademark of AgrEvo Environmental Health, Inc.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(To the user: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 432-454

EPA EST. NO.

KEEP OUT OF REACH OF CHILDREN

EXCISEMENT LABEL

FIRST AID

IF IN EYES OR ON SKIN: Flush with plenty of water. Get medical attention if irritation persists.

See Side Panel For Additional
Precautionary Statements

SHAKE WELL BEFORE USING

NET WEIGHT:

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. Wash thoroughly with soap and water after handling.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After space spraying in meat packing plants, bakeries and other food processing plants, wash all equipment, benches, shelving, etc., where exposed food will be handled, thoroughly with an effective cleaning compound and use with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use.

Remove pets, birds and caged fish aquariums before spraying.

Environmental Hazards

This pesticide is highly toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present onto intertidal areas below the mean high water mark. Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

Physical And Chemical Hazards

Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F (54.5°C) may cause bursting. Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazards. Do not use on surfaces that can be harmed or stained by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL
Storage: Keep in locked storage area.
Container Disposal: Replace cap and discard container in trash.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantings, tapes, ornamental gardens or parks or lawns and grounds.

For use in and around hotels, motels, homes, food handling establishments, packaging and processing plants, seed houses, commercial, industrial, office, institutional and storage buildings, milk handling areas of milk, cheese and ice cream plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in

SBP-1382® INSECTICIDE

horse stables. In horse stables, not for application if horses will be used for food.

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

INDOORS

FOR HOUSE FLIES, MOSQUITOES, GNATS, FLYING MOTHS (Millers), FRUIT FLIES AND VINEGAR FLIES: Close all doors and windows. Point nozzle upward; direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

FOR CRAWLING INSECTS: Cockroaches (Waterbugs), Earwigs, Grain Mites, Sowbugs (Pillbugs), Silverfish (Firebrats), Spiders, Centipedes, Millipedes, Fleas, Ants, Booklice And Crickets: Spray thoroughly and frequently into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, door frames, bookcases and storage areas. Spray directly on insects where possible. Repeat as necessary.

TO KILL ANTS - INCLUDING FIRE ANTS: Spray doors, around window frames, and other points of entry. Spray ant trails, nests, and hills contacting as many ants directly as possible.

TO KILL BEDBUGS AND LICE: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor pests, concentrating on tufts, folds, and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

TO KILL FLEAS AND TICKS IN DOG AND CAT PREMISES: Thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of 2 seconds per linear foot (30 cm) at a distance of 12 to 18 inches (30 to 46 cm). Also spray the entire inside surface of doghouses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible. For best results, dogs and cats should also be treated with an appropriate insecticide.

TO PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, begonias, geraniums and coleus for control of *Aphids, Exposed Thrips, Adult Whiteflies, Spider Mites, And Fungus Gnats*. Use with caution. To ascertain possible phytotoxicity, spray only a few plants, wait 24 hours, and check for possible injury.

OUTDOORS

TO KILL FLYING INSECTS: Spray areas infested with *Mosquitoes, Gnats (Midges), House Flies, Stable Flies, Deer Flies And Flying Moths (Millers)*. Follow precautions listed above. Hold container 18 to 24 inches (46 to 61 cm) from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard (0.76 cu. m) of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

TO PROTECT ORNAMENTAL PLANTS: To control *Aphids, Exposed Thrips, Adult Whiteflies, Leafhoppers, Gypsy Moths, Plant Bugs, Spittlebugs And Japanese Beetles* on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyan, viburnum, euonymus, oaks, maples, locust, for-

sythia and althea, and mimosa webworm larvae on mimosa, spray directly on insects. Hold container 18 inches (46 cm) from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot (30 cm) of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, iris, zinnia, snapdragons, ageratum, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatsy, ivy, tobira, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittlebugs and Japanese beetles. Spray as directed above. Ornamental Quince, Juniper, Cypress, Ornamental Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus may also be treated to control the above pests, but use with extreme caution. Certain varieties have shown some leaf burn under certain conditions of use such as heavy dosage. Repeat as necessary to obtain control.

TO KILL TENT CATERPILLARS: Application should be made when caterpillars are young and tents are first noticed (usually late spring). For best results, apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

TO KILL GYPSY MOTH CATERPILLARS: Application should be made when caterpillars are young and first noticed (usually late spring). Spray plants evenly to the point of run-off, applying to both sides of leaves and other areas of plants where insects are suspected. For dense evergreens and shrubs, be sure spray reaches interior leaves and branches.

TO CONTROL NUISANCE PESTS: CRICKETS, SILVERFISH, EARWIGS, SOWBUGS, CENTIPEDES AND MILLIPEDES: Same as directed above for indoor use.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

FIRE ANTS: Kills and temporarily reduces annoyance of these insects. To reduce annoyance from these ants and keep them from entering homes, spray patios, barbecue areas, driveways, door sills, window sills, other entrance areas, foundations of buildings, ant trails, around pools and areas where the ants are noticed. Spray until the area is wet contacting as many insects as possible. This treatment will not kill the colonies, but will reduce the immediate population for up to four hours. Repeat as necessary. Avoid contacting plants, shrubbery and large areas of lawns.

FLEAS AND TICKS: Same as directed for indoor use.

SPIDERS: (Including Black Widow and Brown Recluse): Direct short bursts of spray on spiders and repeat as necessary.

WASPS, HORNETS, YELLOW JACKETS AND BEES: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

AgrEvo Environmental Health
95 Chestnut Ridge Road
Montvale, NJ 07645

In case of Medical emergencies or health
and safety inquiries or in case of fire,
leaking or damaged containers, information
may be obtained by calling 1-800-471-0660.

For Product Information Call Toll-Free:
1-800-331-2867