

BASE INSECTICIDE 0.20%

GOOD FOR CONTROLLING THE FOLLOWING INSECTS AND CRAWLING INSECTS:

moths (millers), wasps, hornets, yellow jackets, cockroaches, waterbugs, spiders, grain mites, sowbugs, centipedes and millipedes.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing mist. Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. Remove pets and cover fish aquariums before spraying.

FIRST AID: In case of contact, immediately flush skin and eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and ponds). Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS: Extinguish open flame before space spraying. This formulation is combustible. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Store at room temperatures above 32°F.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For use outdoors in yards, patios, picnic areas, campsites, drive-ins, around swimming pools, poultry houses. For use indoors and outside premises of hotels, motels, homes, food handling establishments, packaging and processing plants, seed houses, commercial, industrial, office, institutional and storage buildings, milk handling areas of milk plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins, and hospitals. **IN HOSPITALS,** patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars, ship cargo holds, and in dog kennels and horse stables. **IN HORSE STABLES, NOT FOR APPLICATION IF HORSES WILL BE USED FOR FOOD.**

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas apply only when the facility is not in operation.

Before application, remove plants and animals from area to be treated. Cover or remove fish aquariums. Remove horses before spraying stables.

STORAGE: Store product in original container in a locked storage area.
CONTAINER DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

DIRECTIONS FOR USE (Continued on reverse side)

Reviewed: _____
Registration: _____
Date: _____

0.200%w/w
0.027%
0.265%
0.500%
0.908%
0.000%w/w

P-12/11/89



22

ed)

EPA Reg. 432-504

contact these insects in cracks and
baseboards, around pipes, behind
other harborages. Spray these infested
thoroughly wet. Repeat as necessary.

Spray around doors and window
frames, and hills in addition to direct spray
on insects.

Spray directly on these pests and
apply to the webbing and all
places which harbor spiders.

Applications - This formulation is to be
used in only such portable thermal
fogger that can deliver spray particle
in aerosol-size range.

Use to control annoyance from flies, mosquitoes
around decks, patios, around swimming pools,
camp-sites, drive-ins, and horse stables.
Apply with applicator at least five feet from any
flowering plants and shrubbery. Avoid
spraying foliage. Apply undiluted at the rate
of 1/2 gallon per acre. Fog only when there is no
wind and wind velocity is less than 3 miles per
hour. Time is generally at dusk when the
air and the air temperature and ground
temperature are approximately the same. Under
these conditions the fog will linger in the treated
area for a long time. Apply fog while moving away
from the area. Avoid inhalation of the fog. Do not
enter the treated area until the fog has dissipated.
Repeat application as necessary.

Use: To kill nuisance pests around
lawns and paved areas: Spray directly on
spraying turf or plants.

Directions for indoor application.

Use: Yellow Jackets: Application
in late evening when insects are at
opening away from face. Aim spray
in bushes, under eaves, chimneys,

cracks and crevices. Thoroughly saturate nest with
spray, contacting as many insects as possible. Spray
liberally into hiding and breeding places.

**EARWIGS, sowbugs, spiders, centipedes and
millipedes:** Apply as a contact spray, as for indoor
use.

To control flies, mosquitoes and gnats and crawling
insects around premises, spray thoroughly around
doors, window frames and similar areas of entry.
Also thoroughly spray garbage and refuse areas,
especially around poultry houses, barns, and similar
areas of heavy infestation. Do not spray on turf or
plants.

WARRANTY STATEMENT

Our recommendations for use of this product are
based upon tests believed to be reliable. The use of
this product being beyond the control of the
manufacturer, no guarantee, expressed or implied,
is made as to the effects of such or the results to be
obtained if not used in accordance with directions or
established safe practice. The buyer must assume
all responsibility, including injury or damage,
resulting from its misuse as such, or in combination
with other materials.