[432-507]

YOUR BRAND SBP-1382[®]/BIOALLETHRIN (0.20% + 0.125%) AQUEOUS PRESSURIZED SPRAY FOR HOUSE AND GARDEN

NET WT._____

ACTIVE INGREDIENTS:

| * *(5-Benzyl-3-furyl)methyl 2,2-dimethyl-3- (2-methylpropenyl)cyclopropanecarboxylate | 0.200% w/w |
|--|------------|
| Related compounds | 0.027% |
| d-trans Allethrin (allyl homolog of Cinerin I) | 0.125% |
| Related compounds | 0.009% |
| Aromatic petroleum hydrocarbons | 0.265% |
| INERT INGREDIENTS: | 99.374% |
| | 100.000% |

*Cis/trans isomers ratio: max. 30% (±)cis and min. 70% (±)trans. *U.S. Patent No. 3,465,007

| SKAKE WELL BEFORE USI | ACCEPTED 492-2011 NOV 27 1972 |
|-------------------------|---|
| CAUTION: KEEP OUT OF RE | FUNCT FG ASCH OF CHILDRENDUBLEST TO ATTACHED COMMENTS. |

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Cover or remove fishbowls. Do not use on edible crops.

NOTICE – When used in bakerres, restaurants and food processing establishments, foods should be removed or covered before treatment. All food processing surfaces should be covered before treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130⁰F may cause bursting. Never throw container into fire or incinerator.

KILLS: Houseflies, mosquitoes, gnats, wasps, reaches, earwigs, two spotted mites, sowbugs, spiders, ants, biting flies, aphids (plant lice), leafhoppers, exposed thrips, adult whiteflies.



DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray

CONTROL HOUSEHOLD PESTS:

For Houseflies, Mosquitoes, and Gnats: Close all doors and windows. Point nozzle upward, direct spray mist to all parts of the room especially windows and other sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

For Crawling Insects: Roaches, earwigs, sowbugs and spiders: Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

OUTDOORS- TO REDUCE ANNOYANCE FROM FLYING INSECTS:

Reduce mosquitoes, gnats (midges), houseflies, biting flies and pyralid and noctuid flying moths by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

OUTDOORS-TO PROTECT ORNAMENTAL PLANTS:

To control aphids, exposed thrip, adult whiteflies, leafhoppers and two spotted mites on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyan, viburnum, suonymus, oaks, maples, locust, forsythia and althea, spray directly on insects. Hold container 18 inches from the infested plants. Spray in short spray bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, iris, zinnia, snapdragons, ageratums, chrysanthemums, nasturtiums, daisy, geraniums, marigoids, daphne, fatshedera, ivy, tobira, queen sago and broad teat acacta for control of aphids, exposed thrips, adult whiteflies and leafhoppers. Spray as directed above.

<u>USE WITH CAUTION</u> on young tender foliage. Avoid heavy wetting. Plants such as flowering quince, ash and camellias may be subject to burn under certain conditions.

For adult whiteflies, two spotted mites and mealybugs be sure to direct spray to contact these pests. Thoroughly cover the infested area. Use several repeat applications at two day intervals if necessary.

PROTECT HOUSE PLANTS:

Spray infested African violets, philodendron, geraniums, begonias, coleus for listed pests. Leave the sprayed area when done indoors, and ventilate area after treatment is completed.

NUISANCE PESTS - ON OR ENTERING BUILDINGS:

<u>Wasps:</u> Applications should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary. Spray directly on stray wasps that enter buildings.

Ants: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

Spiders: Direct short burst of spray on spiders and repeat as necessary.

Earwigs: Spray directly on earwigs in crevices and other hiding places on outdoor porches, party from and doorway entrances.

EPA Reg. No. 432-507

E - 2966 Y