## YOUR BRAND SBP-1382 / BIOALLETHRIN (0.20% + 0.125%)**AQUEOUS PRESSURIZED SPRAY FOR HOUSE** & GARDEN

\* KILLS: HOUSE FLIBS, MOSQUITOES, GNATS, WASPS, ROACHES, EARWIGS, TWO-SPOTTED MITES, SOWBUGS, SPIDERS, ANTS, BITING FLIES, APHIDS (PLANT LICE), LEAFHOPPERS, EXPOSED THRIPS, ADULT WHITEFLIES, AND FLEAS.

#### **ACTIVE INGREDIENTS:**

\*#Resmethrin[ 5-(phenylmethyl)-3-furanyl] methyl 2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate ...... 0.200%w/w d-trans Allethrin ...... 0.125%w/w OTHER INGREDIENTS: ...... 99.675%w/w 100.000%w/w

\*Cis/trans isomers ratio: max. 30%(±)cis and min. 70%(±)trans. ‡AgrEvo Environmental Health Inc.'s SBP-1382® brand of resmethrin insecticide. SBP-1382 is a registered trademark of AgrEvo Environmental Health, Inc.

EPA REG. NO. 432-507

EPA EST. NO.

### KEEP OUT OF REACH OF CHILDREN

Additional ionary Statements

NOTICE-WHEN USED IN BAKERIES, RESTAURANTS AND FOOD PROCESSING ESTABLISHMENTS, FOODS SHOULD BE REMOVED OR COVERED BEFORE TREATMENT. ALL FOOD PROCESSING SURFACES AND UTENSILS SHOULD BE COV-ERED OR REMOVED PRIOR TO TREATMENT, OR THOROUGHLY WASHED WITH AN EFFECTIVE CLEANING COM-POUND FOLLOWED BY A POTABLE WATER RINSE PRIOR TO USE. WHEN USING THIS PRODUCT IN THESE AREAS, APPLY ONLY WHEN THE FACILITY IS NOT IN OPERATION.

# ACCEPTET

3 SEP 1998

Under the Federal Insecticide. Fungicide, and Redenticide Act. as amended, for the posticide registered under 52-50 EPA Reg. No.

#### PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals CAUTION

Harmful if swallowed. Avoid contamination of feed and foodstuffs. Remove pets, birds, and cover fish aquariums before spraying. Do not use on edible crops.

#### **Environmental Hazards**

This pesticide is highly toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

Physical And Chemics Flazoros
Contents under pressure. Do let u enclus. Do not use or store
near heat or open flaming Exposer a to simple ratures above 130°F
may cause bursting. Never brown or tainer into fire or incinerator.

#### **DECTIONS FOR USE**

ation of Federal Law to use this product in a manner inconsisted with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Storage: Store in cool, locked storage area inaccessible to children, and away from heat, sparks or open flame. Container Disposal: Replace cap, wrap container in several layers of newspaper, and discard in trash.

SHAKE WELL BEFORE USING.

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

CONTROL HOUSEHOLD PESTS

FOR HOUSE FLIES, MOSQUITOES, AND GNATS: Close all doors and windows. Point nozzle upward. Direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

FOR CRAWLING INSECTS

ROACHES, EARWIGS, SOWBUGS, SPIDERS: Spray thoroughly into hiding places such as cracks, crevices, rapistaréas, openings around pipes and sinks, under refrigerators, beseboards, and storage areas. Spray directly on insects where possible. , , , , , ,

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

#### ANIMAL PESTS

TO KILL FLEAS, IN DOG AND CAT PREMISES: Thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of 2 seconds per linear foot at a distance of 12 to 18 inches. Also spray the entire inside surface of doghouses. Repeat as necessary, thoroughly-ensaving-floor

# SBP-1382°

many insects directly as possible. For best results, dogs and cats should also be treated with an appropriate insecticide.

**OUTDOORS** 

TO REDUCE ANNOYANCE FROM FLYING INSECTS: Reduce Mosquitoes, Gnats (Midges), House Flies, Biting Flies And Flying Moths (Millers) by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes, and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

**OUTDOORS** 

TO PROTECT ORNAMENTAL PLANTS: To control Aphids, Exposed Thrips, Adult Whiteflies, Leafhoppers And Two-spotted Mites on smaller rhododendrons, pines, spruce, hemlock, spirea, dogwood, palm, mimosa, toyan, viburnum, euonymus, oaks, maples, locust, forsythia and althea, spray directly on insects. Hold container 18 inches from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides covering foliage until slightly moist. Avoid heavy wetting. Repeat as necessary to obtain control.

Spray peonies, iris, zinnia, snapdragons, ageratums, chrysanthemums, nasturtiums, daisy, geraniums, marigolds, daphne, fatshedera, ivy, tobira, queen sago and broad leaf acacia for control of aphids, exposed thrips, adult whiteflies and leafhoppers. Spray as directed above.

USE WITH CAUTION: on young tender foliage. Avoid heavy wetting. Plants such as flowering quince, ash and camellias may be subject to burn under certain conditions.

For Adult Whiteflies, Two-spotted Mites And Mealybugs be sure to direct spray to control these pests. Thoroughly cover the infested area. Use several repeat applications at two day intervals if necessary.

PROTECT HOUSE PLANTS: Spray infested African violets, philodendron, geraniums, begonias, coleus for listed pests. Leave the sprayed area when done indoors, and ventilate area after treatment is completed.

NUISANCE PESTS-ON OR ENTERING BUILDINGS: WASPS: Applications should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary. Spray directly on stray wasps that enter buildings.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

SPIDERS: Direct short burst of spray on spiders and repeat as necessary.

EARWICS: Spray directly on earwigs in crevices and other hiding places, on outdoor porches, patio floor and doorway entrances.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645

> in case of Medical emergencies or health and safety inquiries or in case of fire, leaking or camaged containers, information may be obtained by calling 1-800-471-0660.

For Product Information Call Toll-Free: 1-800-331-2867

