

pm10

432-517

1-83

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Ms. Sharon M. Johnston  
Roussel Uclaf Corporation  
95 Chestnut Ridge Road  
Montvale, NJ 07645

POST OFFICE

JUN 21 1993

Dear Ms. Johnston:

**Subject:** Revised Ingredients Statement Incorporating "Related Compounds" into Inert Percentage  
SBP-1382<sup>®</sup> 0.35% Space and Residual Aqueous Pressurized Spray  
EPA Registration Number 432-517  
Submission dated May 18, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable subject to the comment listed below. Submit two (2) copies of the finished labeling prior to releasing the product for shipment.

Expand the STORAGE AND DISPOSAL statement as follows:

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.  
**STORAGE:** Store in cool, locked storage area inaccessible to children, and away from heat, sparks or open flame.  
**CONTAINER DISPOSAL:** Replace cap, wrap container in several layers of newspaper, and discard in trash.

Sincerely,



Rita Kumar  
Acting Product Manager 10  
Insecticide-Rodenticide Branch  
Registration Division

**CONCURRENCES**

SYMBOL							
SURNAME							
DATE							

2 of 3

BEST AVAILABLE

# SBP-1382 0.35% SPACE AND RESIDUAL AQUEOUS PRESSURIZED

### ACTIVE INGREDIENTS:

\*•Resmethrin[5-(phenylmethyl)-3-furanyl]methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate ..... 0.350%/w/w

INERT INGREDIENTS ..... 99.650%  
100.000%/w/w

\*Cis/trans isomers ratio max. 30% (z)cis and min. 70% (z)trans.  
©Roussel Uclaf's SBP-1382® brand of resmethrin insecticide.  
Roussel Uclaf Corp., 95 Chestnut Ridge Road, Montvale, N.J. 07645

**SHAKE WELL BEFORE USING.**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Avoid contamination of food, pets, birds and cover fish aquariums before spraying. Do not use in edible products areas (e.g., restaurants or other areas where food is prepared or processed). Do not use in serving areas while food is being served.

**ENVIRONMENTAL HAZARDS:** This pesticide is highly toxic to aquatic life. Do not apply directly to water, or to areas which drain to water, or to intertidal areas below the mean high water line. Treated sites may be hazardous to fish in adjacent water when disposing of equipment washwaters.

**PHYSICAL AND CHEMICAL HAZARDS:** Contains flammable vapors. Do not puncture or incinerate container. Do not use near open flame. Exposure to temperatures above 130°F may cause container to rupture. Throw container into fire or incinerator.

**KILLS:** House flies, mosquitoes, gnats, wasps, roaches, earwigs, grain mites, sowbugs, spiders, centipedes, ants, biting flies, aphids (plant lice), leafhoppers, exposed termites, adult whiteflies, plant bugs, Japanese beetles, spittlebugs.

**FOR USE IN HOMES, MOTELS, INSTITUTIONS, RESTAURANTS, AND FOOD PROCESSING PLANTS.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with the label directions.

EPA Reg. No. 432 517

EPA Est.: See Lot No. F 14370-TX-1  
A 42545

P-12/12/89

NET WT.

LOT NO.:

**RECEIVED**  
**LABORATORY**  
**IN EPA LABORATORY**  
JUN 21 1993  
P 655-GA-1  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 432-517

**ROUSSEL UCLAF**  
Roussel Uclaf Corporation  
95 Chestnut Ridge Road  
Montvale, New Jersey 07645

**STORAGE:** Keep in locked storage area.  
**CONTAINER DISPOSAL:** Replace cap and recycle in trash.

NOTICE: Buyer assumes all responsibility for safety and use.

3  
2

## SBP-1382® 0.35% SPACE AND RESIDUAL AQUEOUS PRESSURIZED SPRAY

EPA Reg. No.432-517

DIRECTIONS FOR USE

**BEST AVAILABLE COPY**

**DOUBLE ACTION INSECT KILLER:** This is a dual purpose formulation that can be used as a contact spray against flying and crawling insects and as a residual insecticide providing up to 3 to 4 weeks of control of roaches.

### DIRECTIONS FOR USE

Shake well before each use. Remove protective cap. Aim spray opening away from person. Push button to spray.

**HOUSE FLIES, MOSQUITOES AND GNATS: INDOORS:** Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed.

**TO REDUCE ANNOYANCE FROM FLYING INSECTS: OUTDOORS:** Reduce mosquitoes, gnats, midges, house flies, biting flies and flying moths (millers) by spraying their resting areas. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic feet of space while moving away from the treated areas. To reduce annoyance from these flying insects, treat open porches and patios in a similar manner. Spray only when the air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

**ROACHES, EARWIGS, GRAIN MITES, SOWBUGS, SPIDERS AND CENTIPEDES: CONTACT SPRAY:** Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, and storage areas. Spray directly on insects where possible. Repeat as necessary.

**ROACHES, EARWIGS, GRAIN MITES, SOWBUGS, SPIDERS AND CENTIPEDES: RESIDUAL SPRAY:** Spray baseboards, cracks and crevices, around door and

window frames, and other hiding places such as closets, in back of and under appliances, sinks and cupboards in the kitchen and bathroom. In basements, spray hiding places in cracks and crevices, lower walls and dark places. Spray at the rate of 3 to 4 seconds per linear foot at a distance of 12 to 14 inches from the surface. Repeat as necessary.

**WASPS:** Applications should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, contacting as many insects as possible. Repeat as necessary.

**ANTS:** Apply to ant trails and around door sills, window frames and other places where ants may enter.

**ORNAMENTAL PLANT PESTS:** Spray aphids, exposed thrips, adult whiteflies, leafhoppers, plant bugs, spittle bugs, Japanese beetles, mimosa webworm larvae, and oakworm larvae. Direct spray on insects by holding the container 12 to 18 inches from the infested plants. Spray in short bursts of one to three seconds or spray at the rate of one linear foot of growth per second. Direct spray to all sides, covering foliage until slightly moist. Avoid heavy wetting. Repeat applications as necessary.

**USE WITH CAUTION** on the foliage of Flowering Quince, Juniper, Cypress, Mulberry, Ash, Cotoneaster, Rose, Camellia, Azalea, Pyracantha and Caenothus, as certain varieties have shown some leaf browning.

When infested, spray smaller evergreens, deciduous trees and shrubs as rhododendrons, pine, spruces, hemlock, yews, arborvitae, junipers, holly, podocarpus, boxwood, laurel, lilac, spirea, dogwood, palm tree, mimosa, toyon, viburnum, oaks, maples, locust, forsythia, althea and roses for control of aphids, and other pests listed above. Also spray infested flowering plants as peonies, iris, zinnia, snapdragons, ageratums, chrysanthemums, nasturtium, daisy, geraniums, marigolds, daphne, fatsihedera, ivy, tobiira, queen sago, broadleaf acacia for the same listed pests.