

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)**

CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid contamination of feed and foodstuffs. Remove pets, birds, and cover fish aquariums before space spraying. Do not use on edible crops. Do not remain in treated areas. Ventilate the area when treatment is complete.

FIRST AID

In case of contact, immediately flush eyes, and skin with plenty of water for at least 15 minutes. Get medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician at once.

ENVIRONMENTAL HAZARDS

Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place. A container size of 30 gals. or more may be resealed and offered for reconditioning.

GENERAL: Consult federal, state or local disposal authorities for approved alternate procedures.

Do not use in edible product areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. In the home, cover all food handling surfaces and cover or remove all food and cooking utensils.

("Directions For Use" continued on Right Panel.)

**SBP-1382® 0.25%
TRANSPARENT
EMULSION*
SPRAY**

A SYNTHETIC PYRETHROID

ACTIVE INGREDIENTS:

*resmethrin([5-(phenylmethyl)-3-furanyl)methyl 2,2-dimethyl 3-(2-methyl-1-propenyl)cyclopropanecarboxylate)

Related compounds

INERT INGREDIENTS:

0.250%w/w

0.034%

99.716%

100.000%w/w

*Cis/trans isomers ratio max 30% (±)cis and min 70% (±)trans.

†U.S. Patent Nos. 3,465,007 and 3,542,928.

**U.S. Patent No. 3,683,078.

•Penick's SBP-1382® brand of resmethrin insecticide.

Penick Corp. 1050 Wall Street West, Lyndhurst, N.J. 07071

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

EPA Reg. No. 432-548

EPA Est. 432-NJ-1

NET WEIGHT

POUNDS

NET CONTENTS

GALLONS

PENICK Penick Corporation

1050 Wall Street West, Lyndhurst NJ 07071

ACCEPTED

SEP 24 1980

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 432-548

DIRECTIONS FOR USE (continued)

For use in non-food areas of food handling establishments, in and around premises commercial, industrial, office, residential, public, institutional and storage buildings, hotels, motels, homes, preschools, nurseries, day camps, seed houses, packaging and processing plants, milk handling areas of milk plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in campers, mobile homes, buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

Do not use on electrical equipment or on surfaces that can be harmed or stained by water. **INDOORS**

FLYING INSECTS: To kill houseflies, deer flies, stable flies, gnats, fruit flies, flying moths (millers). Apply as a space spray. Mix well before using. Fill spray tank with undiluted SBP-1382® 0.25% Transparent Emulsion Spray. Adjust pressure and nozzle to deliver a fine mist spray. Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract these insects. Keep the spray nozzle at all times at least four feet from fabrics to avoid wetting or staining. Fill the room with mist at the dosage of one fluid ounce per 1000 cubic feet of space. Then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed and before reentry.

To kill wasps, hornets, yellow jackets, bees - Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Repeat as necessary.

CRAWLING INSECTS: To kill fleas, cockroaches, waterbugs, crickets, silverfish, earwigs, spiders, sowbugs, centipedes, millipedes, grain mites and grain insects (adults and accessible stages of confused flour beetles, red flour beetles, sawtoothed grain beetles, shiny spider beetles, carpet beetles, grain weevils, Angoumois grain moths, Indian meal moths) - Apply as a contact spray. Mix well before using. Fill spray tank with undiluted SBP-1382® 0.25% Transparent Emulsion Spray. Keep spray nozzle about 8 to 12 inches from surface to be treated. Adjust pressure and nozzle to deliver a coarse spray with particle size larger than that of the average aerosol. Spray thoroughly into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards, storage areas and other insect harborage areas. Spray directly on insects where possible. Repeat as necessary.

*Grain Insects - For control of the insect pests listed above, remove infested grain, cereal or flour and destroy. Avoid contamination of other food and foodstuffs in the area by removing before spraying. Spray directly on exposed stages and into cracks and crevices where grain insects may be hiding. In homes, dispose of all infested products. The shelves and drawers should be cleaned and thoroughly sprayed so as to contact all pests, and covered with shell paper after allowing shelves to dry. Any utensils that are contacted with spray should be washed before using. Repeat as necessary.

TO KILL SPIDERS: Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

TO KILL ANTS: Spray doors, around window frames, ant trails and hills.

TO KILL BEDBUGS: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

TO KILL FLEAS ON DOGS, CATS & PREMISES: Apply directly to animals along back, under forelegs, and on other areas except scrotum and anus. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not spray directly in eyes or mouth of animal. Repeat within 24 hours of first treatment, if necessary. To reduce reinestation, thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches. Also spray the entire inside surface of dog houses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible.

OUTDOORS

TO KILL HOUSEFLIES, MOSQUITOES, GNATS (Midges), DEER FLIES, STABLE FLIES, FLYING MOTHS (Millers), FRUIT FLIES - Spray resting areas to reduce annoyance from these pests. Follow precautions listed above. Hold container 18 to 24 inches from shrubs, bushes, and grassy areas. Spray these areas with a slow sweeping motion at the rate of from 1 to 3 seconds per cubic yard of space, while moving away from the treated areas. Treat open porches and patios in a similar manner. Spray only when air is still. Allow spray to dissipate before occupying the treated area. Repeat as necessary.

WASPS, HORNETS, YELLOW JACKETS, BEES - Same as directed above for indoor use. Application should be made late in the evening when insects are at rest. Aim spray at nest openings in ground, bushes, and in cracks and crevices which may harbor nests, saturating nest openings and contacting as many insects as possible. Repeat as necessary.

ANTS AND FLEAS - Same as directed above for indoor use.

CRICKETS, SILVERFISH, EARWIGS, SPIDERS, SOWBUGS, CENTIPEDES AND MILLIPEDES - Same as directed above for indoor use.

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effect of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse as such, or in combination with other materials.

432-548

09/24/1980

11