

Reg # 432-562

PM-18

131

1% ROTENONE DUST**CONTROLS**

Beetles (Asparagus, Mexican Bean, Bean Leaf, Spotted &
Striped Cucumber, Flea & Colorado Potato)

Imported Cabbage Worm & Cabbage Looper

Fleas on Dogs and Cats

ACTIVE INGREDIENTS:

Rotenone..... 1.0%

Other Associated Resins 2.0%

INERT INGREDIENTS: 97.0%

100.0%

KEEP OUT OF REACH OF CHILDREN**CAUTION**

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

EPA Est.: See Lot No. For Correlation

EPA Reg. No. 432-562

A 42545-MO-1

F 16370-TX-1

P 655-GA-1

P- 3/21/90

NET CONTENTS:**ACCEPTED**

AUG 19 1994

Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 432-562

ROUSSEL UCLAF

Roussel Uclaf Corporation
95 Chestnut Ridge Road
Montvale, New Jersey 07645

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through skin. Avoid inhalation of dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, get medical attention. If in eyes, flush with plenty of water. If irritation develops or persists, get medical attention. If on skin, wash immediately with soap and water.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwater. Do not feed treated foliage to livestock. Do not apply rotenone in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until sprays have dried or dust has settled.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Dust at the rate of 10 to 20 pounds per acre or 2 oz. to each 50 feet of row in the garden. On flowers and shrubs dust the entire plant thoroughly. Repeat applications as necessary for effective control. When insect infestation is heavy applications are usually necessary at weekly intervals. To control Mexican Bean Beetles, direct the material to the underside of the leaves. NOTE: This product can be applied to vegetables up to within 1 day of harvest.

GARDEN AND TRUCK CROPS

ASPARAGUS: Asparagus Beetle.

BEANS: Mexican Bean Beetle, Bean Leaf Beetle, Spotted and Striped Cucumber Beetles.

BEETS: Flea Beetle.

CABBAGE, OTHER CORD CROPS: Imported Cabbage Worm, Cabbage Looper.

CUCUMBER, SQUASH, MELONS: Spotted and Striped Cucumber Beetles.

EGG PLANT: Colorado Potato Beetle, Flea Beetle.

LETTUCE: Loopers.

PEAS: Pea Aphid.

POTATO: Colorado Potato Beetle, Flea Beetle, Aphid.

RADISH: Flea Beetle.

SPINACH: Flea Beetle.

TOMATO: Colorado Potato Beetle, Aphid, Flea Beetle.

TURNIPS: Flea Beetle, Loopers.

FLOWERS AND SHRUBS

CABBAGE LOOPER & IMPORTED CABBAGE WORM: On Carnation, Chrysanthemum, Geranium and Nasturtium.

ANIMALS

DOGS AND CATS: Do not use on puppies or kittens less than 2 weeks old.

Fleas: Dust thoroughly into hair and rub in with hands. Avoid getting into animal's eyes, nose and mouth. Also thoroughly treat the sleeping quarters, including bedding.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

BEST AVAILABLE COPY

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.