



PENICK CORPORATION

CHLORPYRIFOS INSECTICIDE TRANSPARENT EMULSION** SPRAY 0.5%

OFFERING RESIDUAL KILL OF COCKROACHES FOR UP TO 4 WEKKS

For residual and contact control of cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, centipedes, millipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bed bugs, houseflies, mosquitoes, gnats, flying moths, confused flour beetles, sawtoothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths, clothes moths.

CONTAINS DURSBAN[†] INSECTICIDE

ACTIVE INGREDIENT:

*Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-phosphorothioate)]

INERT INGREDIENTS:

ACCEPTED	
432-624	
SEP 14 1982	
Under the Federal Insecticide, Fungicide, and Rodenticide Act, registered under EPA Reg. No. 432-624	
0.5% v/w	99.5%
100.0% w/w	

BEST DOCUMENT AVAILABLE

*U.S. Patent No. 3,244,586. Dursban is a Trademark of Dow Chemical Co.
**U.S. Patent No. 3,683,078.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
TRANSLATION: (To the user: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

EPA Reg. No. 432-624

EPA Est. 432-NJ-1

NET CONTENTS:

In food handling establishments, can be used in non-food areas as a contact or residual spray. In food areas, application is limited to spot and/or crack and crevice treatment.

Penick Corporation
1050 Wall Street West
Lyndhurst, New Jersey 07071 U.S.A.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash hands after using.

In food handling areas, application is limited to crack and crevice and/or spot treatment. Do not apply directly to food or feedstuffs. In commercial food handling facilities, cover or remove any food or food processing equipment during application. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Get immediate medical attention. If in eyes or on skin: Flush with plenty of water for at least 15 minutes. For eyes, get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition appear, atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

PHYSICAL/CHEMICAL HAZARDS

Do not apply this water-based product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be harmed or stained by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Do not store this water-based formulation at temperatures below 32°F.

BEST DOCUMENT AVAILABLE

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures.

CONTAINER DISPOSAL:

For containers one gallon and under: Do not use empty container. Wrap container and place in trash collection.

For containers over one gallon: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and Local procedures.

FOR USE IN NON-FOOD AREAS INDOORS AND OUTSIDE PREMISES OF HOTELS, MOTELS, HOMES, PRESCHOOLS, APARTMENT BUILDINGS, COMMERCIAL, OFFICE, INSTITUTIONAL AND STORAGE BUILDINGS, RECREATIONAL BUILDINGS, BOATS, SHIP CABINS, AND HOSPITALS. IN HOSPITALS, REMOVE PATIENTS PRIOR TO TREATMENT. VENTILATE ROOMS FOR TWO HOURS BEFORE RETURNING PATIENTS.

THIS PRODUCT CAN BE USED IN NON-FOOD AREAS SUCH AS CORRIDORS, OFFICES, FOYERS, LAVATORIES, GARAGES, UTILITY ROOMS, STORAGE ROOMS AND BASEMENTS, OF PACKAGING AND PROCESSING PLANTS, MILK HANDLING AREAS OF MILK PLANTS, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, SUPERMARKETS, RESTAURANTS, BOTTLING PLANTS, CANNERIES, FLOUR AND FEED MILLS, AND BAKERIES.

IN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, APPLICATION IS LIMITED TO SPOT AND/OR CRACK AND CREVICE TREATMENT.

ALSO FOR USE IN AND AROUND CAMPERS, MOBILE HOMES, BUSES AND AUTOMOBILES. THIS PRODUCT CAN BE USED IN CARGO PLANES, TRUCKS, BOXCARS AND SHIP CARGO HOLDS ONLY IF THE VEHICLES ARE NOT USED FOR TRANSPORT OF FOOD OR FOOD PRODUCTS AT ANY TIME.

ALSO, FOR USE IN DOG KENNELS, AND HORSE STABLES. DO NOT SPRAY ON HORSES AND DOGS. DO NOT SPRAY DOG AND HORSE FOOD OR FEED ITEMS.

NOTE TO PEST CONTROL OPERATORS: One gallon (3.79 L) of spray is sufficient to treat 1500 linear feet (457 m) with a 6 inch (15.2 cm) spray band.

For contact and residual sprays, use this product without diluting. For repeat applications, dilute with equal parts of water.

CONTACT SPRAY-KILLS AND REDUCES INFESTATIONS: Cockroaches (waterbugs), crickets, silverfish (firebrats), clover mites, centipedes, millipedes, earwigs, sowbugs (pillbugs), spiders, brown dog ticks, lone star ticks, ants, fire ants, carpet beetles, fleas, boxelder bugs, scorpions, spider beetles, bed bugs, confused flour beetles, sawtoothed grain beetles, shiny spider beetles, grain mites, red flour beetles, angoumois grain moths, grain weevils, Indian meal moths, Mediterranean flour moths, lesser grain borers, cigarette beetles, drug store beetles, chocolate moths. Use as a spot treatment. When possible, spray directly on pests to reduce infestations. Apply as a coarse spray or with a paint brush into hiding places such as cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators and other appliances, baseboards and storage areas. Apply spray until area is wet, but not to the point of run-off. Repeat as necessary.

DIRECTIONS FOR USE (continued)

432 624

ANTS (Including Fire Ants): To kill ants, spray trails, nests and points of entry within buildings and homes. spray directly on ants where possible. Repeat as necessary.

FLEAS, BROWN DOG AND LONE STAR TICKS: Thoroughly apply as a spot treatment to infested areas such as pets' beds and resting quarters; nearby cracks and crevices; along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean fresh bedding after treatment of pet areas. DO NOT TREAT PETS WITH THIS PRODUCT. Treat pets with a product registered for animal application, such as SBP-1382@/Bioallethrin Aqueous Pressurized Spray, EPA Reg. No. 432-482, to control the source of flea and tick infestation.

CARPET BEETLES AND CLOTHES MOTHS: Do not spray clothing with this product. Do not allow clothing to contact areas where this product has been used. Spray edges of carpeting and under carpeting, rugs, and closets. Make localized application to the floor and baseboards. Spray directly into cracks, crevices, closets, and infested areas of shelving and/or wherever these insects are seen or suspected. Repeat as necessary.

RESIDUAL SPRAY: AIDS IN THE CONTROL OF COCKROACHES (WATERBUGS), CRICKETS, BOXELDER BUGS, SCORPIONS, SPIDER BEETLES, SILVERFISH (FIREBRATS), CENTIPEDES, MILLIPEDES, EARWIGS, SOWBUGS (PILLBUGS). Kills up to 4 weeks. Apply on baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back of and under appliances, sinks and cupboards in kitchens and bathrooms. In basements, apply in hiding places, in cracks and crevices, lower walls and dark places. Spray surfaces until wet, but not to the point of run-off.

PESTS OUT-OF-DOORS

PESTS OUT-OF-DOORS: DO NOT TREAT OR ALLOW THE SPRAY TO CONTACT PLANTS (LAWNS, FLOWERING PLANTS, SHRUBS, TREES, EVERGREENS, ETC.) BECAUSE OF POSSIBLE INJURY.

PESTS ON OUTSIDE SURFACES OF BUILDINGS: Use this product to control ants, clover mites, crickets, earwigs, millipedes and sowbugs (pillbugs) on the outside surfaces of buildings. To help prevent infestation of buildings treat the building foundation to a height of 2 to 3 feet (0.5 to 1 m), where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of buildings, including porches, screens, window frames, eaves, patios, garages, refuse dumps, and other areas where these pests congregate or have been seen. Repeat treatment as needed to maintain effectiveness.

HOUSEFLIES, MOSQUITOES, GNATS, FLYING MOTHS, OUTDOORS: For outdoor use only as aid in reducing annoyance from these insects. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, and other places where these insects might or congregate. Repeat as necessary.

BEST DOCUMENT AVAILABLE

1132-624

ANTS, BROWN DOG TICKS, CRICKETS, FLEAS, AND EARWIGS OUTDOORS: For treatment of localized infestations of these insects in outdoor areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application as infestations warrant and as reinfestations occur.

TO CONTROL FIRE ANTS: Dilute 1.5 pints (0.71 L) of this product in 1 gallon (3.79 L) of water. Apply 1 gallon (3.79 L) of solution as a gentle rain to each fire ant mound using a sprinkler can. Thoroughly wet the mound and surrounding area to a 4 foot diameter (1.16 m). For best results, apply in cool weather, 65-80°F (18.5°C-27°C), or in early morning or late evening hours. Treat new mounds as they appear. Do not disturb the mound prior to treatment. If possible, treat all mounds in the vicinity, and treat all colonies which may not as yet have constructed a mound.

Larger quantities of solution can be prepared for multiple mound treatments. Equipment to deliver the diluted product as a gentle rain at the rate of 1 gallon (3.79 L) per mound is still required.

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.