

SBP-1382® PRESSURIZED SPRAY INSECTICIDE 0.15% FORMULA III For Wasps and Hornets

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. If swallowed, do not induce vomiting. Call a physician immediately.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is underway. After spraying in meat packing plants, bakeries and other food processing plants, thoroughly wash all equipment, benches, shelving, etc., where exposed food will be handled, and rinse with potable water. In the home, all food processing surfaces and utensils should be covered during treatment, or thoroughly washed before use. Remove pets, birds and cover fish aquariums before spraying.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marches, and potholes). Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130°F (54.5°C) may cause bursting. Never throw container into fire or incinerator.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For use in and around hotels, motels, homes, greenhouses, and lathhouses, food handling establishments, packaging and processing plants, food houses, commercial, industrial, office, institutional and storage buildings, milk handling areas of milk, cheese and ice cream plants, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mills, bakeries, boats, ship cabins and hospitals. In hospitals, patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boxcars and ship cargo holds. Also for use in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

STORAGE: Keep in locked storage area.

CONTAINER DISPOSAL: Replace cap and discard container in trash.

**A SYNTHETIC PYRETHROID WITH LONG SPRAY PATTERN
UP TO 10 FEET KILLS WASPS, HORNETS, YELLOW JACKETS,
SPIDERS AND TENT CATERPILLARS**

Authorized by USDA for use in edible product areas of official establishments operating under the Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs.

Dielectric Strength-Non-conducting up to 26,500 volts

ACTIVE INGREDIENTS:

*Resmethrin [5-(phenylmethyl)-3-furanyl]methyl 2,2-dimethyl-3-(2-methyl-1-propenyl)cyclopropanecarboxylate 0.150%w/w
Petroleum Distillate 11.05%

INERT INGREDIENTS: *Centauris Petroleum Distillates* 88.210%
100.000%w/w

*Cis/trans isomers (ratio: max. 30% (1)cis and min. 70% (1)trans.
*Roussel Bio's SBP-1382® Brand of resmethrin insecticide.
Roussel Bio Corp., 400 Sylvan Avenue, Englewood Cliffs, N.J. 07632

**KEEP OUT OF REACH OF CHILDREN
CAUTION**
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

EPA Est.: See Lot No. For Correlation
EPA Reg. No. 432-640 A 42545-MO-1 P-12/08/89
F 10370-TX-1 P 655-GA-1
NET CONTENTS:

ROUSSEL BIO
CORPORATION
P.O. Box 1077 400 Sylvan Avenue
Englewood Cliffs New Jersey 07632

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticides and Fungicides for Pesticide 2
Handbook, Page 2.12, under the Pesticide in Service.

DIRECTIONS FOR USE (Continued)

Remove protective cap. Aim spray opening away from person. Push button to spray.

INDOORS: To kill wasps, hornets, yellow jackets: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places, particularly under rafters, contacting as many insects as possible. Saturate areas in and around nest openings. Repeat as necessary.

TO KILL SPIDERS: Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

OUTDOORS: To kill wasps, hornets, yellow jackets: Application should be made in late evening when insects are at rest. Aim spray at nest openings in trees, bushes, under eaves, chimneys, cracks and crevices to saturate in and around nest openings. Contact as many insects as possible. Repeat as necessary.

TO KILL SPIDERS: Same as directed above for indoor application.

TO KILL TENT CATERPILLARS: Application should be made when caterpillars are young and tents are first noticed. For best results apply in the late afternoon or evening when caterpillars have returned to their tents. Aim spray directly at tents, and totally saturate on all sides. Repeat as necessary.

WARRANTY STATEMENT

Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.