

FOLIAFUME® INSECTICIDE
with ROTENONE/PYRETHRINS
EMULSIFIABLE CONCENTRATE 1.1% + 0.8%

FORMULA II
EMULSIFIABLE PYRETHRUM-ROTENONE PLANT & ANIMAL INSECTICIDE
Contains 0.08 lb/gal (9.9 g/L) of Rotenone & 0.06 lb/gal (7.2 g/L)
of Pyrethrins

ACTIVE INGREDIENTS:

Rotenone	1.1% w/w
Other Cube Extractives	2.2%
Pyrethrins	0.8%
Petroleum Distillate	3.2%
Aromatic Petroleum Solvent	84.7%
INERT INGREDIENTS:	8.0%
	100.0% w/w

*Penick Corporation Trademark

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

TRANSLATION: (To the user: If you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN
WARNING

SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

EPA Reg. No. 432-684

EPA Est. 432-NJ-1

E-30374-P

NET WT.

lbs (

Kg)

PENICK Penick Corporation

1050 Wall Street West Lyndhurst, New Jersey 07071 U.S.A.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

May be fatal if swallowed. Harmful if absorbed through skin. Causes skin irritation. May cause substantial but temporary eye injury. Wear safety glasses. Do not get in eyes, on skin, or on clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call physician or Poison Control Center. Drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Do not induce vomiting. This product contains aromatic petroleum solvent. Aspiration may be a hazard.

If in eyes or on skin: Flush with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Flash Point minimum 140°F (60°C).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State, or Local procedures.

CONTAINER DISPOSAL: For containers one gallon and under: Do not use empty container. Wrap container and place in trash collection.

For containers over one gallon: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State or local procedures.

For use on ornamental and flowering plants such as Poplar, Linden, Dogwood, Willow, Birch, Boxelder, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Spirea, Rose, Lilac, Zinnia, Aster, Petunia, Geranium, Dahlia, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation, Chrysanthemum, and Rose, mix this product with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Do not apply this product within one day of collection. Repeat every 5-10 days or as needed.

DIRECTIONS FOR USE (Continued on Reverse Side)

DIRECTIONS FOR USE (Continued)

EPA Reg. No. 432-684

PESTS	VOLUME OF PRODUCT/GALLON OF WATER (3.79L)
Aphids, Leafhoppers, Exposed Thrips	1 teaspoon (5 ml)
Oblique-banded Leaf Rollers, Sawfly larvae, Boxelder bugs, Japanese beetles, Elm leaf beetle larvae, Mites, Lygus bugs, Lace bugs, Greenhouse Whiteflies, Stink bugs, Garden Fleahoppers, Greenhouse Leafhoppers	2 teaspoons (10 ml)
Cucumber beetles, Rose chafers, Ants, Aster beetles, Flea beetles	4 teaspoons (20 ml)

For use on vegetable plants such as peas, bean, beet, cucumber, squash, muskmelon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot and corn: Mix with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed.

PESTS	VOLUME OF PRODUCT/GALLON OF WATER (3.79L)
Aphids, Leafhoppers, Exposed Thrips	1 teaspoon (5 ml)
Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetles, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Oblique-banded Leaf Rollers, Stink Bugs, Celery Leafhoppers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites	2 teaspoons (10 ml)
Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers- 3 to 4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals.	4 teaspoons (20 ml)

For use on Fruits and Berries such as apple, pear, peach, plum, cherry, grape, currant, gooseberry, raspberry, blackberry, strawberry and blueberry, consult the following chart. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed.

PESTS	VOLUME OF PRODUCT/GALLON OF WATER (3.79 L)
Aphids, Leafhoppers, Exposed Thrips	1 teaspoon (5 ml)
Imported Currant Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms (adult), Red-necked Borers, Japanese beetles, and Mites	2 teaspoons (10 ml)
Strawberry Leafrollers and Rose Chafers	4 teaspoons (20 ml)

TO KILL LICE, FLEAS AND TICKS ON DOGS, CATS AND PREMISES: For lice and fleas, dilute at the rate of 1 tablespoon (15 ml) per gallon (3.79 L). For ticks, dilute at the rate of 4 tablespoons (60 ml) per gallon (3.79 L). Apply directly to animals as a spray or wash. Avoid contact with eyes, nose or mouth of animal. To prevent reinfestation, thoroughly spray the animals' living quarters.

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.