

UltraTEC® INSECTICIDE with ROTENONE/PYRETHRINS TRANSPARENT EMULSION** DILUTABLE CONCENTRATE 2.2% + 1.6%

Contains 0.18 lb/gal of Rotenone, 0.36 lb/gal of Other Cube Extractives, and 0.13 lb/gal of Pyrethrins.
For Control of Vegetable, Ornamental and Flowering Plant Pests, Pests of Fruits and Berries,
and Lice, Fleas and Ticks on Dogs, Cats and Premises.

TO BE APPLIED ONLY BY OR UNDER THE SUPERVISION OF PEST CONTROL OPERATORS, SERVICE PERSONNEL,
SUCH AS COMMERCIAL NURSEYRIMEN, LICENSED VETERINARIANS AND AGRICULTURAL SPECIALISTS
OR OTHER TRAINED PERSONNEL RESPONSIBLE FOR INSECT CONTROL PROGRAMS.

ACCEPTED
JAN 16 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act,
this pesticide is registered under EPA Reg. No. 432-697

ACTIVE INGREDIENTS:

Rotenone..... 2.2% w/w
Other Cube Extractives 4.4%
Pyrethrins 1.6%

*†INERT INGREDIENTS: 91.8%

100.0%w/w

*Contains Petroleum Distillate.

†Contains Methanol.

**U.S. Patent No. 3,683,078.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.
(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

**KEEP OUT OF REACH OF CHILDREN
DANGER
POISON**



DANGER: Methanol may cause blindness. Harmful or fatal if swallowed. Vapor harmful. Avoid breathing mist or vapors. Avoid contact with skin. In case of eye contact: Flush with plenty of water and get medical attention. See side panel for additional precautionary statements.

LOT NO.:

EPA Reg. No. 432-697

EPA Est. 432-NJ-1

P-85-287

NET CONTENTS:

PENICK

Penick Corporation

150 Wall Street West, Lyndhurst, N.J. 07071 U.S.A.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS DANGER

Causes irreversible eye damage. Harmful or fatal if swallowed. Avoid breathing vapors. Do not get in eyes, on skin or on clothing. Wear goggles or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Remove pets and cover fish aquariums before spraying.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a Physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If in eyes: Flush with plenty of water. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

Contains 6.4% Petroleum Distillate.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

FLAMMABLE. Do not use or store near heat or open flame. Do not use in the vicinity of pilot lights. Flash point minimum of 80°F (15.5°C).

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: This material may stratify upon standing, even at room temperature. Thoroughly mix before each use. Do not expose to temperatures below 40°F (4.5°C). Allow to stand until product reaches room temperature and thoroughly mix before using. **DO NOT USE OPEN FLAME.**

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

DIRECTIONS FOR USE (Continued on Reverse Side)

PLANT PESTS

For use on ORNAMENTAL and FLOWERING PLANTS such as Poplar, Linden, Dogwood, Willow, Birch, Boxelder, Maple, Oak, Elm, Pine, Spruce, Cedar, Yew, Spirea, Lilac, Zinnia, Aster, Petunia, Geranium, Dahila, Gladiolus, Phlox, Sweet Pea, Marigold, Carnation, Chrysanthemum, and Rose, mix this product with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant. Do not apply this product within one day of collection. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS	Volume of product/gallon of water (3.79 L)
Aphids, leafhoppers, exposed thrips	1/2 teaspoon (2.5 ml)
Oblique-banded leaf rollers, Sawfly larvae, Boxelder bugs, Japanese beetles, Elm leaf beetle larvae, Mites, Lygus bugs, Lace bugs, Greenhouse Whiteflies, Stink bugs, Garden fleahoppers, Greenhouse leafhoppers	1 teaspoon (5 ml)
Cucumber beetles, Rose chafers, ants, Aster beetles, Flea beetles	2 teaspoons (10 ml)

For use on VEGETABLE PLANTS such as peas, bean, beet, cucumber, squash, muskmelon, potato, tomato, cabbage, spinach, lettuce, celery, asparagus, pepper, eggplant, onion, carrot, and corn. Mix with water as described in the following chart. Use hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed. Keep diluted spray away from light.

PESTS	Volume of product/gallon of water (3.79 L)
Aphids, leafhoppers, exposed thrips	1/2 teaspoon (2.5 ml)
Cabbage Worms, Bean Beetles, Beet Webworm, Asparagus Beetles, Japanese Beetle, Lygus Bugs, White Fly, Garden Fleahopper, Harlequin Bug, Oblique-banded Leaf Roller, Stink Bugs, Celery Leafhopper, Melon Worm, Tortoise Beetle, Garden Webworm, Sod Webworm, Alfalfa Webworm and Mites	1 teaspoon (5 ml)
Cucumber beetles, Flea Beetles, Vegetable Weevils, Squash Bug (nymphs), Squash Vine Borer, European Corn Borer, 3-4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals.	2 teaspoons (10 ml)

For use on FRUITS and BERRIES such as apple, pear, peach, plum, cherry, grape, currant, gooseberry, raspberry, blackberry, strawberry, and blueberry, consult the following chart. Use sufficient spray mix to give complete coverage of the plants. Do not apply this product within one day of harvest. Repeat every 5-10 days or as needed.

PESTS	Volume of product/gallon of water (3.79 L)
Aphids, leafhoppers, exposed thrips	1/2 teaspoon (2.5 ml)
Imported currant worm, gooseberry fruit worm, raspberry fruit worm, blueberry maggot, cherry fruit fly, strawberry root worm (adult), red-necked borer, Japanese beetle, and mites.	1 teaspoon (5 ml)
Strawberry leafroller and rose chafer	2 teaspoons (10 ml)

ANIMAL PESTS

TO KILL FLEAS, TICKS AND LICE ON DOGS, CATS AND PREMISES: (Keep diluted spray away from light).

DO NOT USE ON NURSING PUPPIES OR NURSING KITTENS.

ON DOGS: DO NOT USE MORE THAN ½ FLUID OUNCE (1 Tbsp.)(15 ml) PER GALLON (3.79 L) ON PUPPIES.

For fleas and lice, dilute UltraTEC® Insecticide at the rate of ½ fluid ounce (1 Tbsp)(15 ml) per 1 gallon (3.79 L) of water. [For young animals, use ¼ fluid ounce (½ Tbsp)(7.5 ml) per 1 gallon (3.79 L) of water.] Apply directly to animal as a spray, sponge-on or a wash. Dip until animal's skin is wet. Do not rinse.

For ticks, dilute UltraTEC® Insecticide at the rate of 1 fluid ounce (2Tbsp)(30 ml) per 1 gallon (3.79 L) of water. [For young animals use ½ fluid ounce (1 Tbsp)(15 ml) per 1 gallon (3.79 L) of water]. Apply directly to animal as a spray, sponge-on or a wash. Dip until animal's skin is wet. Do not rinse.

Avoid contact with eyes, nose or mouth of animal. To prevent reinfestation, thoroughly spray the animal's living quarters with UltraTEC® Insecticide.

ON CATS: DO NOT USE ON YOUNG KITTENS. ON ADULT ANIMALS, DO NOT USE MORE THAN ½ FLUID OUNCE (1 Tbsp)(15 ml) PER GALLON (3.79 L).

For fleas, ticks, and lice, dilute UltraTEC® Insecticide at the rate of ½ fluid ounce (1 Tbsp)(15 ml) per gallon (3.79 L) of water. Apply directly to animal as a spray, sponge-on or a wash. Dip until animal's skin is wet. Do not rinse.

Avoid contact with eyes, nose or mouth of animal. To prevent reinfestation, thoroughly spray the animal's living quarters with UltraTEC® Insecticide at the rate of 1 fluid ounce (2 Tbsp)(30 ml) per gallon (3.79 L) of water.

Thoroughly spray the animal's bedding and other resting areas until the surface is slightly moist. Treat these areas at the rate of two seconds per linear foot at a distance of 12 to 18 inches (30 to 46 cm). Also spray entire inside surface of dog houses. Repeat as necessary, thoroughly spraying floors, cracks, crevices and bedding. Hit as many insects directly as possible.

WARRANTY STATEMENT

Our recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage resulting from its misuse as such, or in combination with other materials.

