

PROPOXUR/PYRETHRINS/PIPERONYL BUTOXIDE TRANSPARENT EMULSION** PRESSURIZED SPRAY 0.5%+0.05%+0.5%

An Insecticide Offering Good Flushing Activity, Quick Knockdown and Residual Kill of Cockroaches for 2 to 3 weeks.
FOR RESIDUAL CONTROL OF ANTS, COCKROACHES (Waterbugs), CLOVER MITES, CRICKETS, EARWIGS, FLEAS (Outdoors),
MOSQUITOES (Outdoors), HOUSE FLIES, HORNETS, MILLIPEDES, SAW-TOOTHED GRAIN BEETLE (Exposed stage),
SCORPIONS, SILVERFISH (Firebrats), SOWBUGS (Pillbugs), SPIDERS, BROWN DOG TICKS, WASPS.

ACTIVE INGREDIENTS:

o-isopropoxyphenyl methylcarbamate

Pyrethrins

oPiperonyl Butoxide, Technical

INERT INGREDIENTS:

ACCEPTED
JUL 20 1988
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registration under EPA Reg. No. 432-702.

0.50%w/w

0.05%

0.50%

98.95%

100.00%w/w

*Equivalent to 0.4% (butylcarbityl) (6-propyl piperonyl) ether and 0.1% related compounds.

**U.S. Patent No. 3,683,078.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada amplamente.

TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS AND
STATEMENT OF PRACTICAL TREATMENT

Can be used in non-food areas of food handling establishments and in food areas as a crack and crevice treatment only.

EPA Est.: See Lot No. For Correlation

EPA Reg. No. 432-702

G 42545-MO-1

P-5/15/88

H 5405-AR-1

P 655-GA-1

NET CONTENTS:

ROUSSEL BIO
CORPORATION

PO Box 1077 400 Sylvan Avenue
Englewood Cliffs, New Jersey 07632

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Harmful or Irritant. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. In case of expected prolonged exposure, wear natural rubber gloves, protective clothing and goggles. Do not use as a space spray. Provide adequate ventilation of area being treated. Do not apply to animals or humans. Do not use on food crops. Do not use in animal barns or poultry houses for fly control. In food handling areas, application is limited to crack and crevice treatment. Do not apply directly to food or feedstuffs. In commercial food handling facilities, cover or remove any food or food processing equipment during application. Do not apply while food processing is underway. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes and skin with plenty of water. If irritation develops or persists, seek medical attention.

PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not puncture or incinerate container. Do not use near heat or open flame. Exposure to temperatures above 130°F (54.5°C) may cause bursting.

Do not apply this product in conduits, motor housings, junctions, and switch boxes or other electrical equipment because of possible shock hazard. Do not use on surfaces that can be stained or harmed by water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Keep in locked storage area.

CONTAINER DISPOSAL: Replace cap and discard container in trash.

RESTRICTIONS: Since this is a water suspension material, do not apply to rugs, carpets, draperies, wallpaper, or similar materials that may be stained or watermarked. Shake well before each use. Remove protective cap. Hold can upright and aim spray opening away from person. Push button to spray.

DIRECTIONS FOR USE (Continue on reverse side)

DIRECTIONS FOR USE (Continued)

INDOORS: Pest control in ~~and~~ homes, apartment buildings, stores and warehouses: For residual control of ants, cockroaches, millipedes, silverfish, spiders, waterbugs and to aid in control of saw-toothed grain beetles (exposed stage). Apply as a residual spray, hitting as many insects as possible, to baseboards, window frames, the undersides of shelves and drawers, under sinks and stoves and to other places where insects may hide. Also spray cracks, crevices and surfaces where insects may crawl when they come out of hiding. For ants, spray trails and places where ants enter premises. Repeat as necessary.

Do not spray foodstuffs or house plants. Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated surfaces until surfaces are completely dry.

Pest control in meat packing and food processing plants, food storage areas, restaurants, and other food handling establishments (places other than private residences in which exposed food is held, processed, prepared or served). For residual control of ants, cockroaches, millipedes, silverfish, spiders, waterbugs, and to aid in control of saw-toothed grain beetles (exposed stage).

NON-FOOD AREAS: Includes garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Apply to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

FOOD AREAS: Limited to crack and crevice treatment only - Apply a small amount of material as a pin thin stream directly into cracks and crevices such as expansion joints between different elements of construction or between equipment bases and the floor, wall voids, or hollow equipment legs where the insects noted above hide. Do not use this product in conduits, motor housings and electrical switch boxes.

Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT, ARE NOT PERMITTED.

EPA NO. 432-702

Spraying unloading docks and ~~outside~~ areas where insects congregate will aid in preventing reinfestations. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

Pest control in animal research facilities: the use of this product in and around research animal quarters for pest control will not induce changes in liver microsomal enzyme activity in these animals.

PESTS OUT-OF-DOORS: DO NOT TREAT OR ALLOW THE SPRAY TO CONTACT PLANTS (LAWNS, FLOWERING PLANTS, SHRUBS, TREES, EVERGREENS, ETC.) BECAUSE OF POSSIBLE INJURY.

OUTDOORS: For residual control of insects around industrial buildings, homes, garbage areas, dumps.

RESIDUAL SPRAY for Ants, Cockroaches, Earwigs, House flies, Hornets, Millipedes, Mosquitoes, Spiders, Crickets, Clover Mites, Fleas, Scorpions, Sowbugs, Brown Dog Ticks, Wasps and Waterbugs. Apply as a residual spray hitting as many insects as possible, or with a paint brush to surface of buildings, porches, screens, window frames, patios, and garages where these insects congregate. For control of wasps and hornets application should be made in late evening when insects are at rest. Hold can upright. Point valve opening away from face. With wind at back aim at nest. Thoroughly saturate nest with spray, contacting as many insects as possible. To prevent insects from falling on skin and clothing do not stand directly under the nest. Repeat as necessary.

*Delete
BA*

WARRANTY

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with the directions or established safe practice. The buyer must assume all responsibility including injury or damage resulting from its misuse as such, combination with other materials.