



2/2

# SCOURGE® Insecticide with SBP-1382®/PIPERONYL BUTOXIDE 4% + 12% MF FORMULA II

## DIRECTIONS FOR USE (Continued)

This product is to be used for control of adult mosquitoes, midges (biting and non-biting) and blackflies in ground application with nonthermal or mechanical spray equipment that can deliver spray particles within the aerosol size range and at specified dosage levels.

**NOTICE:** This concentrate cannot be diluted in water. Mix well before using. Avoid storing excess formulation in spray equipment tank beyond the period needed for application.

### ULTRA LOW VOLUME APPLICATIONS

For use in nonthermal ULV portable backpack equipment similar to the Hudson B.P. Mix 70 fl oz (2068 ml) of this product with 1 gal (3.79 L) of refined soybean oil, light mineral oil of 54 second viscosity or other suitable solvent. Adjust equipment to deliver fog particles of 18-50 microns mass median diameter. Apply at the rate of 4.25-8.50 fl oz of finished formulation per acre (311-621 ml/ha) as a 50 ft (15.2 m) swath while walking at a speed of 2 mph (3.2 kph). This is equivalent to 0.0035-0.0070 lb ai SBP-1382/A (3.92-7.85 gm/ha) plus 0.0105-0.0210 lb ai piperonyl butoxide tech./A (11.77-23.54 gm/ha). Where dense vegetation is present, the higher rate is recommended.

For truck mounted nonthermal ULV equipment similar to LECO HD or MICRO-GEN or WHISPERMIST-XL. Adjust equipment to deliver fog particles of 8-20 microns mass median diameter. Consult the following chart for application rates.

Treatment lb ai/A of Scourge Wanted SBP-1382/PBO	Fl oz/A of Undiluted Spray to be Applied	Application Rate-Fl oz/Min	
		5 MPH	10 MPH
0.007/0.021	3.0(90 ml)	9.0(266.2 ml)	18.0(532.3 ml)
0.0035/0.0105	1.5(45 ml)	4.5(133.1 ml)	9.0(266.2 ml)
0.00175/0.00525	0.75(22.5 ml)	2.25(66.6 ml)	4.5(133.1 ml)
0.00117/0.00351	0.50(15 ml)	1.50(45 ml)	3.0(90 ml)

Where dense vegetation is present, the use of the higher rates and/or slower speed is recommended.

Fog only when air currents are 2-8 mph (3.2-12.9 kph). It is preferable to fog during early morning and evening when there is less breeze and convection currents are minimal. Arrange to apply the fog in the direction with breeze to obtain maximum swath length and better distribution. Direct spray head of equipment in a manner to insure even distribution of the fog throughout the area to be treated. Avoid prolonged inhalation of fog.

Guide the direction of the equipment so that the discharge nozzle is maintained at a distance of more than 6 feet (1.83 m) from ornamental plants and 15 feet (4.57 m) or more from painted objects. Temperature fluctuations will require periodical adjustment of equipment to deliver the desired flow rate at the specified speed of travel. The flow rate must be maintained to insure the distribution of the proper dosage of finished formulation.

Spray parks, campsites, woodlands, athletic fields, golf courses, swamps, tidal marshes, residential areas and municipalities around the outside of apartment buildings, restaurants, stores and warehouses. Do not spray on cropland, feed or foodstuffs. Avoid direct application over lakes, ponds and streams.

## EPA Reg. No. 432-716

### DIRECTIONS FOR STABLE FLY, HORSE FLY, DEER FLY CONTROL:

Treat shrubbery and vegetation where the above flies may rest. Shrubbery and vegetation around stagnant pools, marshy areas, ponds and shore lines may be treated. Application of this product to any body of water is prohibited.

For control of adult flies in residential and recreational areas, apply this product undiluted at a rate of 178 fl oz/hr (5.26 L/hr) by use of a suitable ULV generator travelling at 5 mph (8 kph) or at a rate of 356 fl oz/hr (10.53 L/hr) while travelling at 10 mph (16 kph). When spraying, apply across wind direction approximately 300 ft (91.4 m) apart.

Apply when winds range from 1-10 mph (1.6-16.0 kph). Repeat for effective control.

### DIRECTIONS FOR AERIAL APPLICATIONS FOR USE WITH FIXED-WING AND ROTARY AIRCRAFT

This product is used in specially designed aircraft capable of applying ultra low volume of undiluted spray formulation for control of adult mosquitoes (including organophosphate resistant species), midges (biting and non-biting) and blackflies.

Aerial application should be made preferably in the early morning or evening. Application should be made preferably when there is little or no wind.

It is not recommended to make application when wind speeds exceed 10 mph (16 kph). Repeat applications should be made as necessary. Apply preferably when temperatures exceed 50°F (10°C).

May be used as a mosquito adulticide in recreational and residential areas, and in municipalities, around the outside of apartment buildings, golf courses, athletic fields, parks, campsites, woodlands, swamps, tidal marshes, and overgrown waste areas.

Do not spray on cropland, feed or foodstuffs. Avoid direct application over lakes, ponds and streams.

### APPLICATION INSTRUCTIONS FOR PRODUCT IN AIRCRAFT USAGE

lb ai/A Wanted SBP-1382/PBO	Fl oz/A of Undiluted Spray to be Applied
0.007/0.021	3.0 (90 ml)
0.0035/0.0105	1.5 (45 ml)
0.00175/0.00525	0.75 (22.5 ml)
0.00117/0.00351	0.50 (15 ml)

PENICK-BIO UCLAF CORPORATION  
1050 Wall Street West, Lyndhurst, N.J. 07071