US ENVIRONMENTAL PROTECTION AGENCY		<u> </u>
	EPA REGISTRATION NO.	SEP - Q IQQQ
OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS. 767)	TERM OF ISSUANCE	0 1309
WASHINGTON, DC 20460		
NOTICE OF PESTICIDE: HE GISTRATION	NAME OF PESTICIDE PRODUC	, T
*Under the Federal Inserticide, Fungacide and Rodentia de Actais and medical		
AME AND ADDRESS OF REGISTRANT (Include ZIP code)		
,—	٦	18/2 200.36
Ĺ	7	,
NGTE. Changes in labeling formula differing in substance submitted to and accepted by the Registration Division priproduct always refer to the above U.S. EPA registration nu	or to use of the label in commerc	
On the basis of information furnished by the registrant, the	above named pesticide is hereb	y Registered/Reregistered under
the Federal Insecticide, Fungicide, and Rodenticide Act.		
A copy of the labeling accepted in connection with this Re		
Registration is in no way to be construed as an indorsement all hand the environment, the Administrator, on his moticule in accordance with the Act. The acceptance of any nact is not to be construed as giving the registrant a right toy others.	on, may at any time suspend or c ame in connection with the regis	ance) the registration of a pest- tration of a product under this
		e de la companie de l
,		

BEST AVAILABLE COPY

ATTACHMENT IS APPLICABLE		
SIGNATURE OF APPROVING OFFICIAL	Asil dellas	9/4/09
EPA Form 8570-6 (Rev. 5-76)	PRIVIOUS EDITION MAY BE USED UNTIL SUBS	

57500: I:CR-31: Heyward: LR-2: KENCO: 9/01/89:10/15/39:dg: aw: vo: ek:dg

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FTFKA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

George T. LaRocca Product Manager (15) Insecticide-Rodenticide Branch Registration Division (N7505C)

Enclosures

and Marcella.

Cyper-Active TM 2.14 EC SMALL

Crack And Crevice, Spot Spray For Residual Cockroach Control

For residual and contact control of cockroaches (waterbugs), crickets, sliverlish (firebrats), centipedes, millipedes, earwigs, nowbugt (pill) dog ticks, lone star ticks, ants, fless (including pre-adult fless), flies, mosquitoes, gnats, flying moths, wasps, horners, yellow jeci: Cyper-Active Contains 2.14 Pounds Active Ingredient Per Gallon

ONLY FOR SALE TO, USE AND STORAGE BY DE UNDER SUPERISION OF TRAINED PEST CON

ACTIVE INGREDIENTS:

Cypermethrin

 $(\pm)\alpha$ -cyano-(3-phenoxyphenyl) methyl (\pm) -cls,trans-3-

(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate*.......26.0% Oinert ingredients:74.0%

100.00%

OCONTAINS XY LENE RANGE Aronatic Solvent
Please read the entire label before using this product.

PRECAUCION AL CONSUMIDOR: \$1 usted no lee Ingles, no use este producto hasta que la etiquele le haya eldo explicada ampliamente.

[TO THE USER; if you cannot read English, do not use this product until the label has been fully explained to you.)

KEEP OUT OF REACH OF CHILDREN WARNING

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NIOT FOR USE IN USDA HEATAND POULTRY

Gen be used in non-load areas of load handling establishments such as; t ut auانتها، به au (to sewers), entries, floor drains (to sewers), entries, auانتها، au vestibules.and hellways, offices, locker rooms, machine rooms, basemer ta, boiler rooms, garages, utility sooms, mop closets and storage rooms (after canning and bottling).

EPA Reg. No. 432-

NET CONTENTS: LOT NO.

SEE ATTACHHENT

in EPA Lett r Dated:

ACCEPTED 1

with COMMENTS

Under the Federal in catacide for side and Rod word for an animaled, for the ge that. registered under EPA Reg. No.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARMING

MAY BERATAL IF SWALLOWED. CAUSES FYE AND SKIN IRRITATION on skin, or on clothing. Were acolective clothing when apraying. Wear impermeable gloves and full feet shield or goppies when handling or r Avoid breathing vapors or spray mist. Wash thoroughly with soap and a Barrove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Gall a physician immediately. IF IN EYES; Flush eyes with plenty of water. Call a physician if irritation IF ON SKIN: Wash with plenty of soap and water. Get medical attention NOTE TO PHYSICIAN

Inducing vomiting as first aid for this substance may result in increase pneumonia or pulmonary edema caused by aspiration of the hydrocarbo should be induced only under professional supervision.

ENVIRONMENTAL HAZARDS when This product is extremely toxic to fish. Do not contaminate water by-clear disposal of wattre. Do not apply directly to any body of water, Apply i specified on this label. Care should be used when apraying to avoid rish

PHYSICAL OR CHEMICAL HAZARDS Do not use or store near heat or open flame, Flashpoint 140°F Min. Do n product in conduits, motor housings, junction and switch boxes or other because of possible shock hazard. On not use on surfaces that can be h

DIRECTIONS FOR USE

PIANTS

STURAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dump not reuse empty container.

STORAGE: Store product in original container and keep container closed not store near load or feed. Protect from freezing, in case of spill or le surfaces, soak up with eand, earth or synthetic absorbent. Remove to ci PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Impropposticide, apray mixture or rinsale is a violation of Federal Law. If the disposed of by use according to label instructions, contact your Environmental Control Agency, or the Hazardour, Waste representative Regional Office for Guidance.

CONTAINER DISPOSAL: Metal Containers: Triple rinse (or equipment), Th or reconditioning, or puncture and dispose of in a sanitary lendfill, or approved by state and local authorities. Plactic Containers: Triple rinse offer for recycling or reconditioning, or puncture and dispose of in a incineration, or, if allowed by state and local authorities, by burning, i smake.

HOT FOR USE IN FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

FOR RESIDUAL PEST CONROL III AND ON BUILDINGS AND STRUCTURES AND THEIR NAMEDIATE SURROUNDINGS AND ON MODES OF TRANSPORT. PERMITTED AREAS OF USE INCLUDE, BUT ARE NOT LIMITED TO, NON-FOOD AREAS INDOORS AND OUTSIDE PREMISES OF MOTELS, MOTELS, HOMES, PRESCHOOLS, NURSERIES, DAY DAMPS, GREENHOUSES, LATHHOUSES, SEED HOUSES, FOOD HANDLING ESTABLISHMEN'I S, PACKAGING AND PROCESSING PLANTS, COMPACIAL, INDUSTRIAL, OFFICE, INSTITU-TIONAL AND STORAGE BUILDINGS, DAIRIES, MILK STORAGE ROOMS, EGG PROCESSING PLANTS, SUZZÁMARKETS, RESTAURANTS, BOTTLING PLANTS. CANNERIES, FLOUR AND FEED MILL, BAKERIES, LABORATORIES, AND HOS-PITALS. IN HOSPILATS: PATIENTS SHOULD BE REMOVED PRIOR TO TREAT-MENT. ALSO, JANUSE IN CAMPERS, MOBILE HOMES, BUSES, AUTOMOBILES, TRUCKS MAILERS, VESSELS, AIRCRAFT, AND BOXCARS. ALSO FOR USE IN DOGMENNELS AND HORSE STABLES. IN HORSE STABLES. NOT FOR APPLICA-FION IF HORSES WILL BE USED FOR FOOD.

Syper-Active insecticide is intended for dilution with water for apray application.

fill sprayer with the desired volume of water and add Cyper-Active insecticide. Close and shake before use in order to insure proper mixing. Shake or reagitate iprayer before use if spraying is interrupted. Make up only as required; repeat realments when needed.

		Concentration		
		of Active	Dilution	
Peets		Ingredient	Rate	
Ante	Files	0.1%	₩ fl. oz.	
Centipedes	Millipedes		per 1 gal.	
Cockroaches	Pilibugs		water	
(Maintenance)	Silverlish			
Pickela	Sowbuge			
irebrais	Gnais			
arwigs	Mosquitoes			
lees	Hornels	0.2%	1 fl, oz.	
:ockraeches	Yellow Jeckets		per 1 gal.	
(Clean-out)	Wasps		water	
lying Moline SMALL	cleas (& pre-eduit fless)			
lying Moths St	Brown Dog Ticks	DECT AND		
one Star Ticks		I DES! RVA!	I ARIT PROM	

INDOORS

:OCKROACHES, SPIDERS, CRICKETS, SILVERFISH AND FIREBRATS: Apply as a parse, low pressure spray to areas where these peats hide, such as baseboards, orners, storage areas, closets, around water pipes, doors and windows, attics, and aves, behind and under refrigerators, cabinets, sinks, furnaces, and stoves, the inderside of shelves, drawers and similar areas. Pay particular attention to cracks nd crevices.

> ANTS: To kill ante within buildings and homes, spray Iralia, neets and points of entry. Spray directly on anta where possible. Repeat M necessary.

EPAREG. No. 432--DO NET allow ! LEEN OR PETS TO CONTACT TREATME SURFACES UNTIL THE SHITAY has deced.

FLEAS (& PRE-ABULT FLEAS), BROWN DOG AND LONE STAR TICKS: Thoroughly apply as a spot treatment to infected areas such as pets' beds and resting quarters; nearby cracks and crevices; along and behind baseboards, window and doo; drames and localized areas of floors where these posts may be present. Gld-bcdding of pets should be removed and replaced with clean fresh bedding sifter traditions of pet reas, DO NOT TREAT PETS WITH THIS PRODUCT. Treat pets with a product registered for animal application such " a BP-13879/Bioallethrin Aqueous Pressurized Spray, EPA Reg. No. 432-482, to control the conrect of flee and tick infestation.

WASPS, HORNETS, YELLOW JACKETS, BEES: Application should be made in the late evening when insects are at rest. Spray liberally into hiding and breeding places, especially under attic rafters, contacting as many insects as possible. Repeat as necessary.

SOWBUGS, PILLBUGS, MILLIPEDES, EARWIGS, AND CENTIPEDES: Apply around doors and windows and other places where these peats may be found or where they may enter premises. Spray baseboards, storage areas and other locations. Ref. - to barrier treatment directions to prevent infer auon.

-SUA-L FLIES, MOSQUITOES, GNATS, FLYING MOTHS: Apply directly to wells, ceilings, window screens, and other flying insect resting areas as a residual surface treatment. May be used inside residential buildings as well as in and around carports, garages, and storage sheds.

OUTDOCAS

Do not apply this product to edible crops.

-SHALL

ANTS, BEES, "ENTIPEDES, COCKROACHES, CRICKETS, "REBRATS/MILLIPEDES, SILVERFISH, SOWBUGS, SPIDERS, FLIES, MOSQUITOES, GNATS, FLYING MOTHS, WASPS, HORNETS, YELLOW JACKETS, FLEAS , # PRE-ADULT FLEAS], BROWN DOG AND LONE STAR TICKS. Apply as a residual spray to surfaces of buildings, porches, acreens, wirdow frames, eaves patios, lawns, refuse dumps, garages in either hand or power sprayer. Treat other areas where these pests are found,

BARRIER TREATMENT: To help prevent infestation of buildings, apply to a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building. Also, treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Apply as a coarse spray to thoroughly and uniformly wel the band area, using 1 gallon of spray mix per 400 square feet.

WARRANTY STATEMENT

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established sate practice. The buyer must assume all responsibilities, including injury or damage, resulting from Its misuse as such, or in combination with other materials.

- #1. IF SWALLOWED: Do not induce vomiting. Call a physician or Poison Control Center. Do not give anything by mouth to an unconscious person.
- #2. IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration preferably mouth-to-mouth. Get medical attention.

#3. PRECAUTIONARY STATEMENTS

Causes substantial but temporary eye injury. Causes skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles, face shield or safety glasses. Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing spray. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

- #4. FOR RESIDUAL PEST CONTROL IN AND ON buildings and structures and their immediate surroundings and on modes of transport. PERMITTED AREAS OF USE INCLUDE industrial buildings, houses, apartment buildings, laboratories, greenhouses, and the nonfood/feed areas of stores, warehouses, vessels, railcars, trucks, trailers, aircraft, schools, nurseries, day camps, lath houses, seed houses, food handling establishments, packaging and processing plants, commercial, industrial, office, institutional and storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, flour and feed mill, bakeries, laboratories and hospitals. IN HOSPITALS: patients should be removed prior to treatment. Also for use in campers, mobile homes, buses, automobiles, and boxcars. Also for use in dog kennels and horse stables. IN HORSE STABLES: not for application if horses will be used for food.
- prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as diming rooms but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed.

Nonfood areas are areas such as (see 1st page of draft label for listing)

