

GOLD CREST®

Vengeance®

ACCEPTED
FEB 5 1992
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 432-748

**RODENTICIDE
SMALL BAIT PACKS**

Specimen Label

**For Control of Commensal Rodents
Kills Norway Rats, Roof Rats,
and House Mice**

Rats and mice consume a lethal dose in a single feeding with dead rodents appearing two to three days after treatment begins. Rodents cease feeding after consuming a lethal dose in a single-night's feeding. Vengeance bait is effective against anticoagulant-resistant rats and mice.

ACTIVE INGREDIENT:

Bromethalin: N-methyl-2,4-dinitro-N-(2,4,6-tribromophenyl)-6-(trifluoromethyl) benzenamine 0.01%

INERT INGREDIENTS: 99.99%
TOTAL: 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION
MAY BE HARMFUL OR FATAL IF SWALLOWED
SEE SIDE PANEL FOR
ADDITIONAL PRECAUTIONARY STATEMENTS

**U.S. Patent No. 4,815,923
E.P.A. Reg. No. 432-748
E.P.A. Est. No. 61282-WI-1(H) *
E.P.A. Est. No. 602-MO-1(P) ***

NET CONTENTS 3 POUNDS

(Contains 96 x 1/2 oz. Bait Packs)
* Letter refers to lot number of producing establishment.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Keep away from humans, domestic animals, and pets. May be harmful or fatal if swallowed. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Wash hands thoroughly after handling Vengeance bait.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician or local poison control center. Drink 1-2 glasses of water and induce vomiting by touching the back of throat with finger. Do not try to induce vomiting or give anything by mouth to an unconscious person. In case of accidental contact with eyes or skin, flush with plenty of water, and seek medical attention if irritation persists. Wash all contaminated clothing thoroughly. If bait is eaten by animals or pets contact a veterinarian.

Note to Physician: Vengeance bait is not an anticoagulant type of rodenticide. If ingested, limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store only in original container, in a cool, dry place inaccessible to children and pets and away from other chemicals. Keep containers closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse empty containers. Triple rinse container, and then dispose of in a sanitary landfill or by incineration if allowed by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice in and around homes, industrial and agricultural buildings, and similar man-made structures. This product may also be used in alleys located in urban areas, inside transport vehicles (ships, trains, aircraft), and in and around related port or terminal buildings. Do not use Vengeance bait in sewers. Vengeance baits must be placed in tamperproof bait boxes or in locations inaccessible to children, pets, domestic animals, or wildlife. Do not place baits in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

BEST AVAILABLE COPY

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners, and concealed places between floors and walls or in locations where rodents or their signs have been observed. Remove as much food accessible to rodents as possible.

APPLICATION DIRECTIONS: Vengeance bait is a new rodenticide with a mode of action completely unlike the anticoagulants. Generally rodents will consume a lethal dose in a single feeding, with death occurring 2 to 3 days later. A rodent will cease feeding after consuming a lethal dose.

NORWAY AND ROOF RATS: Place 1 to 8 ounces of bait per placement (usually at intervals of 20 to 30 feet). Adjust the amount applied to the level of rat feeding expected at each bait placement.

For burrow baiting, use 1 to 2 ounces per burrow.

When baiting burrows, insert each bait pack into the burrow far enough so that only a person who knows that bait is there would be likely to see it. Do not bait burrows in any location accessible to children or nontarget animals unless there is no other way to bait the target population effectively. If burrows in such an area are baited, check burrows every morning and evening to make sure that rodents have not pushed bait packs out of burrows and that baits have not become more accessible to children or nontarget animals in any way. Remove packs that become more accessible to nontarget animals. Remove the packages of bait from burrows if the bait has not been touched by target rodents for seven days. These packs may be relocated or properly disposed.

HOUSE MICE: Place ¼ ounce to 3 ounces of bait per placement (usually at intervals of 8 to 12 feet). Adjust the amount of bait applied to the level of mouse feeding expected at each bait placement.

BAITING STRATEGY: Highest bait consumption is expected to occur on the first day or two after treatment. After several days, inspect the placements and replenish bait at sites where there is evidence of heavy feeding. Normally, volumes of bait consumed will be significantly less with bromethalin baits than with anticoagulant baits. Although one night's feeding will usually provide a lethal dose, maintain bait for at least a week to provide all rodents a chance to feed on the bait. Baiting should be continued until all signs of feeding have ceased. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.


LIMITED WARRANTY AND LIABILITY

NOTICE: Read this Limited Warranty and Liability before buying or using this product. If the terms are not acceptable, return it at once unopened.

It is critical that this product be used only as specified on the label. The laws of a State may make some or all of this paragraph inapplicable or may give you rights in addition to your rights hereunder. Except to the extent prohibited by applicable law, the exclusive remedy of the user or buyer and the limit of liability of this Company or any other Seller for any and all losses, personal injuries or damages resulting from the use of this product, shall be the purchase price paid by the user or buyer for the quantity of product involved. Except to the extent prohibited by State law, there is no warranty, and this Company and other Sellers disclaim all liability for losses, personal injury or damages, whether indirect, special or consequential, when such claim is not reported to this company within one year of discovery and: (i) arises from any use of this product in a manner or for a purpose not recommended in its label directions or (ii) arises from handling or storage in violation of label instructions. THERE ARE NO IMPLIED WARRANTIES AND NO WARRANTIES OF MERCHANTABILITY OR FITNESS.

ROUSSEL BIO 
CORPORATION

P.O. Box 1077, 400 Sylvan Avenue
Englewood Cliffs, New Jersey 07632
Telephone: (201) 871-0771 Telex: ITT 420839



Gold Crest® and Vengeance® are registered trademarks of Roussel Bio Corporation

©Roussel Bio Corporation, 1989

SP-353R

MM 9/89

GOLD CREST®

Vengeance®

NOTE: If bait is eaten by animals or
pets: call for assistance
1-800-331-2867.

RODENTICIDE

ACTIVE INGREDIENT:

Bromethalin: N-methyl-2,4-dinitro-N-
(2,4,6-tribromo-phenyl)-6-(trifluoro-
methyl) benzenamine 0.01%
Inert Ingredients 99.99%

100.00%

Note to Physician: Vengeance bait is not
an anticoagulant type of rodenticide. If
ingested, limit absorption by either
emesis or gastric lavage. Sublethal
symptoms, if present, would be the
result of cerebral edema and should
be treated accordingly through
administration of an osmotic diu-
retic and corticosteroid.
Check appropriate E.P.A. registration
number below.

- E.P.A. Reg. No. 432-748
- E.P.A. Reg. No. 432-746

This sticker was applied on _____
at time Vengeance was added.

Pest Control Company _____
Address _____
City _____ State _____
Phone _____
Local Poison Control Center Phone _____

KEEP OUT OF REACH OF CHILDREN
CAUTION
 MAY BE HARMFUL OR FATAL IF SWALLOWED
 SEE PRODUCT LABEL FOR
 ADDITIONAL PRECAUTIONARY STATEMENTS

ROUSSEL BIO 
 CORPORATION

P.O. Box 1077, 400 Sylvan Avenue
Englewood Cliffs, New Jersey 07632

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
registered under the pesticide
EPA Reg. No. 432-748

ACCEPTED
FEB - 5 1992

BEST AVAILABLE COPY