Under the Federal Insecticid Fungicide, and Redenticide

AquaPy®

- Specially Formulated to Give Equal Performance with Oil-Based Sprays when Applied Undiluted or Diluted with Water
- For Indoor or Outdoor Application as a Space, Area or Contact Spray. Dependent upon Pests to be Controlled and the Area to be Treated, May be Applied Through Mechanical Aerosol Generators (ULV) or Conventional Fogging or Spraying Equipment
- Indoor Use Areas Unoccupied Livestock Shelters (Cattle Barns, Poultry Houses, Swine Houses) Bakeries, Beverage Plants, Canneries, Commercial Transportation Facilities, Daycare Centers, Dried Food Products in Storage or Processing, Flour Mills, Food and Feed Processing and Handling Sites, Grain Elevators, Granaries, Horse Barns, Hospices, Hospitals, Hotels, Homes Industrial Installations, Meat Packing Plants, Motels, Nonresidential Buildings, Nursing Homes, Office Buildings, Private and Commercial Dwellings, Railroad Cars, Restaurants, Schools, Ships' Holds, Stored Food in Multi-Walled Paper or Cloth Bags, Supermarkets, Truck Trailers, Warehouses, Zoos

ACTIVE INGREDIENTS:

Pyrethrins	3.0%
*Piperonyl Butoxide, Technical	
OTHER INGREDIENTS:	
	100.0%

*Equivalent to 12% (butylcarbityl)(6-propytpiperonyl) ether and 3% related compounds

AquaPy is a registered trademark of AgrEvo Environmental Health, Inc.

EPA Reg. No. 432-799

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN CAUTION FIRST AID

If Inhaled: Remove affected person to fresh air. Apply artificial respiration, preferably mouth-to-mouth, if indicated. Get medical attention.

See Side Panel for Additional Precautions

NET CONTENTS:

IN CASE OF MEDICAL EMERGENCIES OR HEALTH AND SAFETY INQUIRIES, OR IN CASE OFFIRE, LEAKING OR DAMAGED CONTAINERS, INFORMATION MAY BE OBTAINED BY CALLING 1-800-471-0660.

FOR PRODUCT INFORMATION, CALL TOLL FREE: 1-800-331-2867.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if inhaled. Avoid breathing vapors or spray mist. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

Do not use water-based sprays of AquaPy in conduits, motor housings, junction boxes or other electrical equipment because of possible shock hazard.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE OR AGITATE THOROUGHLY BEFORE DILUTING.

AquaPy is designed to be diluted with water. The unique patented formulation greatly reduces the evaporation rate of water from the spray droplets and enables the waterbased spray cloud to behave as that produced by a typical oil-based formulation.

AquaPy effectively controls the following insects when the applicator follows the directions for use as given below.

Angoumois Grain	Drugstore Beetles	Mosquitoes
Moths	_	Mushroom Flies
Ants	Fleas	Red Four Beetles
Bees	Fruit Flies	Rice Weevils
Boxelder Bugs	Fungus Gnats	Saw-toothed Grain
Cadelles	Grain Mites	Beetles
Cheese Mites	Granary Weevils	Silverfi sh
Cheese Skippers	Horn Flies	Small Flying Moths
Cigarette Beetles	Hornets	Spiders
Clover Mites	Horse Flies	Spider Beetles
Cockroaches	House Plies	Stable Flies
Confused Flour	Indian Meal Moths	Tobacco Moths
Beetles	Lice	Wasps
Crickers	Mediterranean	Vellow Mealworms

A CONTRACTOR OF THE PARTY OF TH

Deer Flies Dried Fruit Beetles Flour Moths

Dark Mealworms

INDOOR USE, UNDILUTED

The use of this product in food/feed processing or food/feed handling establishments (bakeries, beverage plants, canneries, fnod/feed processing plants, hospitals hotels, industrial installations, meat packing plants, restaurants, schools, supermarkets and warehouses) should be confined to time periods when the plants or facilities are not in operation. Food/feed should be removed or covered during treatment. All food/feed processing surfaces should be covered during treatment or thoroughly cleaned befire reusing.

Space and Contact Soray Application

To control accessible adult stages of:

Ante Boxelder Bugs Cheese Mites Cheese Skippers Clover Mites *Cockroaches (German) Crickets

Flics Fruit Flies Homers Mosquitoes Silvertish Small Flying Moths

Wasps

and the following common stored product pests:

Angoumois Grain Moths Cadelles Cigarette Beetles Confused Flour Beerles Dark Mealworms Drugstore Beetles

Granary Weevils Grain Mites

Indian Meal Moths Mediterranean Flour Motiss Red Flour Beetles Rice Weevils Saw-toothed Grain Beerles

Spider Beetles Tobacco Moths Yellow Mealworms

Close doors and windows and shut off ventilating systems. Vacate the area. Only the applicator should be in the treatment area during application. Use a mechanical acrosol generator or a fogger adjusted to deliver acrosol droplets (30 microns or less). Direct the spray first into all cracks and crevices, behind all equipment, cupboards, obstructions or dark harborage areas to provide maximum penetration and impingement on the insects. Then treat as a space spray, directing the spray towards the ceiling and upper corners of the area and behind any obstructions. apply at a rate of 6 milliliters or 0.20 ounces per 1000 cubic feet of space (or I oz per 5,000 cu. ft.). Keep area closed for at least one hour. Leave the treated area and ventilate before reoccupying. Repeat treatment as necessary.

*Rates up to 9 ml or 0.30 oz. per 1,000 cu. ft. may be required for control of some German Cockroach

populations (dependent on temperature, sanitation and other factors).

Treatment of Stored Food/Feed Areas

Foods/feeds in multi-walled bags or in cloth bags stored in warehouses, storage rooms, flour mills, grain elevators and

To control accessible adult stages of:

Angoumois Grain Moths Cadelles Confused Flour Beetles Indian Meal Moths Mediterranean Flour Moths Rice Weevils Saw-toothed Grain Beetles Yellow Mealworms

Cheese Mites Cheese Skippers Cigarette Beetles Grain Mites Granary Woovils Dark Mealworms German Cockroaches Red Flour Beetles

Infested stored products should be destroyed, furnigated or treated by other effective methods using an approved product intended for this purpose. Vacate the area. Only the applicator should be in the treatment area during application. As noted above, disperse AquaPy first by directing spray into cracks and crevices, corners, under pallets, and around stacks of packaged goods - hitting as many insects as possible and then apply as a space spray at a dosage of 4 milliliters or 0.13 ounces per 1000 cubic feet of room space (or 1/2 oz. per 4,000 cu. ft.). Repeat application as necessary for insect control. Leave the treated areas closed for one hour. Do not remain in the treated area and ventilate before reentry.

ACTISOL SPRAY EQUIPMENT

Deco Harborage: To kill Cockroaches, Carpenter Ants, Termites, Spiders, Bees, Wasps and other labeled pests, treat wall voids, insect tunnels, boxed beams and other hollow elements of construction, using UNDILUTED product through mechanical aerosol generators such as ACTISOL equipment, inject void for 5-10 seconds followed by sufficient air to pressurize the void.

Crack and Crevice Treatment: For treatment of Cockroaches, Silverfish, Ants, Waterbugs, Crickets, Centipedes, Sowbugs, Millipedes, Spiders, Scorpions, Mites, Bedbugs, Lice, Carpet Beetles, Wasps, Hornets, Yellow jackets and Grain Insects, crack and crevice treatments with ACTISOL equipment, using undiluted product, represents a one second injection into openings. seams, holes, etc.

INDOOR USE, DILUTED

Space Spray Application: For use in mechanical fogging or spray equipment.

To control accessible stages of:

FliesGnats Fruit Flies Small Flying Moths Mosquitoes

Cover or remove exposed food/feed and cover food/feed handling surfaces in edible product areas of food/feed handling establishments. Close rooms and shut off air conditioning or ventilating equipment. Vacate the area. Only the applicator should be in the treatment area during application. Mix 16 ounces of AquaPy with sufficient water to equal 1 gallon of diluted spray. Apply at the rate of 8 fluid ounces per 1000 cubic feet, filling the room with mist. Keep area closed for at least 15 minutes. Leave the treated area and ventilate before reoccupying. Repeat treatment if reinfestation occurs. Do not apply while food/feed processing is underway.

ANIMAL QUARTER USE

(Cattle barns, horse barns, poultry houses, swine houses, zoos): As a space spray, dilute for use in conventional fogging or spraying equipment.

To control Fleas, Flies, Gnats, Mosquitoes, Small Flying Moths: Mix 16 ounces of AquaPy with sufficient water to equal one gallon of diluted spray. Vacate the treatment area. Animals must be removed prior to treatment. Only the applicator should be in the area during application. Apply at the rate of 8 ounces per 1000 cubic feet of space. Direct spray towards the upper portions of the enclosure filling the room with mist. Keep area closed for at least 15 minutes. Leave the treated area and ventilate before reoccupying. Repeat treatment if reinfestation occurs.

SURFACE SPRAY APPLICATION

To control accessible stages of:

Ants Crickets Fless

German Cockroaches

Mix 2 to 4 ounces of AquaPy with sufficient water to equal one gallon of diluted spray. Apply as a coarse droplet spray using approximately one gallon to 1000 square feet of surface area. Spray walls, floors and other surfaces of bins and storage and handling areas thoroughly.

NOTE: Over-application may cause discoloration of light colored surfaces.

Fleas: AquaPy is effective as part of a total management program for Flea Control. It is suggested that infested areas be thoroughly vacuumed prior to treatment. These areas include all carpet, upholstered furniture, baseboards and other indoor areas frequented by pets. The vacuum cleaner bag should be discarded in an outdoor trash container. Pet bedding should be cleaned or replaced.

Apply AquaPy alone or in combination with an appropriate IGR (see below), to rugs, carpets, upholstered furniture, pet beds, floor coverings, pet resting areas and other infested habitats. The quantity of finished spray required is dependent upon carpet/fabric density. Provide uniform coverage, apply until infested area is moist.

Effective flea control includes controlling the source of flea infestations. This can require treatment of pets with a product registered for such use and treating outside areas frequented by pets. AquaPy can also be applied out-of-doors for flea control. Do not treat pets with this product. Do not allow pets to contact treated areas until spray has dried. Remove animal and feed before spraying.

In Combination with Insect Growth Regulators (IGRs): To extend control of immature stages of cockroaches and fleas, tank-mix the recommended rate of AquaPy with the correct amount of IGR per finished gallon. Follow all directions for use patterns and spray intervals from the IGR label to achieve maximum benefit.

For In-Transit Protection: Clean and spray truck beds, boxcars and ships' holds before loading. Thoroughly treat walls and floors and spray each layer of cartons or containers lightly as stacked.

Treatment of Stored Grain: To control accessible adult stages of Indian Meal Moths and Angoumois Grain Moths: Apply to the surface of binned grain at a rate of 1 gallon per 1000 square feet. Apply at monthly intervals with the first appearance of moths and continue until early fall. In severe infestations, break up webbing with a rake before spraying and make a second application in two weeks.

<u>Treatment of Flying Insects' Resting Places:</u> To aid in the control of Mosquitoes, Small Flying Moths and Gnats: Treat walls, ceiling, moldings, screens, door frames, window frames, beams, light cords and similar resting places with the above dilution of AquaPy.

OUTDOOR APPLICATION, DILUTED

For use on horses:

NOT FOR USE ON HORSES WHICH WILL BE USED FOR FOOD.

To protect against Horn Flies, House Flies, Fleas, Gnats and Mosquitoes: Mix 13 ounces of AquaPy with sufficient water to equal one gallon of diluted spray. Apply a light mist sufficient to wet the tips of the hair.

To protect against Stable Flies, Horse Flies, Deer Flies: Apply the above dilution at two ounces per adult horse to sufficiently wet the hair but not to soak the hide. Repeat this application once or twice daily or as needed to give continued protection.

AccounTable To System

		Food Processing Plants		Commercial		Residential	
AEROSOL SPI	RAY	Operating 1	Nonoperating	Fabric	Without Fabric	Fabric	Without Fabric
ACTISOL SPRAY	Space Spray		x	x	x	x	x
EQUIPMENT	Deep Harborage/ Crack & Crevice	х	X	x	x	х	X
COARSE WET	SPRAY						
	Surface Spray	x	x		x	х	x
	Crack & Crevice	х	х		х	х	x

¹ Pesticide applications are permitted in non-USDA inspected facilities when they are operating.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or discosal.

Persicide Storage and Spill Procedures: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

Pessicide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved State and local procedures.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645