03 432-864 8-28-48

160 5544012 /Of 4



U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Program Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

432-864

AU3 28 1998

Term of Issuance: Conditional

NOTICE OF PESTICIDE:

 $\underline{x}$  Registration

Reregistration

(under FIFRA, as amended)

Name of Pesticide Product;

Permanone Insect

Killer R400

Name and Address of Registrant (include ZIP Code):

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- Make the labeling changes listed below before you release the product for shipment:
  - a. Add the phase "EPA Registration No. 432-864".

Signature of Approving Official:	Date:	

EPA Form 8570-6

page 2 EPA Reg. No. 432-864

- b. Made the following additions to the label to be in compliance with PR Notice 96-6 (Pet Pesticide Product Label Statements). "READ ENTIRE LABEL BEFORE USE." "USE ONLY ON DOGS." Clarify Reapplication Limitation. "Do not treat puppies less than 12 weeks old." (unless you have domestic animal safety data on file at the Agency to support use on younger animals.). "Consult a veterinarian before using this product on debilitated, aged, pregnant, medicated or nursing animals.". "Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.".
- 3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

#### **CODE 72053**

## PERMANONE®

- \* FOR DOMESTIC USE BOTH INDOORS AND OUTDOORS.
- \* A READY-TO-USE PERMANONE STABLE EMULSION INSECTICIDE
- FOR CONTROL OF FLEAS, BROWN DOG TICES AND LICE ON PREMISES AND ON DOGS
- \* CONTROLS FLEAS AND TICKS ON DOGS FOR 35 DAYS
- AIDS IN CONTROL OF TICKS ON DOGS FOR AN ADDITIONAL 2 WEEKS
- FOR CONTROL OF PESTS ON LIVESTOCK
- WITH COUNTRY FRESH FRAGRANCE
- PROVIDES 60-DAY (TWO MONTHS) CONTBOL OF COCK-ROACHES
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF FLEAS
- PROVIDES 60-DAY (TWO MONTHS) CONTROL OF STORED PRODUCT PESTS

#### **ACTIVE INGREDIENT:**

*Permethrin	0.25%
INERT INGREDIENTS:	
	100.00%

\*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-2,2-dichlorzethenyl) 2,2-di-methylcyclopropanecarboxylate. Cis/trans ratio: min. 35% (+/-) cis and max. 65% (+/-) trans.

PERMANONE - Registered Trademark of Agreeo Environmental Health, Inc.

EPA REG. NO. 432-JAV EPA EST. NO.

# KEEP OUT OF REACH OF CHILDREN CAUTION

See Side Panel For Additional Precautions.

**NET CONTENTS:** 

ACCEPTED
with COMMENTS
in EPA Letter Dated

/AUG 2 8 1998

Under the Federal Insertibile, Fungiside, and Rodentiest. Act as amenated, for the permitte registered under EPA Rea. No.

#### 756mastr 091096tk

### PRECAUTIONARY STATEMENTS Hazards To Humans & Domestic Animals

CAUTION

Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

#### **Environmental Hazards**

This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

SHAKE WELL BEFORE USING

INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors.

SURFACE SPRAYING: FOR 60-DAY (TWO MONTHS) CONTROL OF: COCKROACHES, WATERBUGS, PALMETTO BUGS, SCORPIONS, MILLIPEDES, CENTIPEDES, SOWBUGS, PILLBUGS, ANTS, SILVERFISH, FIREBRATS, SPIDERS, CRICKETS, CLOVER MITES, CHEESE MITES, GRANARY WEEVILS, RICE WEEVILS, CONFUSED FLOUR BEETLES, RUST RED FLOUR BEETLES, DRUGSTORE BEETLES, MEAL WORMS, GRAIN MITES AND CADELLES: Direct the spray into hiding places, cracks and crevices, and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

TO CONTROL CARPET BEETLES: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving. Repeat treatment as needed.

TO CONTROL BEDBUCS: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

TO CONTROL FLEAS (ADULTS AND LARVAE) AND TICKS (ADULTS AND LARVAE): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment: Repeat treatment as needed. Allow to dry thoroughly before all cyring pets to contact treated areas.

TO CONTROL CLOTHES MOTHS: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

TO KILL LICE AND LOUSE EGGS: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated particles to dry thoroughly before use.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from Hom Flies, Face Flies, House Flies, Mosquitoes and Gnals: Apply a light mist sufficient to wet the surface of the hair. For face flies, spray face and head, but do not spray into eyes.

TO CONTROL STABLE FLIES, HORSE FLIES AND DEER FLIES: Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

TO CONTROL BLOOD-SUCKING LICE: Apply to the infested areas

of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required.

TO CONTROL POULTRY LICE: Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

TO CONTROL BEDBUCS AND MITES IN POULTRY HOUSES: Spray crevices of roost poles, cracks in walls, and cracks in nests where the *Bedbugs* and *Mites* hide.

#### ON DOCS:

TO CONTROL FLEAS, TICKS, AND LICE: Start spraying at the tail, moving the dispenser rapidly and making sure that the animal's entire body is covered, including the legs and underbody. While spraying, fluff the hair so that the spray will penetrate to the skin. Make sure spray wets thoroughly, but do not saturate animal. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as necessary.

NOTE: Not recommended for use on puppies less than four weeks old. DO NOT USE ON CATS OR KITTENS.

#### **OUTDOOR APPLICATION**

To help control ticks in outdoor activity areas, treat a band of vegetation and leaf litter 6-10 feet wide and adjacent to activity areas, especially where dense vegetation occurs.

FOR CONTROL OF GARDEN INSECTS: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as Whitellies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, camations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms.

When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90°F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given becervith.

AgrEvo Environmental Health 95 Chestnut Ridge Road Montvale, NJ 07645