



United States  
Environmental Protection Agency  
Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number 432-887	2. EPA Product Manager Eugene Wilson	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Chipco Ronstar 50WP	PM# Team 23	
5. Name and Address of Applicant (Include ZIP Code) Bayer Environmental Science 95 Chestnut Ridge Road Montvale, NJ 07645 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

**Section - II**

Amendment - Explain below.  
 Resubmission in response to Agency letter dated \_\_\_\_\_  
 Notification - Explain below.

Final printed labels in response to Agency letter dated \_\_\_\_\_  
 "Me Too" Application.  
 Other - Explain below.

**NOTIFICATION**

MAR 14 2003

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.  
(1) Added Bayer Environmental Science to Warranty Statement replacement the Aventis name.

**Section - III**

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt. No. per container	

3. Location of Net Contents Information  
 Label  Container

4. Size(s) Retail Container

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product  
 Lithograph  
 Paper glued  
 Stenciled  Other \_\_\_\_\_

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Richard M. Gorrell	Title Manager, Registrations	Telephone No. (Include Area Code) (201) 307-3376
----------------------------	---------------------------------	---

**Certification**  
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Manager, Registrations	6. Date Application Received (Stamp)
4. Typed Name Richard M. Gorrell	5. Date March 6, 2003	

# CHIPCO® RONSTAR® 50 WP

Herbicide

MASTER LABEL

---



## General label revisions:

- Added the Bayer Environmental Science Company name in the Warranty Statement replacing the Aventis name.

## Printing Restrictions:

**NOTIFICATION**

MAR 14 2003

3/10

# CHIPCO® RONSTAR® 50 WP

## Herbicide

Selective Preemergent Herbicide for the Control of Annual Grasses and Broadleaf Weeds In Turf and Ornamentals

FOR SALE TO, USE AND STORAGE BY COMMERCIAL NURSERY, TURF AND LANDSCAPE PERSONNEL ONLY. NOT FOR USE ON HOME LAWNS.

### ACTIVE INGREDIENT:

Oxadiazon (2-tert-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)-Δ-1, 3, 4-oxadiazolin-5-one) ..... 50%

### INERT INGREDIENTS:.....

50%

EPA Reg. No. 432-887

EPA Est.

## KEEP OUT OF REACH OF CHILDREN WARNING AVISO

NOTIFICATION

MAR 14 2008

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-334-9745

### FIRST AID

If swallowed:	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told t by a poison control center or doctor.</li> </ul>
If in eyes:	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin:	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled:	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

### PRECAUTIONARY STATEMENTS

#### WARNING

#### HAZARDS TO HUMANS

Causes substantial but temporary eye injury. Causes skin irritation. Harmful if inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear coveralls over a short-sleeved shirt and short pants; waterproof gloves; a chemical-resistant apron when cleaning equipment, mixing or loading; protective eyewear; chemical resistant headgear for overhead exposure; and chemical resistant footwear plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

4/10

### **User Safety Recommendations**

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash body thoroughly and put on clean clothing.

### **ENVIRONMENTAL HAZARDS**

Do not contaminate water by cleaning of equipment or disposal of wastes. RONSTAR® should not be used in areas where irrigation or rainfall results in the direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxidiazon is adsorbed.

### **DIRECTIONS FOR USE**

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls over short-sleeved shirt and short pants, waterproof gloves, protective eyewear, chemical-resistant footwear plus socks and chemical-resistant headgear for overhead exposure.

### **STORAGE AND DISPOSAL**

#### **STORAGE**

Do not contaminate water, food or feed by storage or disposal. Do not store in or around the home.

#### **PESTICIDE DISPOSAL**

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### **CONTAINER DISPOSAL**

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### **GENERAL INFORMATION**

CHIPCO® RONSTAR® 50 WP is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf and on a variety of woody ornamental shrubs and trees. CHIPCO® RONSTAR® 50 WP controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Apply CHIPCO® RONSTAR® 50 WP with a properly calibrated sprayer that will apply the spray solution uniformly (uniform application is essential for satisfactory weed control). Calibrate the sprayer prior to use, according to the manufacturer's directions and check daily to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

5/10

## TURF

For use in dormant, established Bermudagrass, St. Augustinegrass and Zoysia turf such as fairways, parks, golf courses and lawns. CHIPCO® RONSTAR® 50 WP controls Crabgrass, Goosegrass, Annual Bluegrass, Annual Sedge and Virginia Buttonweed.

CHIPCO® RONSTAR® 50 WP should be applied at least 2-3 weeks prior to green up. Rake turf area free of leaves, mowing if necessary so the spray solution reaches the soil surface. Measure the area to be treated and uniformly apply the amount required. If rain is not expected shortly after application, sprinkle the turf thoroughly.

**DO NOT USE ON ST. AUGUSTINE GRASS OR ZOYSIA TURF IN CALIFORNIA.**

### PREEMERGENCE WEED CONTROL

WEEDS CONTROLLED	AMOUNT OF CHIPCO® RONSTAR® 50 WP		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass ( <i>Digitaria spp.</i> ) Goosegrass ( <i>Eleusine indica</i> ) Annual Bluegrass ( <i>Poa annua</i> )* Annual Sedge ( <i>Cyperus compressus</i> )*	4 - 6 lbs.	1.5 - 2.2 ounces	Apply in late winter or early spring prior to weed seed germination and turf greenup. Treatment is effective only against spring germinating <i>Poa annua</i> . Apply in 30 - 150 gallons of water per acre or 0.75 - 3.5 gallons of water per 1000 sq. ft.
Virginia Buttonweed ( <i>Diodia virginiana</i> )	6 lbs.	2.2 ounces	

\*Not currently registered in California.

### ANNUAL WEED CONTROL DURING BERMUDAGRASS, AND ZOYSIAGRASS ESTABLISHMENT FROM SPRIGS.

WEEDS CONTROLLED	AMOUNT OF CHIPCO® RONSTAR® 50 WP		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass, Goosegrass, and Annual Sedge*	4 - 6 lbs.	1.5 - 2.2 ounces	Make application immediately prior to or immediately after sprigging. Disturbance of the soil surface after CHIPCO® RONSTAR® 50 WP application may disrupt the herbicide barrier and reduce effectiveness. Irrigate immediately after application for best results.

Do not apply RONSTAR® 50 WP to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. RONSTAR® 50 WP should not be used on putting greens or tees because of the varying cultural practices used.

\*Not currently registered in California.

## ORNAMENTALS AND CONIFER NURSERIES

**WOODY ORNAMENTAL SHRUBS, TREES, AND CONIFER NURSERY APPLICATION:** Apply CHIPCO® RONSTAR® 50 WP prior to weed seed germination, as a directed spray to the soil surface or as an over the top spray only on labeled species. CHIPCO® RONSTAR® 50 WP can be applied to both newly transplanted and established plants. Application can be made to actively growing or dormant plants. However, CHIPCO® RONSTAR® 50 WP should not be applied during bud break or during the 4 weeks after bud break.

Rainfall or overhead irrigation (one-half inch) immediately after application of CHIPCO® RONSTAR® 50 WP will improve weed control activity. Rainfall or overhead irrigation immediately after over the top sprays is required to move the CHIPCO® RONSTAR® 50 WP from the foliage to the soil surface.

CHIPCO® RONSTAR® 50 WP should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of CHIPCO® RONSTAR® 50 WP. **Do not apply in greenhouses** as plant injury may result from codistillation of CHIPCO® RONSTAR® active ingredient onto leaf surfaces in condensation.

6/10

**AMOUNT TO USE:** Ground Application - Apply CHIPCO® RONSTAR® 50 WP broadcast, in a minimum of 50 gallons per acre, to ornamentals at 4 to 8 pounds of product per acre, depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

CHIPCO® RONSTAR® 50 WP at 4 to 8 pounds per acre is effective in controlling annual grasses and broadleaf weeds such as:

ANNUAL BROADLEAF WEEDS		ANNUAL GRASSES	
Bittercress	( <i>Cardamine spp.</i> )	Annual Bluegrass <sup>†</sup>	( <i>Poa annua</i> )
Bristly Oxtongue <sup>†</sup>	( <i>Picris echioides</i> )	Annual Sedge <sup>†</sup>	( <i>Cyperus compressus</i> )
Carpetweed	( <i>Mollugo verticillata</i> )	*Barnyardgrass	( <i>Echinochloa crus-galli</i> )
Cheeseweed <sup>†</sup>	( <i>Malva parviflora</i> )	Carpetgrass	( <i>Axonopris offinis</i> )
*Common Groundsel	( <i>Senecio vulgaris</i> )	Crabgrass	( <i>Digitaria spp.</i> )
Common Purslane	( <i>Portulaca oleracea</i> )	Fall Panicum	( <i>Panicum dichotomiflorum</i> )
*Evening Primrose	( <i>Oenothera spp.</i> )	**Goosegrass	( <i>Eleusine indica</i> )
Fiddleneck <sup>†</sup>	( <i>Amsinckia intermedia</i> )	Green Foxtail	( <i>Setaria viridis</i> )
Fireweed <sup>†</sup>	( <i>Epilobium angustifolium</i> )	Ripgut Bromegrass <sup>†</sup>	( <i>Bromus catharticus</i> )
Florida Pusley	( <i>Richardia scabia</i> )	Wild Oats <sup>†</sup>	( <i>Avena fatua</i> )
*Galinsoga	( <i>Galinsoga spp.</i> )		
Golden Ragwort	( <i>Senecio aureus</i> )		
Lambsquarters	( <i>Chenopodium album</i> )		
**Liverwort	( <i>Marchantia spp.</i> )		
Oxalis	( <i>Oxalis spp.</i> )		
Pennsylvania Smartweed	( <i>Polygonum pensylvanicum</i> )		
Petty Spurge	( <i>Euphorbia pepus</i> )		
**Prostrate Spurge	( <i>Euphorbia supina</i> )		
Redroot Pigweed	( <i>Amaranthus retroflexus</i> )		
Shepherdspurse <sup>†</sup>	( <i>Capsella bursa-pastoris</i> )		
Speedwell	( <i>Veronica spp.</i> )		
Sow Thistle <sup>†</sup>	( <i>Sonchus oleraceus</i> )		
Spotted Catsear	( <i>Hypochaeris radicata</i> )		
*Swinecress	( <i>Coronopus didymus</i> )		
Yellow Woodsorrel	( <i>Oxalis stricta</i> )		

\*Use 6 to 8 pounds of product per acre.

\*\*Use 8 pounds product per acre.

<sup>†</sup> Not currently registered in California for use on Bristly Oxtongue, Cheeseweed, Fiddleneck, Fireweed, Shepherdspurse, Sow Thistle, Annual Bluegrass, Annual Sedge, Ripgut Bromegrass, and Wild Oats.

7/10

**CROP TOLERANCE:** CHIPCO® RONSTAR® 50 WP can be used as directed spray to the soil surface on a wide variety of container and field grown woody ornamental shrubs and trees and on conifers in nurseries. CHIPCO® RONSTAR® 50 WP may be used as an over the top spray in a minimum of 50 gallons per acre, at the recommended rates, on the following ornamental species only:

<b>SPECIES .</b>	<b>VARIETIES</b>
Arborvitae	American - Smaragd, Rheingold Giant - Zebrina
Baby's Breath	<i>Gypsophila paniculata</i>
Barberry	Japanese
Boxwood	Dazzler
Cedar	Eastern Red
Cranberry	American
Dodonaea	Purpurea
Euonymus	Evergreen - Golden Japanese, Gold Spot
False Cypress	Nootka -Aurea Hinoki - Graciosa, Pygmaea Sawara - Aurea Nana
Fir*	Douglas, Fraser, Grand, Noble
Gladiolas	
Hemlock*	Canada
Holly	Japanese - Helli Kurogane Chinese - Parsonii
Juniper	Common - Berkshire, Effusa Creeping - Blue Chip, Blue Rug, Andorra Singleseed - Blue Star Savin - Buffalo, Tamariscifolia
Oleander	Common
Ophiopogon	
Pine	Eastern White, Mugho, Scotch
Raphiolepis	
Spruce*	Blue
Statice	<i>Limonium spp.</i>

Local conditions may affect the use of this chemical. Consult State Agricultural Experiment Station or Extension Service weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

\* Not currently registered for use on Fir, Hemlock, and Blue Spruce in California.

**LANDSCAPE PLANTINGS: COMMERCIAL, ROADSIDE, & INDUSTRIAL SITES.** Apply CHIPCO® RONSTAR® 50 WP prior to weed seed germination as an over the top spray only on labeled ornamentals.

**AMOUNT TO USE:** Apply CHIPCO® RONSTAR® 50 WP broadcast, in a minimum of 50 gallons of water per acre, to established ornamentals at 4 pounds of product per acre. Remove existing weed growth before application. Do not apply within 60 days of transplanting. Do not apply with adjuvants or drift control agents. Sprinkle irrigate within 2 hours after application, otherwise crop injury may result.

**CROP TOLERANCE:** CHIPCO® RONSTAR® 50 WP may be used as an over the top spray on the following ornamental species only:

<b>SPECIES</b>	<b>VARIETIES</b>
Ice Plant	Large Leaf
Ivy	English, Algerian
Daisy	African Trailing
Periwinkle	Major, Minor

8/10

**TANK-MIX COMBINATIONS:** Ornamental Plantings and Noncropland areas:

Tank-mix combinations of RONSTAR® plus a non-selective herbicide such as FINALE® are recommended to control existing undesirable vegetation. Applied as a directed spray, RONSTAR® plus a non-selective herbicide such as FINALE® will provide pre and postemergence control of susceptible species listed on the respective labels of the two herbicides.

A compatibility test must be conducted with any potential tank mix partner with Ronstar ® WP Herbicide. Using a clear glass quart jar, conduct the test as described below:

1. Fill the jar three-quarters full with water
2. Add the appropriate amount of herbicide in the following order: (a) dry flowable, (b) wettable powder, (c) aqueous suspensions, (d) flowables, (e) liquids and (f) solutions and emulsifiable or liquid concentrates. Shake or gently stir jar after each addition to thoroughly mix.
3. After adding all ingredients, let the mixture stand for 15 minutes and then look for separation, large flakes, precipitates, gels, and heavy oily film on the jar or other signs of incompatibility.
4. If the compatibility test shows signs of incompatibility, do not tank mix the product tested with Ronstar ® WP Herbicide.

**MIXING INSTRUCTIONS:** RONSTAR® 50 WP should be added to the tank and agitated prior to introduction of a non-selective herbicide such as FINALE®. Please refer to the non-selective herbicide label for information on surfactant and spray volume to use. Only mix the amount of spray needed for immediate application needs as settling can occur if spray mixture is allowed to sit over an extended period of time.

**PRECAUTION:** Non-selective herbicides such as FINALE® are not recommended as an over the top spray in ornamentals. Extreme care must be exercised to avoid contact of spray containing a non-selective herbicide with foliage, stems, green bark, or bare roots of desirable turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or destruction may result. If spraying areas adjacent to desirable plants with a non-selective herbicide, use a shield made of cardboard, sheet metal, or plyboard while spraying to help prevent spray from contacting foliage of desirable plants.

**BROADCAST APPLICATION RATES:**

**RONSTAR®/FINALE®**

Existing Annual and Broadleaf Weeds	RONSTAR® 50 WP	FINALE®
	Pounds Per Acre	Quarts Per Acre
Less Than 6" Tall	4 - 8	3
Greater Than 6" Tall	4 - 8	4
Existing Perennial	4 - 8	4 - 6

These rates are intended as general information. Consult rate sections in individual product labels for detailed information on specific weed species controlled and rates of application.



9/10

## **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer Environmental Science. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE ELECTION, THE REPLACEMENT OF PRODUCT

CHIPCO, RONSTAR and FINALE are registered trademarks of the Bayer Ag.

Bayer Environmental Science  
95 Chestnut Ridge Road  
Montvale, NJ 07645

CHIPCO® RONSTAR® 50 WP (MASTER) Notification Corrected Typo Errors 3/3/03.

# Bayer Environmental Science



March 6, 2003

EPA03RMG0306Ronstar50WPBayertoWarranty  
Fedex

Ms. Sherada Hobgood  
U.S. Environmental Protection Agency  
Office of Pesticide Programs (7504C)  
Registration Division (7504C)  
Document Processing Desk (Notification)  
Room 266A  
1921 Jefferson Davis Hwy  
Arlington, VA 22202-4501

RE: Chipco Ronstar 50WP (EPA Reg. No. 432-887Notification)

Dear Ms. Sherada Hobgood:

Bayer Environmental Science is notifying the Agency of a change to the Warranty Statement. The company name Bayer Environmental Science replaced the Aventis name in the Warranty Statement.

Bayer Environmental  
Science  
95 Chestnut Ridge Road  
Montvale, NJ 07645  
Phone : 201.307.9700

If you have any questions, contact me at (215) 757-3185 or email me at [mike.gorrell@bayercropscience.com](mailto:mike.gorrell@bayercropscience.com).

Sincerely,

Richard M. Gorrell  
Manager, Registrations