



## **FUNGICIDE**

Supplemental Label

\* A Fungicide for the Prevention and Control of Certain Diseases of Turfgrass

**EPA Reg No. 432-888** 

E.P.A. Est. No. 264-MO-2

## **ATTENTION**

- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for 26GT Fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of 26 GT according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for 26 GT.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Refer to the 25 GT product label for General Use Precautions, Mixing and Application Instructions.

TURF: 26GT Fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf coures, sod farms, and institutional areas where fine turf is grown. When used in conjunction with good turf management practices, 26GT Fungicide is effective in controlling the following disease.

Apply the recommended rates as indicated in the table in 0.5 to 10 gallons of water per 1000 square feet. Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

## USE ON BERMUDAGRASS TURF ONLY APPLICATION DIRECTIONS

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

DISEASE	INTERVAL OF APPLICATIONS	FLUID 0Z./1000 FT.2
Curvularia (Curvularia sp.)	Use as a preventative every 14 days as long as required.	4

26 CT is a registered trademark of Bayer.



Bayer Environmental Science
A Business Group of Bayer CropScience LP

Bayer Environmental Science 95 Chestnut Ridge Road, Montvale, NJ 07645 www.bayerprocentral.com

