



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 20, 2022

Nina S. Rao
Regulatory Affairs
Bayer CropScience, Environmental Science Division
5000 CentreGreen Way, Suite 400
Cary, NC 27513

Subject: Label Amendment – Change REI to 24 hours
Product Name: Aliette WDG Brand Fungicide
EPA Registration Number: 432-890
Application Date: 10/20/2021
Decision Number: 579433

Dear Ms. Rao:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 432-890
Decision No. 579433

with FIFRA section 6. If you have any questions, please contact Jennifer Drobish by phone at 202-566-2642 or via email at Drobish.jennifer@epa.gov.

Sincerely,

A handwritten signature in black ink, reading "Shaja B. Joyner". The signature is written in a cursive style with a large, stylized initial "S".

Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P

Enclosure

ALUMINUM TRIS

GROUP

P07

FUNGICIDE

ALIETTE® WDG brand Fungicide

ABN: Aliette

ACTIVE INGREDIENT: Aluminum tris (O-ethyl phosphonate)..... 80.0%

OTHER INGREDIENTS: 20.0%

TOTAL: 100.0%

E P A Reg. No. 432-890

EPA EST. No.

KEEP OUT OF REACH OF CHILDREN CAUTION

See [Back] [Side] Panel for First Aid Instructions and [Leaflet] [Booklet] for Complete Precautionary Statements and Directions for Use.
(Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-800-331-2867.

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks, and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic and estuarine invertebrates. Drift and runoff may be hazardous to aquatic and estuarine organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters or rinsate.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours.

Exception: If the product is soil injected, soil incorporated, or applied by drenching, the Worker Protection Standard under certain circumstances, allows workers to enter the treated area without restrictions if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of the treated areas until sprays have dried.

DIRECTIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including micro jet, drip, solid set, and center pivot. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First, prepare a suspension of ALIETTE® WDG brand Fungicide in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of ALIETTE WDG brand Fungicide, and then the remaining volume of water. Then set sprinkler to deliver 0.1 to 0.3 inches of water per acre. Start sprinkler and uniformly inject the suspension of ALIETTE WDG brand Fungicide into the irrigation water line so as to deliver the desired rate per acre. The suspension of ALIETTE WDG brand Fungicide should be injected with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: For Micro jet and Drip Irrigation Systems: When treatment with ALIETTE WDG brand Fungicide has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours. **For Solid Set and Center Pivot Irrigation Systems:** When treatment with ALIETTE WDG brand Fungicide has been completed, further field irrigation over the treated area should be avoided until foliage is dry to prevent washing the chemical off the plants.

INSTRUCTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e. g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

Resistance Management

For resistance management, please note that ALIETTE WDG brand Fungicide contains a Group P07 [phosphonates] fungicide. Any fungal population may contain/develop individuals naturally resistant to ALIETTE WDG brand Fungicide and other Group P07 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

ORNAMENTALS AND BEDDING PLANTS

Use ALIETTE WDG brand Fungicide on ornamentals and bedding plants grown in field nursery, greenhouse, landscaping, and conifer nursery situations, for control of diseases caused by *Pythium* and *Phytophthora*. ALIETTE WDG brand Fungicide can also be used on ornamentals for the control of downy mildew and fire blight and for the suppression of bacterial blight caused by certain pathovars of *Xanthomonas campestris*. Applications should be made prior to disease development and should be made in conjunction with good cultural management practices. Use the higher rate when disease pressure is severe. Do not exceed recommended rates or apply more frequently than at specified intervals or plant injury may occur.

RESTRICTIONS

- When applying use spray nozzles that will deliver fine or larger spray droplets as defined in the American Society of Agricultural and Biological Engineers (ASABE) standard ANSI/ASAE S572.1 (March 2009).
- For foliar applications on ornamental herbaceous plants, ornamental nonflowering plants, and ornamental woody shrubs and vines, do not apply more than 11 times per year, [per] [container] [pot] [or] [plant].
- For applications using chemigation on ornamental herbaceous plants, ornamental nonflowering plants, and ornamental woody shrubs and vines, do not apply more than 11 times per year, [per] [container] [pot] [or] [plant].
- For soil drench applications on ornamental herbaceous plants, ornamental nonflowering plants, and ornamental woody shrubs and vines, do not apply more than 4 times per year, [per] [container] [pot] [or] [plant].
- For applications to ornamental conifers and conifers in nurseries, do not apply more than 9 times per year, [per] [container] [pot] [or] [plant].
- Apply by ground application equipment only. Do not apply by air unless directed by Supplemental or Special Local Need labeling.

ORNAMENTALS

FOLIAR APPLICATIONS to plants such as *Aglaonema*, *Aphelandra*, Azalea, *Bougainvillea*, Boxwood, *Cattelya skinneri*, *Cissus*, *Dieffenbachia*, *Hibiscus*, Juniper, Leather-leaf Fern, *Pittosporum*, *Philodendron*, Pothos, *Rhododendron*, *Spathiphyllum* and *Taxus media*.

Mix 2.5 to 5.0 pounds of product with 100 gallons of water and spray to wet. If less than 100 gallons of spray solution is applied per acre, then apply 2.5 to 5.0 pounds of product per acre. Do not exceed 400 gallons of spray solution per acre. Repeat as necessary but do not exceed one application every 30 days.

DRENCH APPLICATIONS to plants such as *Aphelandra*, Azalea, Boxwood, *Cissus*, *Dieffenbachia*, Japanese Holly, Juniper, Monterey Pine, *Philodendron*, *Pieris*, *Pittosporum*, *Rhododendron*, *Schefflera*, *Spathiphyllum*, and *Taxus media*:

Mix 0.4 to 0.8 pounds (6.4 to 12.8 ounces) of product with 100 gallons of water and apply to 400 square feet (equivalent to 2 pints. solution per square feet). Repeat as necessary but do not exceed one application every 30 days.

SOIL INCORPORATION to plants such as *Rhododendron*, Azalea, or *Pieris* for control of *Phytophthora* species only:

Mix 0.5 to 0.8 pounds (8 to 12.8 ounce) of product per cubic yard of soil mix immediately before potting. If conditions are favorable for disease development, applications as a foliar spray or drench should be made as needed but are not to exceed once every 30 days. Use soil incorporation only with well-rooted plants.

BEDDING PLANTS

FOLIAR APPLICATIONS to plants such as Begonia, Pansy, Vinca, Marigold, Zinnia, Petunia, Geranium, and Impatiens:

Mix 1.25 to 4 pounds of product with 100 gallons of water and spray to wet using no more than 400 gallons of water per acre. Repeat as necessary but do not exceed one application every fourteen days.

FOR USE ON CONIFERS IN NURSERIES* TO PREVENT PHYTOPHTHORA ROOT ROT

DIP TREATMENTS to conifers such as Douglas Firs, Spruce, and Pines. Dip transplants in 2.5 pounds product per 100 gallons to thoroughly wet plant and root mass. Dip immediately before transplanting. When making dip applications, wear chemical/water resistant gloves, goggles or face shield, long pants (coveralls), long sleeved shirt, shoes, and socks.

FOLIAR APPLICATIONS to conifers such as Douglas Firs, Spruce, and Pines. Mix 2.5 to 5.0 pounds of product in 100 gallons of water and spray to wet. If less than 100 gallons of spray solution is applied per acre, then apply 2.5 to 5.0 pounds of product per acre. Do not exceed 400 gallons of spray solution per acre. Do not exceed more than one application every thirty days.

***Not currently registered in California.**

DOWNY MILDEW CONTROL IN ROSES

FOLIAR APPLICATIONS TO ROSES: field, container, landscape and mini varieties to control downy mildew (*Peronospora sparsa*).

Applications should be made in conjunction with a disease sanitation program to reduce the spread of the disease to uninfected plants. Mix 2.5 pounds of product with 100 gallons of water and spray to wet. If less than 100 gallons of spray solution is applied per acre, then apply 2.5 to 5.0 pounds of product per acre. Do not exceed 400 gallons of spray solution per acre. Repeat as necessary but do not exceed one application every seven days.

FIRE BLIGHT SUPPRESSION

FOLIAR APPLICATIONS to plants such as Ornamental Pear, *Pyracantha*, and Hawthorn. Applications should be made in conjunction with a strict sanitation program to reduce the spread of the disease to uninfected plants.

Mix 2.5 pounds of the product in 100 gallons of water and spray to wet. If less than 100 gallons of spray solution is applied per acre, then apply 2.5 to 5.0 pounds of product per acre. Do not exceed 400 gallons of spray solution per acre. ALIETTE works solely as a preventive treatment. Begin spray treatments at pre-bloom stage and continue at 7-day intervals until bloom period ends. Do not exceed one application every seven days.

BACTERIAL BLIGHT SUPPRESSION**

FOLIAR APPLICATIONS to plants such as *English Ivy*, *Schefflera*, *Anthurium*, *Dieffenbachia*, *Spathaphyllum*, *Syngonium*, and *Ficus* for the suppression of the *Xanthomonas campestris* pathovars *hederae*, *dieffenbachiae*, *syngonii*, and *fici*. Applications should be made in conjunction with a disease sanitation program to reduce the spread of the disease to uninfected plants.

Mix 2 to 4 pounds of product with 100 gallons of water and spray to wet. If less than 100 gallons of spray solution is applied per acre, then apply 2.5 to 5.0 pounds of product per acre. Do not exceed 400 gallons of spray solution per acre. Repeat as necessary but do not exceed one application on every 7 days.

Refer to compatibility statements concerning use of coppers or other compounds.

**** Not currently registered in California.**

PLANT TOLERANCE

Plant tolerances to ALIETTE® WDG Fungicide have been found to be acceptable in the specific genera and species listed on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to ALIETTE WDG brand Fungicide. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

ORNAMENTAL AND BEDDING PLANT TANK MIXTURES

Additional Disease Control: If ornamentals are threatened by additional diseases, ALIETTE WDG brand Fungicide is compatible with most commonly used fungicides. For control of diseases such as Aerial Web Blight, Alternaria Leaf Blight, Botrytis Blight, Fusarium Leaf Spot, Helminthosporium Leaf Spot, and Ray Blight, ALIETTE WDG brand Fungicide can be tank mixed with CHIPCO® 26019 Brand Fungicide. Consult the CHIPCO® 26019 Brand Fungicide label for a complete listing of diseases controlled and for application rates.

If a tank mix with other fungicides is used, follow label directions for the use of that product and apply at the recommended rate for control of the target disease organism. ALIETTE WDG brand Fungicide is not compatible with the flowable formulation of Rubigan® A.S. (fenarimol; EPA Reg. No. 10163-274) and Spectro® 90WDG (chlorothalonil and thiophanate-methyl; EPA Reg. No. 1001-72). These products should not be combined for application.

COMPATIBILITY

ALIETTE WDG brand Fungicide may not be compatible with foliar fertilizers or with copper based fungicides. To avoid undesirable copper phytotoxicity, do not apply ALIETTE WDG brand Fungicide and copper-based compounds at less than 14-day intervals. Also, if copper based products and ALIETTE WDG brand Fungicide are included in the same foliar spray program it is advised to avoid application with air blast equipment.

Do not mix ALIETTE WDG brand Fungicide with any sticker, extender or wetting agent.

To determine the physical compatibility of ALIETTE WDG brand Fungicide with any other product, use a small container to mix a small amount (e.g. 1 pint) of spray solution, containing all ingredients in the same order and ratio as the anticipated use. If any indication of physical incompatibility develops, do not use this mixture for spraying. Indication of incompatibility usually appears within 5-15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

ALIETTE WDG brand Fungicide is not physically compatible with flowable formulations of Rubigan® A.S. (fenarimol; EPA Reg. No. 10163-274) and Spectro® 90 WDG (chlorothalonil and thiophanate-methyl; EPA Reg. No. 1001-72) as a tank mixture. These products should not be combined for application.

TURF

ALIETTE WDG brand Fungicide is a systemic fungicide, which may be used in a seasonal program for the control of *Pythium* diseases, such as blight and root rot, and yellow tuft on common turf grasses on golf courses, sod farms sports fields, residential, commercial and other turfgrass areas. Apply as a foliar spray, using 1 to 5 gallons of water per 1000 sq. ft., as indicated in the table below. Apply with a properly calibrated sprayer.

When applying use spray nozzles that will deliver fine or larger spray droplets as defined in the American Society of Agricultural and Biological Engineers (ASABE) standard ANSI/ASAE S572.1 (March 2009).

Do not exceed a total of 180 lb. of product per acre (144 lb. a.i. /A), per year.

Apply by ground application equipment only. Do not apply by air unless directed by Supplemental or Special Local Need labeling.

DISEASE	INTERVAL OF APPLICATIONS	RATE OZ. PRODUCT/1000 SQ. FT.
<i>Pythium</i> diseases, yellow tuft	14 days	4.0
	21 days	8.0

Begin preventive applications when conditions first favor disease and repeat at recommended interval. Do not mow and/or water treated areas until foliage is completely dry. Maintain agitation during spray operations.

TURF TANK MIXTURES

An ALIETTE WDG brand Fungicide program encourages the development of a thick, vigorous turf stand that is better able to tolerate various stresses (heat, humidity, drying winds, traffic etc.) that can have a negative effect on turf health. A thick, vigorous ALIETTE WDG brand Fungicide treated turf is better able to resist disease development and suppress the development of other pests such as weeds and algae.

Summer Decline: For control of summer decline caused by a complex of *Pythium* and *Rhizoctonia* diseases, ALIETTE WDG brand Fungicide can be tank mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

Anthracnose: For improved control of anthracnose ALIETTE WDG brand Fungicide can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. A preventative, multi-application (repeated every 14 days) program must be started prior to the appearance of the disease and the conditions favorable to anthracnose.

Bentgrass Dead Spot: For control of bentgrass dead spot, ALIETTE WDG brand Fungicide can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. A preventative, multi-application (repeated every 14 days) program must be started prior to the appearance of the disease and the conditions favorable for development of bentgrass dead spot.

PRODUCT [†]	RATE OZ. PRODUCT/1000 SQ. FT.	DISEASE	APPLICATION INTERVAL
ALLETTE WDG brand Fungicide	4.0 to 8.0	Summer Decline (<i>Pythium</i> , Anthracnose, and <i>Rhizoctonia spp.</i>)	14 days
+	+	<i>and/or</i>	
26GT [®]	3.0 to 4.0	Anthracnose (<i>Colletotrichum graminicola</i>)	
or	or		
Tartan [®]	1.0 to 2.0		
Daconil Ultrex [®]	3.2		
or	or		
Manicure [®] 6FL	3.3		
or	or		
Dithane [®] 75DF Rainshield [®]	8.0		
or	or		
Protect [™] DF	8.0		
or	or		
Mancozeb DG	8.0		
or	or		
Fore [®] 80WP Rainshield [®]	4.0 to 8.0		
or	or		
Heritage [®]	0.2 to 0.4		
or	or		
Banner [®] MAXX [®]	1.0 to 2.0		
or	or		
ProStar [®] 70WP	1.5 to 3.0		

ALLETTE WDG brand Fungicide		Bentgrass Dead Spot (<i>Ophiosphaerella agrostis</i>)	14 days
+	4.0		
+	+		
Fore® 80WP Rainshield®	8.0		
or	or		
Daconil Ultrex®	3.2		

Do not exceed a total of 180 lb. of product per acre (144 lb. a.i. /A), per year.

‡26GT® (iprodione; EPA Reg. No. 432-888); Tartan® (triadimefon and trifloxystrobin; EPA Reg. No. 432-1446); Daconil Ultrex® (chlorothalonil; EPA Reg. No. 100-1694); Manicure® 6FL (chlorothalonil; EPA Reg. No. 60063-7-10404); Dithane® 75DF Rainshield® (mancozeb; EPA Reg. No. 62719-402); Protect™ DF (mancozeb; EPA Reg. No. 1001-77); Mancozeb DG (mancozeb; EPA Reg. No. 62719-402-10404); Fore® 80WP Rainshield® (mancozeb; EPA Reg. No. 62719-388); Heritage® (azoxystrobin; EPA Reg. No. 100-1093); Banner® MAXX® (propiconazole; EPA Reg. No. 100-741); ProStar® 70WP (flutolanil; EPA Reg. No. 432-1223)

Mixing Instructions

Add ALIETTE WDG brand Fungicide to the tank after the tank-mix partner has been added and allowed to fully disperse. The order in which products are added to the tank can affect its ability to go into solution.

Apply as a foliar spray using 1 to 5 gallons of water per 1000 sq. ft. Apply with a properly calibrated sprayer. Begin preventative applications when conditions first favor disease and repeat at recommended interval. Do not mow and/or water treated areas until foliage is completely dry. Maintain agitation during spray operations.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

COMPATIBILITY

ALIETTE WDG brand Fungicide may not be chemically compatible with foliar fertilizers or with copper-based fungicides, and phytotoxicity may occur. Do not mix ALIETTE WDG brand Fungicide with any sticker, extender, or wetting agent.

To determine the physical compatibility of ALIETTE WDG brand Fungicide with any other product, use a small container to mix a small amount (e.g. 1 pint) of spray solution, containing all ingredients in the same order and ratio as the anticipated use. If any indication of physical incompatibility develops, do not use this mixture for spraying. Indication of incompatibility usually appears within 5 - 15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

ALIETTE WDG brand Fungicide is not physically compatible with Rubigan™ A.S. (fenarimol; EPA Reg. No. 10163-274) and Spectro™ 90WDG Fungicide (chlorothalonil and thiophanate-methyl; EPA Reg. No. 1001-72) as a tank mixture. These products should not be combined for application.

Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

PESTICIDE DISPOSAL

Pesticides wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

CONTAINER HANDLING

Rigid Non-refillable Containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs.)

Non-refillable container. Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Non-Seed Treatment Products in Non-Refillable Fiber Drums with Liners

Non-refillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Non-Seed Treatment Products in Non-Refillable Foil outer pouches of Water soluble Packets (WSP)

Offer foil pouch for recycling if available or dispose of empty pouch in the trash as long as WSP is unbroken.

Rigid Non-Refillable containers with capacities smaller or equal to 5 gallons**PLASTIC CONTAINERS:**

Non refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

LIQUID Dilutable formulations:

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

SOLID Dilutable formulations:

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Non-Seed Treatment Products in Non-Rigid, Non-refillable Containers = BAGS

Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, and Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Warranty, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury or other property damage, or other unintended consequences, may result because of factors beyond the control of Bayer CropScience LP such as weather conditions, presence of other materials or the manner of use or application. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 5 lbs.

Aliette, Tartan, 26GT, CHIPCO and ProStar are registered trademarks of Bayer.
Banner MAXX, Daconil Ultrex and Heritage are registered trademarks of Syngenta.
Fore, Rainshield and Dithane are registered trademarks of Corteva Agriscience.
Rubigan is a registered trademark of Gowan Company.
Manicure is a registered trademark of LESCO, INC.
Protect is a trademark and Spectro is a registered trademark of Cleary Chemicals, LLC.

PRODUCED FOR



Bayer Environmental Science
A Division of Bayer CropScience, LP
5000 CentreGreen Way, Suite 400
Cary, NC 27513

[Marketing Claims]

- For Greenhouse and Nursery Use
- Fungicide
- Fungicide / Bactericide
- Ornamental Fungicide

ALUMINUM TRIS

GROUP

P07

FUNGICIDE

Sub Label of Aliette WDG

CHIPCO® SIGNATURE Fungicide

ACTIVE INGREDIENT: Aluminum tris (O-ethyl phosphonate)..... 80.0%
 OTHER INGREDIENTS: 20.0%
 TOTAL: 100.0%

EPA REG NO. 432-890

EPA EST NO.

[This XX lb. bag is not for individual sale]

[This case can only be sold as a single unit]

KEEP OUT OF REACH OF CHILDREN

CAUTION

See [Back] [Side] Panel for First Aid Instructions and [Leaflet] [Booklet] for Complete Precautionary Statements and Directions for Use.
 (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For PRODUCT USE Information Call 1-800-331-2867.

FIRST AID

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin or inhaled. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. Applicators and other handlers must wear long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks, and protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic and estuarine invertebrates. Drift and runoff may be hazardous to aquatic and estuarine organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters or rinsate.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove PPE clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical resistant gloves made of any waterproof material, shoes plus socks, and protective eyewear.

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of the treated areas until sprays have dried.

DIRECTIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product only through sprinkler irrigation systems including microjet, drip, solid set, and center pivot. Do not apply this product through any other type of irrigation system.

SPRAY PREPARATION: Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: First, prepare a suspension of CHIPCO® SIGNATURE in a mix tank. Fill tank with 1/2 to 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Add the required amount CHIPCO SIGNATURE, and then the remaining volume of water. Then set sprinkler to deliver 0.1 to 0.3 inches of water per acre. Start sprinkler and uniformly inject the suspension of CHIPCO SIGNATURE into the irrigation water line so as to deliver the desired rate per acre. The suspension of CHIPCO SIGNATURE should be injected with a positive displacement pump into the main line ahead of a right angle turn to ensure adequate mixing. If you should have any other questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: For Microjet and Drip Irrigation Systems: When treatment with CHIPCO SIGNATURE has been completed, further field irrigation over the treated area should be avoided for 24 to 48 hours. **For Solid Set and Center Pivot Irrigation Systems:** When treatment with CHIPCO SIGNATURE has been completed, further field irrigation over the treated area should be avoided until foliage is dry to prevent washing the chemical off the plants.

INSTRUCTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connection or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop may result from nonuniform distribution of treated water. Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

This product may cause staining. Avoid off-target sites such as sidewalks, patios, driveways, pavers, or similar materials. Do not enter treated areas until spray has dried. Rinse spray equipment thoroughly with warm water and detergent.

Resistance Management

For resistance management, please note that ALIETTE WDG brand Fungicide contains a Group P07 [phosphonates] fungicide. Any fungal population may contain/develop individuals naturally resistant to ALIETTE WDG brand Fungicide and other Group P07 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

TURF

CHIPCO SIGNATURE is a systemic fungicide, which may be used in a seasonal program for the control of *Pythium* diseases, such as blight and root rot, and yellow tuft on common turfgrasses on golf courses, sod farms, sports fields, residential, commercial and other turfgrass areas. Apply as a foliar spray, using 1 to 5 gallons of water per 1000 sq. ft., as indicated in the table below. Apply with a properly calibrated sprayer.

When applying use spray nozzles that will deliver fine or larger spray droplets as defined in the American Society of Agricultural and Biological Engineers (ASABE) standard ANSI/ASAE S572.1 (March 2009).

Do not exceed a total of 180 lb. of product per acre (144 lb. a.i. /A), per year.

Apply by ground application equipment only. Do not apply by air unless directed by Supplemental or Special Local Need labeling.

DISEASE	INTERVAL OF APPLICATIONS	RATE OZ. PRODUCT/1000 SQ. FT.
<i>Pythium</i> diseases, yellow tuft	14 days	4.0
	21 days	8.0

Begin preventive applications when conditions first favor disease and repeat at recommended interval. Do not mow and/or water treated areas until foliage is completely dry. Maintain agitation during spray operations.

Bacterial Wilt

Chipco Signature is prescribed for the alleviation of symptoms on golf course greens affected with bacterial wilt. Apply 8 oz. of Chipco Signature in 2 to 4 gallons of water per 1000 sq. ft. at an interval of 7 to 14 days depending on severity of the outbreak. Use the more frequent interval when severe symptoms appear.

TURF TANK MIXTURES

A CHIPCO SIGNATURE program encourages the development of a thick, vigorous turf stand that is better able to tolerate various stresses (heat, humidity, drying winds, traffic etc.) that can have a negative effect on turf health. A thick, vigorous CHIPCO SIGNATURE treated turf is better able to resist disease development and suppress the development of other pests such as weeds and algae.

Summer Decline: For control of summer decline caused by a complex of *Pythium* and *Rhizoctonia* diseases, CHIPCO SIGNATURE can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

Anthracnose: For improved control of anthracnose CHIPCO SIGNATURE can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. A preventative, multi-application (repeated every 14 days) program must be started prior to the appearance of the disease and the conditions favorable to anthracnose.

Bentgrass Dead Spot: For control of bentgrass dead spot, CHIPCO SIGNATURE can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. A preventative, multi-application (repeated every 14 days) program must be started prior to the appearance of the disease and the conditions favorable for development of bentgrass dead spot.

PRODUCT ¹	RATE OZ. PRODUCT/1000 SQ. FT.	DISEASE	APPLICATION INTERVAL
CHIPCO [®] SIGNATURE	4.0 to 8.0	Summer Decline (<i>Pythium</i> , Anthracnose, and <i>Rhizoctonia spp.</i>)	14 days
+	+		
26GT [®]	3.0 to 4.0	<i>and/or</i>	
or	or	Anthracnose (<i>Colletotrichum graminicola</i>)	
Tartan [®]	1.0 to 2.0		
Daconil Ultrex [®]	3.2		
or	or		
Fore [®] 80WP Rainshield [®]	4.0 to 8.0		
or	or		
ProStar [®] 70WP	1.5 to 3.0		

CHIPCO® SIGNATURE	4.0	Bentgrass Dead Spot (<i>Ophiosphaerella agrostis</i>)	14 days
+	+		
Fore® 80WP Rainshield®	8.0		
or	or		
Daconil Ultrex®	3.2		

Do not exceed a total of 180 lb. of product per acre (144 lb. a.i. /A), per year.

'26GT® (iprodione; EPA Reg. No. 432-888); Tartan® (triadimefon and trifloxystrobin; EPA Reg. No. 432-1446); Daconil Ultrex® (chlorothalonil; EPA Reg. No. 100-1694); Fore® 80WP Rainshield® (mancozeb; EPA Reg. No. 62719-388); ProStar® 70WP (flutolanil; EPA Reg. No. 432-1223)

Mixing Instructions

Add CHIPCO SIGNATURE to the tank after the tank mix partner has been added and allowed to fully disperse. The order in which products are added to the tank can affect its ability to go into solution.

Apply as a foliar spray using 1 to 5 gallons of water per 1000 sq. ft. Apply with a properly calibrated sprayer. Begin preventative applications when conditions first favor disease and repeat at recommended interval. Do not mow and/or water treated areas until foliage is completely dry. Maintain agitation during spray operations.

If a tank mix with other fungicides is used, follow label directions for the use of that product and apply at the recommended rate for control of the target disease organism.

COMPATIBILITY

CHIPCO SIGNATURE may not be chemically compatible with foliar fertilizers or with copper-based fungicides and phytotoxicity may occur. Do not mix Chipco® Signature with any sticker, extender, or wetting agent.

To determine the physical compatibility of CHIPCO SIGNATURE with any other product, use a small container to mix a small amount (e.g. 1 pint) of spray solution, containing all ingredients in the same order and ratio as the anticipated use. If any indication of physical incompatibility develops, do not use this mixture for spraying. Indication of incompatibility usually appears within 5 - 15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

CHIPCO SIGNATURE is not physically compatible Rubigan™ A.S. (fenarimol; EPA Reg. No. 10163-274) and Spectro™ 90 WDG (chlorothalonil and thiophanate-methyl; EPA Reg. No. 1001-72) Fungicide as a tank mixture. These products should not be combined for application.

Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

PESTICIDE DISPOSAL

Pesticides wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

CONTAINER HANDLING

Rigid Non-refillable Containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs.)

Non-refillable container. Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums,

Kegs information as follows.

Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

Non-Seed Treatment Products in Non-Refillable Fiber Drums with Liners

Non-refillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

Non-Seed Treatment Products in Non-Refillable Foil outer pouches of Water soluble Packets (WSP)

Offer foil pouch for recycling if available or dispose of empty pouch in the trash as long as WSP is unbroken.

Rigid Non-Refillable containers with capacities smaller or equal to 5 gallons

PLASTIC CONTAINERS:

Non refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

LIQUID Dilutable formulations:

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

SOLID Dilutable formulations:

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Non-Seed Treatment Products in Non-Rigid, Non-refillable Containers = BAGS

Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, and Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Warranty, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury or other properly damage, or other unintended consequences, may result because of factors beyond the control of Bayer CropScience LP such as weather conditions, presence of other materials or the manner of use or application. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 5.5 lbs.

Aliette, Tartan, 26GT, CHIPCO and ProStar are registered trademarks of Bayer.
Banner MAXX, Daconil Ultrex and Heritage are registered trademarks of Syngenta.
Fore, Rainshield and Dithane are registered trademarks of Corteva Agriscience.
Rubigan is a registered trademark of Gowan Company.
Manicure is a registered trademark of LESCO, INC.
Protect is a trademark and Spectro is a registered trademark of Cleary Chemicals, LLC.

PRODUCED FOR



Bayer Environmental Science
A Division of Bayer CropScience, LP
5000 CentreGreen Way, Suite 400
Cary, NC 27513

CHIPCO SIGNATURE FUNGICIDE (PENDING) 10/01/2019, 12/07/2021, 12/08/2021, 01/06/2022, 01/07/2022