

432-898

9/8/2009

Page 18 14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

SEP 8 2009

Mr. Richard M. Gorrell
Registrations Manager
Bayer Environmental Science
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Subject: Label Notifications for Pesticide Registration Notice 2007-4

Dear Mr. Gorrell,

The Agency is in receipt of your Applications for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated August 7, 2009 and your label resubmissions on September 2, 2009 for the following products:

Chipco® Ronstar® G T/L (EPA Registration Number 432-898)
Ronstar® 2% Plus Turf Fertilizer (EPA Registration Number 432-1430)
Herbicide

The Registration Division (RD) has conducted its review of these requests for their applicability under PRN 2007-4 and finds that the label changes requested fall within the scope of PRN 2007-4. The labels submitted with the applications have been stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please call me directly at 703-305-6249 or Steve Schaible of my staff at 703-308-9362.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2814



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 432- 1400 898	2. EPA Product Manager Ms. Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Ronstar G T/L	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Bayer Environmental Science 2 T. W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input checked="" type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application. NOTIFICATION
<input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below. SEP - 8 2009

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance of PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and it may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

- Container Disposal Statement Revision as per EPA mandate


Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
				<input checked="" type="checkbox"/> Paper	Other (Specify) Paper
3. Location of Net Contents Information <input type="checkbox"/> Label <input checked="" type="checkbox"/> Container		4. Size(s) Retail Container 50 lb paper bag		5. Location of Label Directions <input checked="" type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Richard M. Gorrell	Title Manager, Registrations	Telephone No. (Include Area Code) (919) 549-2423
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Richard M. Gorrell</i>	3. Title Manager, Registrations	
4. Typed Name Richard M. Gorrell	5. Date August 7, 2009	

3 8 14



Bayer Environmental Science

August 7, 2009

EPA09RMG0807RonstarGT/LNotifContDisp
Fedex

Ms. Linda Arrington
U.S. Environmental Protection Agency
Office of Pesticide Programs (7504P)
Document Processing Desk (Notif)
Room S-4900, One Potmac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

RE: Ronstar G T/L Herbicide (EPA Reg No. 432-898) Notification to
Add Container Disposal Statement

Dear Ms. Arrington:

Bayer Environmental Science (BES) is notifying the Agency of the addition of a revised container disposal statement for the Ronstar G T/L Herbicide label. The label language is exactly as prescribed by the Agency in PR Notice 2007-4.

Bayer Environmental Science
2 T.W. Alexander Drive
RTP, NC 27709
Phone : 919.549.2000

The following information is included with this amendment:


- 8570-1 form
- Revised Label (4 paper copies; one highlighted copy)

If you have any questions, contact me at (919) 549-2423 or email me at mike.gorrell@bayercropscience.com.

Sincerely,

Richard M. Gorrell
Manager, Registrations

Cc: Karen Shearer



4 8 14

CHIPCO® RONSTAR® G T/L Herbicide

NOT FOR USE ON HOME LAWNS. FOR SALE TO, USE AND STORAGE BY COMMERCIAL TURF AND LANDSCAPE PERSONNEL ONLY. Not for sale to or use by homeowners/consumers.

Selective, preemergent herbicide for the control of annual grasses and broadleaf weeds in Turf and Woody Ornamental shrubs, vines, and trees.

ACTIVE INGREDIENT:
Oxadiazon [2-tert-butyl-4-(2,4 dichloro-5-isopropoxyphenyl)-?-1,3,4-oxadiazolin-5-one] 2.0%
INERT INGREDIENTS: 98.0%

EPA Reg. No. 432-898

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN
WARNING AVISO **NOTIFICATION**
SEP - 8 2009

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-331-2867

If in eyes:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Harmful if absorbed through the skin. Harmful if inhaled. Do not get in eyes or on clothing. Avoid contact with skin.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Loaders, applicators and other handlers must wear:
Long-sleeved shirt and long pants,
Shoes plus socks,
Chemical resistant gloves, and
Goggles, face shield or safety glasses.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Users should remove clothing/ PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. RONSTAR® should not be used in areas where irrigation or rainfall results in the direct or indirect (via ditches and canals) contamination of surface waters through dissolved runoff or erosion of soil particles to which oxidiazon is adsorbed.

DIRECTIONS FOR USE

**It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read entire label before using this product.**

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.
Do not apply CHIPCO® RONSTAR® G T/L through any type of irrigation system.

NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter treated area until sprays have dried.

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Do not contaminate water, food or feed by storage or disposal. Do not store in or around the home.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary land fill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

CHIPCO® RONSTAR® G T/L is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in turf and on a wide variety of woody ornamental shrubs, vines, and trees. CHIPCO® RONSTAR® G T/L controls weeds by killing the young weed seedlings as they come in contact with the herbicide during germination.

Apply CHIPCO® RONSTAR® G T/L with a properly calibrated granular applicator that will apply the granules uniformly (uniform application is essential for satisfactory weed control). Calibrate the applicator prior to use, according to the manufacturer's directions, and check frequently during the application to be certain that the equipment is working properly.

Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

TURF

For use in established perennial Bluegrass, Bentgrass, Bermudagrass, Buffalograss, Perennial Ryegrass, St. Augustinegrass, Seashore Paspalum, Tall Fescue, and Zoysia turf; such as, fairways, parks, golf courses, and lawns. Rake turf area free of leaves, mowing if necessary, so granules reach soil surface. Measure area to be treated and uniformly apply required amount, using a fertilizer spreader. Cross-treating with one-half required amount in each direction will give best results. **Do not** apply to wet turf. If rain is not expected shortly after application, sprinkling turf thoroughly will increase effectiveness.

PREEMERGENT WEED CONTROL

WEEDS CONTROLLED	AMOUNT OF CHIPCO® RONSTAR® G T/L		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass (<i>Digitaria spp.</i>) Goosegrass (<i>Eleusine indica</i>) Field Sandbur (<i>Cenchrus incertus</i>) Annual Sedge (<i>Cyperus compressus</i>)	100 - 200 lbs.	2 1/4 - 4 1/2 lbs.	Apply in late winter or early spring prior to seed germination. Application made after March or April may not be fully effective.
Annual Bluegrass (<i>Poa annua</i>)	100 - 200 lbs.	2 1/4 - 4 1/2 lbs.	Apply in late summer to early fall or in early spring prior to seed germination.
Virginia Buttonweed (<i>Diodia virginiana</i>) Pigweed (<i>Amaranthus spp.</i>) Florida Pusley (<i>Richardia scabra</i>) Oxalis (<i>Oxalis spp.</i>) Stinging Nettle (<i>Urtica dioica</i>) Carpetgrass (<i>Axonopus offinis</i>)	200 lbs.	4 1/2 lbs.	Apply in early spring prior to seed germination.

FALL APPLICATION FOR SPRING WEED CONTROL IN COOL SEASON TURF

WEEDS CONTROLLED	AMOUNT OF CHIPCO® RONSTAR® G T/L		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass, Goosegrass and other annual weeds	150 - 200 lbs.	3 3/8 - 4 1/2 lbs.	For use in Bluegrass, Bentgrass, Perennial Ryegrass and Tall Fescue turf. Make application in the fall after turf has become dormant. For late, fall applications, seeding into treated areas should be delayed until 4 months after greenup.

7 8.14

ANNUAL WEED CONTROL DURING BERMUDAGRASS, ZOYSIAGRASS, AND SEASHORE PASPALUM ESTABLISHMENT FROM SPRIGS

WEEDS CONTROLLED	AMOUNT OF CHIPCO® RONSTAR® G T/L		DIRECTIONS
	PER ACRE	PER 1000 SQ. FT.	
Crabgrass, Goosegrass, Annual Sedge, and other annual weeds	100 to 150 lbs.	2 1/4 to 3 3/8 lbs.	For Bermudagrass, make application immediately prior to or immediately after sprigging. For Seashore Paspalum, make application 10 to 14 days after sprigging. Applications made close to the time of sprigging may cause objectionable Seashore Paspalum injury. Disturbance of the soil surface after CHIPCO® RONSTAR® G T/L application may disrupt the herbicide barrier and reduce effectiveness. Irrigate immediately after application for best results.

For St. Augustinegrass and Bentgrass turf, do not exceed the rate of 150 pounds per acre (3 pounds active ingredient) of CHIPCO® RONSTAR® G T/L.

Do not apply to Bentgrass turf mowed at less than 3/8 inch.

CHIPCO® RONSTAR® G T/L may cause discoloration of Bentgrass, Bermudagrass and St. Augustinegrass. This discoloration is temporary and normally outgrown within 2 to 3 weeks.

Avoid contact with flowers and shrubs except as recommended elsewhere on this label. Do not apply CHIPCO® RONSTAR® G T/L to newly seeded areas. Seeding into treated areas should be delayed until 4 months after treatment. Do not apply to Dichondra or Centipedegrass. **CHIPCO® RONSTAR® G T/L should not be used on putting greens or tees because of the varying cultural practices used.**

WOODY ORNAMENTAL SHRUBS, VINES, AND TREES APPLICATION

Apply CHIPCO® RONSTAR® G T/L anytime during the year, prior to weed seed germination, for weed control. CHIPCO® RONSTAR® G T/L can be applied to both newly transplanted and established ornamentals. Application can be made to actively growing or dormant ornamentals. Rainfall or overhead irrigation after application of CHIPCO® RONSTAR® G T/L will improve weed control activity. CHIPCO® RONSTAR® G T/L should not be mixed into the soil. Cultivating or disturbing the soil may reduce the weed control activity of CHIPCO® RONSTAR® G T/L. **Do not** apply to wet foliage or under conditions in which granules will collect on leaves. **Do not apply in greenhouses** as plant injury may result from codistillation of CHIPCO® RONSTAR® active ingredient onto leaf surfaces in condensation.

AMOUNT TO USE

Apply CHIPCO® RONSTAR® G T/L broadcast to ornamentals at 100 to 200 pounds of product per acre (or 2 1/4 to 4 1/2 pounds per 1,000 sq. ft.) depending on the weeds to be controlled, as listed below. Remove existing weed growth before application.

Do not apply more than 6 pounds ai per acre per year (equivalent to 300 pounds of product /A/year), except in areas where there is heavy weed infestation. In areas of heavy weed infestation, apply no more than 8 pounds ai per acre per year (equivalent to 400 pounds of product/A/year).

Do not apply more than 4 pounds of active ingredient per acre per application (equivalent to 200 pounds of product per application).

ANNUAL BROADLEAF WEEDS		ANNUAL GRASSES	
Bittercress	(<i>Cardamine spp.</i>)	Annual Bluegrass	(<i>Poa annua</i>)
Bristly Oxtongue	(<i>Picris echioides</i>)	Annual Sedge	(<i>Cyperus compressus</i>)
Carpetweed	(<i>Mollugo verticillata</i>)	Barnyardgrass*	(<i>Echinochloa crus-galli</i>)
Cheeseweed	(<i>Malva parviflora</i>)	Bristly Foxtail	(<i>Setaria verticillata</i>)
Common Groundsel	(<i>Senecio vulgaris</i>)	Crabgrass	(<i>Digitaria spp.</i>)
Common Purslane	(<i>Portulaca olreacea</i>)	Fall Panicum	(<i>Panicum dichotomiflorum</i>)
Evening Primrose*	(<i>Oenothera spp.</i>)	Green Foxtail	(<i>Setaria viridis</i>)
Fiddleneck	(<i>Amsinckia intermedia</i>)	Goosegrass	(<i>Eleusine indica</i>)
Fireweed	(<i>Epilobium angustifolium</i>)	Jungle Ricegrass**	(<i>Echinochloa colonum</i>)
Galinsoga*	(<i>Galinsoga spp.</i>)	Panama Paspalum**	(<i>Paspalum fimbriaatum</i>)
Garden Spurge	(<i>Euphorbia hirta</i>)	Ripgut Bromegrass	(<i>Bromus catharticus</i>)
Golden Ragwort	(<i>Senecio aureus</i>)	Wild Oats	(<i>Avena fatua</i>)
Lambsquarters	(<i>Chenopodium album</i>)		
Liverwort**	(<i>Marchantia spp.</i>)		
Niruri	(<i>Phyllanthus niruri</i>)		
Pennsylvania Smartweed	(<i>Polygonum pennsylvanicum</i>)		
Petty Spurge	(<i>Euphorbia peplus</i>)		
Prostrate Spurge**	(<i>Euphorbia supina</i>)		
Redroot Pigweed	(<i>Amaranthus retroflexus</i>)		
Shepherdspurse	(<i>Capsella bursa-pastoris</i>)		
Sensitive Plant	(<i>Mimosa pudica</i>)		
Sow Thistle	(<i>Sonchus oleraceus</i>)		
Speedwell	(<i>Veronica spp</i>)		
Spiny Amaranth	(<i>Amaranthus spinosus</i>)		
Spotted Catsear	(<i>Hypochaeris radicata</i>)		
Swinecress*	(<i>Coronopus didymus</i>)		
Yellow Woodsorrel	(<i>Oxalis stricta</i>)		

*Use 150 to 200 pounds (3.0 to 4.0 pounds of active ingredient) per acre for less susceptible weeds.

**Use 200 pounds (4.0 pounds of active ingredient) per acre for less susceptible weeds.

How To Calibrate Spreaders For Applications Of CHIPCO® RONSTAR®

Spreader Models	Ground Speed (MPH)	Rate RONSTAR® G (Pounds Per Acre) Swath Width			(Feet)
		100	150	200	
CYCLONE B3-Pro	3	Spreader Settings			8
		2.75	3.25	3.9	
LELY HR*	3	Spreader Settings			
	4	3.25	3.75	4.25	
	5	3.5	4.25	4.5	
		3.75	4.5	5.0	21
LESCO ROTARY	3	Spreader Settings			7
		C	D	E	
SCOTT'S PRO R-8A	3	Spreader Settings			9
		H	I+	J+	
SPYKER 24-22	3	Spreader Settings			8
		3.5	4.0	4.25	
VICON	3	Spreader Settings			
	4	13	15	18	
	5	15	18	22	
		16	21	25	22

* Settings based on use of small seed feeding with central spreader setting

NOTE: These settings are for initial calibration only. It is strongly recommended that your equipment be calibrated prior to actual application, using these settings as a starting point.

PLANT TOLERANCE

Plant tolerances to this product have been found to be acceptable in the specific genera and species on this label. It is not possible to evaluate every species or variety of ornamental plant for its tolerance to this product. The user should test for possible phytotoxic responses in other plants on a small area basis using recommended rates prior to commercial use.

ESTABLISHED ORNAMENTALS AND GROUND COVERS

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Abutilon	<i>Abutilon pictum</i>	Citrus, Ornamental*:	
Acacia	<i>Acacia spp</i>	Grapefruit	<i>Citrus paradisi</i>
Agapanthus	<i>Agapanthus africanus</i>	Lemon	<i>Citrus limon</i>
Allamanda, Common	<i>Allamanda cathartica</i>	Lime	<i>Citrus aurantifolia</i> Bearss
Andromeda	<i>Pieris spp</i>	Mandarin	<i>Citrus reticulata</i>
Aralia	<i>Aralia spinosa</i>	Tangelo	<i>Citrus XTangelo Minneola</i>
Arborvitae	<i>Thuja spp</i>	Tangerine	<i>Citrus reticulata var. tangerina</i>
Ash, Green	<i>Fraxinus pennsylvanica</i>	Temple	<i>Citrus Xnobilis Temple</i>
Azalea	<i>Rhododendron spp</i>	Cuban Shaddock	<i>Citrus maxima</i>
Bamboo, Feathery	<i>Bambusa vulgaris</i>	Valencia	<i>Citrus sinensis</i>
Banyan, Weeping	<i>Ficus benghalensis</i>	Washington Navel	<i>Citrus sinensis</i>
Barberry	<i>Berberis spp</i>	Coromandel	<i>Asystasia gangentica</i>
Beafsteak plant	<i>Acalypha wilkesiana</i>	Cotoneaster	<i>Cotoneaster spp</i>
Bearberry	<i>Arctostaphylos uva-ursi</i>	Crabapple*	<i>Malus spp</i>
Beech, European	<i>Fagus sylvatica</i>	Crapemyrtle	<i>Lagerstroemia indica</i>
Birch, Paper	<i>Betula papyrifera</i>	Chrysanthemum	<i>Chrysanthemum spp</i>
Boxwood, Japanese	<i>Buxus microphylla</i>	Cypress	<i>Cupressus spp</i>
Boxwood, Common	<i>Buxus sempervirens</i>	Cypress, Leyland	<i>Cupressocyparis leylandii</i>
Bouganvillea	<i>Bouganvillea spectabilis</i>	Daphne, Fragrant	<i>Daphne odora</i>
Brassaia, Dwarf	<i>Brassaia actinophylla</i>	Deutzia, Slender	<i>Deutzia gracilis</i>
Buddleia	<i>Buddleia davidii</i>	Dieffenbachia	<i>Dieffenbachia spp</i>
Bugleflower	<i>Ajuga reptans</i>	Dogwood, Common	<i>Cornus florida</i>
Callistemon	<i>Callistemon lanceolatus</i>	Dogwood, Redosier	<i>Cornus stolonifera</i>
Camellia, Common, Japanese	<i>Camellia japonica</i>	Dogwood, Koussa	<i>Cornus koussa</i>
Camellia, Sasanqua	<i>Camellia sasanqua</i>	Douglas fir	<i>Pseudotsuga menziesii</i>
Candytuft	<i>Iberis sempervirens</i>	Elm	<i>Ulmus spp</i>
Carissa	<i>Carissa macrocarpa</i>	Escallonia	<i>Escallonia spp</i>
Carnation	<i>Dianthus caryophyllus</i>	Eucalyptus	<i>Eucalyptus spp</i>
Ceanothus	<i>Ceanothus americanus</i>	Euryops, Gray Leaved	<i>Euryops pectinatus</i>
Cedar, Red	<i>Juniperus virginia</i>	Euonymus, Spreading	<i>Euonymus kiautshovicus</i>
Cedar, Atlas	<i>Cedrus atlantica</i>	Euonymus, Wintercreeper	<i>Euonymus fortunei</i>
Cistus	<i>Cistus purpureus</i>	Falsecypress, Hinoki	<i>Chaemaecyparis obtusa</i>
		Falsecypress, Japanese	<i>Chaemaecyparis pisifera</i>
		Flowering Quince	<i>Chaenomeles speciosa</i>
		Forsythia	<i>Forsythia intermedia</i>
		Gardenia	<i>Gardenia jasminoides</i>
		Gazania, Trailing	<i>Gazania uniflora leucoleana</i>
		Ginko	<i>Ginko biloba</i>
		Heath, Spring	<i>Erica carnea</i>
		Heather, Scotch	<i>Calluna vulgaris</i>

ESTABLISHED ORNAMENTALS AND GROUND COVERS (con't)

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Russian Olive	<i>Eleagnus angustifolia</i>	Syzygium	<i>Syzygium paniculatum</i>
Scotch Broom	<i>Cytisus scoparius</i>	Trailing African Daisy	<i>Osteospermum fruticosum</i>
Sedum	<i>Sedum album</i>	Viburnum	<i>Viburnum spp</i>
Serviceberry, Saskatoon*	<i>Amelanchier alnifolia</i>	Wedelia	<i>Wedelia trilobata</i>
Spruce	<i>Picea spp</i>	Witchhazel, Common	<i>Hamamelis virginiana</i>
St. Johnswort, Golden	<i>Hypericum frondosum</i>	Woadwaxen, Common	<i>Genista tinctoria</i>
St. Johnswort, Gokdencup	<i>Hypericum patulum</i>	Xylosma	<i>Xylosma congestum</i>
Strawberry Tree	<i>Arbutus unedo</i>	Yew	<i>Taxus spp</i>
Sumac, Staghorn	<i>Rhus typhina</i>		

NEWLY TRANSPLANTED ORNAMENTALS AND GROUND COVERS

<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>	<u>COMMON NAME</u>	<u>SCIENTIFIC NAME</u>
Brassaia, Dwarf	<i>Brassia actinophylla</i>	Privet	<i>Ligustrum spp</i>
Camellia	<i>Camellia japonica</i>	Protea Species	<i>Banksia menziesii</i> <i>Leucadendron discolor</i> <i>Leucospermum cordifolium</i> <i>Protea exinia</i>
Carnation	<i>Dianthus caryophyllus</i>	Rhododendron	<i>Rhododendron spp</i>
Cedar, Red	<i>Juniperus atlantica</i>	Russian Olive	<i>Eleagnus angustifolia</i>
Coromandel	<i>Asystasia gangetica</i>	Spruce	<i>Picea spp</i>
Dogwood, Flowering	<i>Cornus florida</i>	Trailing African Daisy	<i>Osteospermum fruticosum</i>
Elm	<i>Ulmus spp</i>	Wedelia	<i>Wedelia trilobata</i>
Forsythia	<i>Forsythia intermedia</i>	Yew	<i>Taxus spp</i>
Gazania, Trailing	<i>Gazania uniflora leucoleana</i>		
Hackberry	<i>Celtis occidentalis</i>		
Ivy, Algerian	<i>Hedera canariensis</i>		
Ivy, English	<i>Hedera helix</i>		
Hemlock, Canada	<i>Tsuga canadensis</i>		
Holly	<i>Ilex spp</i>		
Ice Plant	<i>Carpobrotus edule</i>		
Juniper	<i>Juniperus spp</i>		
Metallic Plant	<i>Hemigraphis colorata</i>		
Oak	<i>Quercus spp</i>		
Periwinkle, Common	<i>Vinca minor</i>		
Periwinkle, Large	<i>Vinca major</i>		
Pine	<i>Pinus spp</i>		
Poplar	<i>Populus spp</i>		

SENSITIVE VARIETIES

Under certain growing conditions, the plant varieties listed below may be sensitive to CHIPCO® RONSTAR® G T/L Herbicide. Growers should evaluate usage on a small scale prior to large scale applications.

<u>PLANT SPECIES</u>	<u>VARIETIES</u>
Andromeda (<i>Pieris japonica</i>)	Snowdrift, Temple Bell
Azalea	Pink Champagne, Hershey, (<i>Rhododendron spp</i>), Leuchtenfuer, Memoire John Hearrens, Hino-Crimson, White Grandeur, Everest, Girard Roberta
Cotoneaster	Eichholz, Himalayan, Shangri-la (<i>Cotoneaster spp</i>)
Crapemyrtle (<i>Lagerstroemia indica</i>)	Petite Red Imp, Seminole Bush
Heath, Spring (<i>Erica carnea</i>)	Mediterranean Pink
Hemlock, Canada (<i>Tsuga canadensis</i>)	Pendula
Hibiscus (<i>Hibiscus syriacus</i>)	Red Heart
Rhododendron	Daphnoides, Nova Zembla, (<i>Rhododendron spp</i>), Catawba Album
Spruce, Norway (<i>Picea abies</i>)	Pumila
Spruce, White (<i>Picea glauca</i>)	Densata
Viburnum (<i>Viburnum spp</i>)	Davidii

* For use only on plants that will not bear edible fruit within one year of CHIPCO® RONSTAR® G T/L Herbicide application.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

CHIPCO® and RONSTAR® are registered trademarks of Bayer.



Bayer Environmental Science

A Business Group of Bayer CropScience LP
PO Box 12014 2 T. W. Alexander Drive
Research Triangle Park, NC 27709

CHIPCO® RONSTAR® G T/L (MASTER) 07/31/07, Notification 08/06/09