


**PERMANONE® MULTI-PURPOSE 10% E.C.**

- \* RECOMMENDED FOR AGRICULTURAL/COMMERCIAL USE
- \* BROAD SPECTRUM MULTI-USE INSECTICIDE
- \* QUICK KNOCKDOWN
- \* FOR USE OUTDOORS AND IN HOMES, KENNELS, AND NON-FOOD/FEED AREAS OF COMMERCIAL BUILDINGS, INSTITUTIONS, WAREHOUSES, THEATERS, OFFICE BUILDINGS, SCHOOLS, MOTELS, HOTELS, RESTAURANTS AND FOOD/FEED PROCESSING ESTABLISHMENTS.
- \* FOR CONTROL OF FLEAS, BROWN DOG TICKS ON PREMISES
- \* FOR USE ON LIVESTOCK: CATTLE (BEEF AND DAIRY), GOATS, SHEEP, SWINE, HORSES AND POULTRY

**ACTIVE INGREDIENT:**

\*Permethrin: ..... 10.0%  
 †OTHER INGREDIENTS: ..... 90.0%  
 100.0%

\*Cis/trans ratio: Max. 65% (+/-) trans and min. 35% (+/-) cis.

†Contains Petroleum Distillates

**EPA Reg. No. 432-1132**

**EPA Est. No.**

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

<b>FIRST AID</b>	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give any liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER: 1-800-331-2867</b> [Synthetic Pyrethroid]	

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

**Note To Physician:** Do not administer or instill milk, cream or other substances containing vegetable or animal fats which enhance absorption of lipophilic substances.

**See Side Panel For Additional Precautionary Statements.**

#### **NET CONTENTS:**

#### **READ ENTIRE LABEL BEFORE EACH USE**

#### **PRECAUTIONARY STATEMENTS** **Hazards To Humans & Domestic Animals**

##### **WARNING**

Causes skin irritation. Do not get on skin or on clothing. Wear coveralls worn over short-sleeved shirt and short pants, socks, chemical resistant foot wear and gloves when handling or mixing concentrate. Harmful if inhaled or absorbed through skin. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

##### **Environmental Hazards**

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when wind speeds exceed 10 mph. Do not contaminate water when disposing of equipment washwaters. Shrimp and crab may be killed if application rates recommended on this label are exceeded. Use with caution where these are important resources. This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply this product or allow drift when bees are actively visiting the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

##### **Physical & Chemical Hazards**

Do not use or store near heat or open flame. Do not use this product in or on electrical equipment due to the possibility of shock hazard.

#### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To prepare dilutions, the concentrate should first be stirred or agitated well. Add the required amount of concentrate to water or oil and blend thoroughly. Do not hold dilutions for more than 24 hours.

For maximum effectiveness, a combination of localized application and space treatment is recommended. Remove pets, birds and cover fish aquaria before spraying.

Do not allow spray treatment to contaminate pasture land, cropland, poultry ranges or water supplies. Do not use on crops used for food, forage or pasture.

{Alternate Text For Labels with Indoor Uses: Do not use in food/feed areas of food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared food is served such as dining rooms, but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food/feed should be covered or removed.}

{Alternate Text for Labels with Indoor Uses: NON FOOD/FEED AREAS include (but are not limited to) garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).}

{Alternate Text for Labels with Indoor Uses: Do not apply to classrooms when in use. Do not apply this product in patient rooms or in any rooms when occupied by the elderly or infirm.}

Do not allow spray treatment to contaminate pasture land, cropland, poultry ranges or water supplies. Do not use on crops used for food, forage or pasture.

#### INDOOR USE AREAS

Bakeries	Kennels
Beverage Plants	Meat Packing Plants
Canneries	Motels
Flour Mills	Office Buildings
Food Processing Plants	Railroad Cars
Grain Elevators	Restaurants
Granaries	Schools
Homes	Ships' Holds
Hospitals (Non-Patient Areas)	Supermarkets
Hotels	Truck Trailers
Industrial Installations	Warehouses

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage And Spill Procedures: Store upright at room temperature.

Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste. Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved State and local procedures.

CONTAINERS ONE GALLON AND SMALLER: Do not reuse container. Wrap container in several layers of newspaper and discard in trash.

CONTAINERS LARGER THAN ONE GALLON: Metal Containers – Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers - Triple rinse or equivalent, then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke. Glass Containers - Triple rinse or equivalent, then dispose of in a sanitary landfill or by other approved state and local procedures.

**SEE REVERSE FOR SPECIFIC USE INSTRUCTIONS**

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer Environmental Science. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG, 2002

PERMANONE is a registered trademark of Bayer.

Bayer Environmental Science  
A Business Group of Bayer CropScience LP  
95 Chestnut Ridge Road  
Montvale, NJ 07645

APPLICATION SITES	PESTS CONTROLLED	USE DIRECTIONS	DILUENT	DILUTION RATE	APPLICATION RATE	DOSE RATE
AS SURFACE SPRAY Bakeries, Beverage Plants, Canneries, Flour Mills, Food Processing Plants, Grain Elevators, Granaries, Homes, Hospitals, Hotels, Industrial Installations, Kennels, Meat Packaging Plants, Motels, Office Buildings, Railroad Cars, Restaurants, Schools, Ships' Holds, Supermarkets, Truck Trailers and Warehouses.	Cockroaches, waterbugs, Palmetto Bugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Millipedes, Centipedes, Sowbugs, Pillbugs, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Red Rust Flour Beetles, Sawtoothed Grain Beetles, Cigarette Beetles, Drugstore Beetles, Mealworms, Grain Mites and Cadelles	For initial clean up of severe insect infestations: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets around containers of stored foods, around the base of machinery behind shelves and drawers. Spray bookcases for Silverfish. Spray ant trails, nests and points of entry.  For normal infestations: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into the hiding places, cracks and crevices under pallets, around containers of stored foods, around the base of machinery and behind shelves and drawers. Spray bookcases for Silverfish. Spray ant trails, nests and points of entry.	Water	1 part concentrate in 10 parts water (13 fl. ounces per gallon)	1 gallon to 750 square feet of surface	
	Carpet Beetles	Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets, and infested area of shelving.	Water	1 part concentrate in 20 parts water (6.5 fl. ounces per gallon)	1 gallon to 750 square feet of surface	
	Bedbugs	Spray mattresses lightly particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, floors. Allow all treated articles to thoroughly dry before use.	Water	1 part concentrate in 20 parts water (6.5 fl. ounces per gallon)		
	Fleas and Ticks (Adults and Larvae)	Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment.	Water	From 1 to 2 parts concentrate to 40 parts water (3.25 - 6.50 fl. ounces per gallon)		
	Walls, ceilings, moldings, screens, door frames, beams, light cords, and similar resting places	Treat surfaces using coarse wet spray. Spray surfaces thoroughly, but do not allow runoff to occur.	Water	1 part concentrate in 20 parts water (6.5 fl. ounces per gallon)		
LIVESTOCK SPRAY: Dairy & Beef Cattle, Sheep, Goats, Swine, Horses, and Poultry (For swine, do not ship animals for slaughter	Horn Flies, House Flies, Mosquitoes, and Gnats	Apply to wet the hair thoroughly, with particular attention to topline, underline, flanks, withers and other infested areas. Repeat treatment at intervals of 5 to 12 days for small insect populations or as needed when flies are emerging in large numbers.	Water	1 part concentrate in 100 parts water (1.3 fl ounces per gallon)		

5/8

animals for slaughter within 5 days of last treatment).

	Stable Flies, Horse Flies, and Deer Flies	Apply a quart per adult animal to wet the hair thoroughly with particular attention to the legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat daily as needed.	Water	1 part concentrate in 100 parts water (1.3 fl ounces per gallon)		
	Face Flies	Apply using spray which produces large wetting droplets. Apply to the face of the animal in the morning before release to pasture. Apply sufficiently to wet the face but not more than 1.5 ounces per animal. Repeat daily as needed.	Water	1 part concentrate in 100 parts water (1.3 fl ounces per gallon)		
	Lice (Including Poultry Lice) Northern Fowl Mites, Mange	Spray roots, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist. For swine, thoroughly soak animal, including ears. Retreat at 14 days for Mange. Treat walls and bedding.	Water	1 part concentrate in 100 parts water (1.3 fl ounces per gallon)		
	Bedbugs and Mites in Poultry Houses	Spray crevices of roost poles, cracks in walls and cracks in nests where the bedbugs and mites hide.	Water	1 part concentrate in 100 parts water (1.3 fl ounces per gallon)		
	Sheep "ticks" or keds	Thoroughly wet all portions of the body by dipping or by spraying with sufficient pressure and with a nozzle adjustment to give penetration of the wool. Treat at a rate sufficient to wet the animal	Water	1 part concentrate in 100 parts water (1.3 fl ounces per gallon)		
	Fleas on Livestock (To kill and to protect against reinfestation)	Wet the animal by dipping or spraying	Water	1 part concentrate in 20 parts water (6.5 fl ounces per gallon)		
	Horn Flies, Face Flies, Stable Flies	Back rubber/Self Oiler (Keep rubbing device charged). Results can be improved by forced use.		1 pint concentrate in 10 parts diesel oil		
AS A SPACE SPRAY LIVESTOCK PREMISES: Barns, Milking Parlors, Milk Rooms, Dairies, Poultry Houses, Swine and Livestock Housing	Fleas Mosquitoes and Gnats	Apply as a fog or fine mist, directing the spray toward the ceiling and upper corners until the area is filled with mist, using about 2 ounces per 1000 cubic feet of space. For best results, close doors and windows before spraying and keep them closed for ten to fifteen minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment daily as needed.	Water	1 part concentrate in 20 parts water (6.5 fl ounces per gallon)		

<p>Bakeries, Beverage Plants, Canneries, Flour Mills, Food Processing Plants, Granaries, Homes, Hospitals, Hotels, Industrial Installations, Kennels, Meat Packaging Plants, Motels, Office Buildings, Railroad Cars, Restaurants, Schools, Ships' Holds, Supermarkets, Truck Trailers, Warehouses</p>	<p>House Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black Flies, Small Flying Moths.</p>	<p>Apply with mechanical or compressed air equipment (non-thermal) adjusted to deliver a fine mist. Close doors and windows and shut off ventilating systems. When using an oil dilution for space spraying extinguish all flames and pilot lights when applying.</p> <p>For rapid control of Houseflies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Bees, Black flies, Small Flying Moths, direct spray at an upward angle distributing uniformly through the entire area at a rate of 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate areas after treatment and ventilate and sweep up dead insects before reoccupying.</p> <p>For rapid kill of exposed or accessible stages of other insects named on this label, apply using conventional mechanical or compressed air equipment (non-thermal) following directions for space spraying.</p>	<p>Oil or Water</p>	<p>1 part concentrate in 20 parts oil or water (6.5 fl. ounces per gallon)</p>	<p>1 ounce per 1000 cubic feet</p>	
<p>Animal Houses, Corrals</p>	<p>Mosquitoes, Gnats, biting and non-biting Midges, Black Flies, and other biting Flies.</p>	<p>Apply using vehicle mounted ULV equipment to create an insecticidal swath. For best results, apply when there is a light breeze (i.e., about 5 mph) and apply in direction of breeze to obtain maximum swath and better distribution. Do not apply if winds exceed 10 mph.</p> <p>The target dose rate to be applied is up to 0.007 lb a.i. per acre. This can be achieved under different conditions by altering the flow of insecticide from the equipment and/or the speed of the vehicle.</p> <p>Apply 3.5 to 7.0 fluid ounces per minute at vehicle speeds of 5-10 miles per hour using swath widths of 150-300 feet as appropriate to local conditions.</p> <p>At higher vehicle speeds, flow rates should be increased proportionately to achieve the target dose rate.</p> <p>Not to be used within 100 feet (30 meters) of lakes and streams.</p> <p>In treatment of corrals, feed lots and swine lots, cover any exposed drinking water, drinking fountains and animal feed before application.</p> <p>Allow 24 hours before retreating. Do not apply more than once in a 24-hour period.</p>	<p>Oil or Water</p>	<p>1 part concentrate with 1 part suitable oil solvent or at proportional oil dilutions calculated on the basis of vehicle speed and swath width to achieve the target dose.</p>	<p>Dependent on swath width, flow rate and speed of vehicle</p>	<p>Up to 0.007 lbs of active ingredient per acre</p>

<b>BARRIER SPRAY</b> Perimeter of residential yards, public facilities, paths and recreational areas.	As an aid in reducing mosquito annoyance. To kill and control biting and non-biting midges, deer flies and other biting flies	Apply when walking at a speed of approximately 2 mph and a swath of 50 feet treating approximately 0.2 acre/minute.	Oil or Water	1 part concentrate to 1 part of a suitable diluent, or at proportional dilutions calculated on the basis of applicator speed and swath width to achieve the target dose	Apply 36.6 fluid ounces of diluted material per acre.	Approximately 0.1 lb per acre
--	---	---	--------------	---	---	-------------------------------

**HOSE-END SPRAYER:** To control listed insects, shake well before applying. Connect nozzle to hose; remove cap, insert diptube and screw bottle onto nozzle. Turn water at the tap. When ready to treat, place finger over the hole on right side of nozzle and the liquid will mix automatically at the proper ratio. Apply to surfaces in dairies, barns, feed lots, stable, poultry houses, swine and livestock housing: to control House Flies, Stable Flies and other manure breeding flies. Also aids in control of Cockroaches, Ants, Spiders, Mosquitoes, Crickets and Face Flies. Walk and spray at a steady pace. Spray walls and surfaces thoroughly but do not allow runoff to occur. Treat no more than once every two weeks. Do not contaminate animal feed or water by spray.

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Bayer Environmental Science's control, it is impossible for Bayer Environmental Science to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATION OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG 2002

Permanone is a registered trademark of Bayer

**BAYER ENVIRONMENTAL SCIENCE**

A Business Group of Bayer CropScience LP  
95 Chestnut Ridge Road, Montvale, NJ 07645