

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

December 2, 2010

Gregory Mattern
Bayer Environmental Science
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Subject:

Application for Pesticide Notification (PRN 98-10)

Optional Marketing Claims

✓ CP Carbaryl Insecticide Spray Ready-To-Use, EPA Reg. No. 432-1211 Sevin Grub Killer Granules, EPA Reg. No. 432-1212 R&M Garden Dust 5%, EPA Reg. No. 432-1209 Application Dated November 8, 2010

Dear Mr. Mattern;

The Agency is in receipt of your filling of a revised Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6736 or electronically at Edwards.joanne@epa.gov.

Sincerely,

Joanne S. Edwards

Insecticide-Rodenticide Branch Registration Division (7505P)

Office of Pesticide Programs

Please read instructions	n reverse before comple	ving form.			Form Apr	oroved	OMB No	. 2070-00	80. Approval expires 2-28-9
United States Environmental Protection Washington, DC 204						✓	Regist Amend Other		OPP Identifier Number
		Application	on for	Pesticid	e - Sec	tion	l		
1. Company/Product Number 432-1211			2. EPA Product Manager Joanne Edwards			3. P	3. Proposed Classification None Restricted		
4. Company/Product (Name) CP Carbaryl Insecticide Spray Ready-To-Use			PM# 7						
5. Name and Address of Applicant (Include ZIP Code)			6. Expedited Reveiw. In accordance with FIFRA Section 3(c)(3)						
Bayer Environmental Science P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, NC 27709			(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No.						
Check if	this is a new address			Produc	t Name				
			Se	ction - II					
Amendment - Exp		r deted			Final printed Agency lett	ter dat	ted		OTIFICATION
Resubmission in response to Agency letter dated Notification - Explain below.				"Me Too" Application. Other - Explain below.			DEC 5 - 5010		
no other changes have be	een made to the labeling of false statement to EPA. I	or the confidenti further unders	al statemo tand that i pject to en	ent of formul f this notifica	a of this prod tion is not co ction and pe	duct. I onsiste	understand ent with the	that it is a vector terms of PR	Notice 98-10 and 40 CFR
1. Material This Product	Will Be Packaged In:								
Child-Resistant Packaging Unit Packaging			Water Soluble Packaging 2. Type o			of Containe	or .		
Yes	Yes		Yes			Metal	Metal Plastic		
✓ No	✓ No		✓	✓ No			Glass		
* Certification must be submitted	If "Yes" Unit Packaging wgt	No. per container	If "Ye Packs	es" age wgt	No. per containe	r		Paper Other	(Specify)
3. Location of Net Conter	nts Information	4. Size(s) Re	tail Cont	ainer	1	5. Lo	cation of L	abel Direct	ions
✓ Label or ✓ Container		Various (up to 2 gal) Cor			Conta	ontainer label / booklet			
6. Manner in Which Label is Affixed to Product J Lithograph or Other Other									
			Sec	tion - IV	/				
1. Contact Point (Compl	ete items directly below	for identificati	on of indi	ividual to be	contacted,	if nec	essary, to	process th	is application.)
Name Gregory C. Mattern			Title Regulatory Product Manager			1	Telephone No. (krolude Area Code) (919) 549-2630		
	atements I have made o any knowlinglly false o ole law.		d all attac				urate and o imprisonm		6. Date Application Received
2. Signature Gregory C. Mattern					- C C C				
4. Typed Name			5. Date						
Gregory C. Mattern		November 8, 2010					(((((((((((((((((((

CP CARBARYL INSECTICIDE SPRAY Ready-to-Use

[FOR OUTDOOR RESIDENTIAL USE ONLY]

[Ready to Use]

[Shake Well and Spray]

[Easy to Apply]

[Kills Japanese Beetles]

[Kills over 100 Insect Pests: 48 Different Insects on Vegetable Crops]

[74 Different Insects on Fruit Crops]

[80 Different Insects on Trees, Ornamentals, Shrubs, Flowers]

NOTIFICATION

roses, and flowers)

Got Japanese Beetles? Get Sevin!

Got [insert listed pest here]? Get Sevin!

Hate The Bugs. Love The Garden. Get Sevin!

Carbaryl*..... OTHER INGREDIENTS.....

EPA Reg. No.432-1211

DEC 2 - 2010 99.874%

Total..... 100.000%

[12 different Insects Around the outside of Homes]

For use on fruits, trees, vegetable gardens, ornamentals, shrubs,

*CAS Number 63-25-2

ACTIVE INGREDIENT:

EPA Est.67572-GA-1

KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-XXX-XXX-XXXX

	FIRST AID
This	product is a N-methyl carbamate and is a cholinesterase inhibitor
If Swallowed	Call poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow
	Do not induce vomiting unless told to do so by the poison control center or doctor.
<u> </u>	Do not give anything by mouth to an unconscious person.
lf on Skin or Clothing	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15-20 minutes.
	Call a poison control center or doctor for treatment advice.
If in Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-334-7577 for emergency treatment information.

NOTE TO PHYSICIAN: CARBARYL is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are not recommended.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. For overhead applications, also wear a hat and eye protection. Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use. Wash the outside of gloves before removing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or allow drift to water. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

BEE CAUTION: MAY KILL HONEYBEES AND OTHER BEES. This product is highly toxic to honeybees and other bees. Do not apply to plants in bloom.

[Optional Back Panel claims statement(s)]:

This product is ready to use and easy to apply. Simply shake well and spray. This product kills over 100 insect pests and can be used on vegetable gardens, fruits, trees, ornamentals, shrubs, flowers, and around the outside of homes.

QUICK FACTS Read the Label First!	
KILLS	Japanese Beetles, Imported Cabbageworms, Tomato Hormworms, Ants, Ticks and over 100 other insect pests.
WHERE TO USE	On vegetables, fruits, trees, shrubs, ornamentals, flowers, and around the outside of homes.
[Phone Symbol]	For questions or comments, call toll-free 1-877-BAYER-AG (1-877-229-3724).

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read the entire label before using this product. Strictly observe label direction, precautions, and restrictions.

Restrictions

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

For outdoor use only. Not for use on plants grown for sale, commercial use, research, or for commercial seed production. Not for use on lawns.

STORAGE AND DISPOSAL

Storage: Store unused product in an area out of reach of children and animals. Do not store in areas where temperatures frequently exceed 100°F.

Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-XXX-XXXX for disposal instructions. Never place unused product down any indoor or outdoor drain.

[Optional Sprayer Directions]

Option 1:

How to Attach Sprayer to Bottle: Remove Power Sprayer and hose from side carrier. Replace bottle cap with cap attached to Power Sprayer hose.

Prepare to Spray: Point Sprayer away from body. Adjust sprayer to desired spray pattern (e.g., stream setting for directly spraying pest or wide setting for treating larger areas), do not turn past "X" position. Slide safety switch to ON position.

Spray: Squeeze trigger to dispense product. Sprayer may require several seconds to prime before spraying.

After Use: Slide safety switch to LOCK position. Turn sprayer nozzle to OFF position. Return Power Sprayer to side carrier for storage.

Option 2:

How to Attach Sprayer to Bottle: Remove Power Sprayer from side carrier. Pull battery tab to remove. Fully extend hose from bottom of sprayer. Open flip-cap on bottle. Insert hose and plug into flip-cap opening until it snaps into place.

How to Spray: Point Power Sprayer away from body. Adjust sprayer to desired spray pattern (e.g., stream setting for directly spraying pest or wide setting for treating larger areas). Squeeze trigger to dispense product. Sprayer may require several seconds to prime before spraying.

After Use: Turn sprayer nozzle to X (off) position. Do not unplug hose. Return Power Sprayer to side carrier for storage. Do not reuse Power Sprayer with any other product or on any other container.

Proper Use of Batteries: Never mix old batteries with new. Do not mix Alkaline with Standard (carbon zinc) or Rechargeable (nickel cadmium) batteries. When replacing batteries, follow diagram inside battery compartment. When storing sprayer at end of season, remove batteries to prevent leakage. Follow manufacturer's recommendations for disposal of batteries.

[HOW TO USE] SHAKE CONTAINER BEFORE USING

Adjust sprayer to desired spray pattern (e.g., stream setting for directly spraying pest or wide setting for treating larger areas). Hold sprayer 8 to 12 inches from the area to be treated. Spray until surface is wet. Do not apply when temperatures exceed 85°F as temporary leaf burn may occur. Start spraying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Do not use on large trees. For trees taller than 10 feet consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

VEGETABLE CROPS

To effectively kill the vegetable pests listed below, apply this product in adequate volume to effectively cover both the upper and lower surfaces of the plant.

PHI refers to the minimum interval in days that must be observed between the date of the last application and date of harvest. APP refers to the maximum number of applications per crop per year.

Vegetable	PHI	APP
Asparagus	1	3
Brassica head and stem vegetables*: Broccoli, Brussels sprouts, Cabbage, Cauliflower, Chinese cabbage, Kohlrabi	3	4
Brassica leafy green vegetables*: Chinese cabbage, Collards, Kale, Mustard greens, Mustard spinach, Turnip greens	14	4
Corn (sweet)	2	8
Cucurbit vegetables: Cucumber, Melons, Pumpkin, Squash	3	6
Dried shelled legume vegetables**: Adzuki bean, Blackeyed pea, Cowpea, Field bean, Kidney bean, Lentils, Lima been (dry), Mung bean, Navy bean, Pinto bean, Southern pea	21	4
Edible-podded legume vegetables**: Dwarf pea, Edible-pod pea, Jack bean, Snap bean, Snow pea, Sugar snap pea, Wax bean, Yardlong bean,	3	4
Fruiting vegetables: Eggplant, Groundcherry, Okra, Peppers, Tomatoes	3	7
Leafy vegetables: Cardoon, Celery, Celtuce, Florence fennel, Dandelion, Endive, Lettuce (head and leaf), Parsley, Rhubarb, Spinach, Swiss chard	14	5
Okra	3	4
Peanuts	14	5
Root and Tuber vegetables (except sugar beets and sweet potatoes): Artichoke (Chinese and Jerusalem), Carrots, Garden Beets (roots and tops), Ginger, Ginseng, Horseradish, Parsnip, Potato, Radish, Rutabega, Salsify, Turnip (roots and tops), Yams	7	6
Sweet Potatoes	7	8 .

PESTS

Alfalfa Caterpillar, Apache cicada, Armyworms, Asparagus beetle, Aster leafhopper, Bean leaf beetle, Chinch bug, Colorado potato beetle, Corn earworm, Corn rootworm Adult, Cowpea curculio, Cucumber beetle, Cutworms, European corn borer, Fall armyworm, Flea beetles, Grasshoppers, Green Cloverworm, Harlequin bug, Imported cabbageworm, Japanese beetle, Lace bugs, Leafhoppers, Lima Bean pod Borer, Lygus bugs, Melon worm, Mexican bean beetle, Pea weevil, Rednecked peanutworm, Sap Beetles, Southwestern corn borer, Spittlebugs, Squash bugs, Stink bugs, Sweet potato weevil, Three cornered alfalfa hopper, Thrips, Tomato hornworm, Tomato pinworm, Tortoise beetle, Velvetbean caterpillar, Webworms, Western bean Cutworm, Whitefringed Beetle (adults), Yellowstriped Armyworm

* Application is only permitted within 30 days of crop emergence or the date or transplanting.

** Do not use on fresh/succulent peas and beans.

FRUIT CROPS

This product may be used as a spot treatment for localized pest infestations on the tree fruits listed below. Do not use on large trees or where pest populations are generally inaccessible to spray deposits, as inadequate performance will result. Direct spray toward upper and lower leaf surfaces and between fruit clusters and on small limbs and trunks to the point of runoff where pest appear.

Repeat applications as necessary up to a total of eight times for tree fruits or five times for small fruits, but not more often than once every seven days.

NOTE: To avoid undesired apple thinning delay use until at least 30 days after full bloom. Carbaryl insecticide may injure early dawn and sunrise strawberry varieties on the Delmarva Peninsula. The number in parenthesis following each fruit and nut crop grouping refers to the minimum interval that must be observed between the date of the last application and the date of harvest.

TREE FRUITS	PESTS		
Apples, Apricots, Cherries, Nectarines, Peaches, Pears, Plums, Prunes (3 days until harvest)	Apple aphid, Apple maggot, Apple mealybug, Apple pandemis, Apple rust mite, Apple sucker, Avocado Leafroller, Bagworms, Black Cherry Aphid, Black scale, Blueberry maggot, Brown soft scale, California red scale, Cherry fruitworm, Cherry maggot, Citrus cutworm, Citrus root weevil, Citricola scale, Citrus snow scale, Codling moth,		
FRUITS Blackberries, Blueberries, Dewberries, Grapes, Raspberries, Strawberries (7 days Until harvest)	Cucumber beetles, Cutworms, Eastern tent caterpillar, Eightspotted forester, European apple sawfly, European earwig, European raspberry aphid, Eyespotted budmoth, Forbes scale, Fruit tree leafroller, Grape berry moth, Grape leaffolder, Grape leafhopper, Green fruitworm, Japanese beetle, June beetles, Leafhoppers, Leafrollers, Lecanium scales, Lesser appleworm, Lesser peachtree borer, Meadow spittlebug, Omnivorous leaftier, Orange tortrix, Oriental fruit moth, Oystershell scale, Peach twig borer, Pear leaf blister mite, Pear psylla, Pear rust mite, Periodical cicada, Plum curculio, Prune leafhopper, Raspberry sawfly, Red-banded leafroller, Rose chafer, Rosy apple aphid, Saltmarsh caterpillar, San Jose		
	scale, Snowy tree cricket, Strawberry fruitworm, Strawberry weevil, Tarnished plant bug, Tentiform leafminers, Tussock moth, Variegated leafroller, Western grapeleaf skeletonizer, Western tussock moth, West Indian sugarcaneborer (adults), Western yellowstriped armyworm, White apple leafhopper, Wooly apple aphid, Yellow headed fireworm.		

ORNAMENTAL TREES, SHRUBS AND FLOWERS

This product may be used as a spot treatment for localized pest infestations on trees. Do not use on large trees or where pest populations are generally inaccessible to spray deposits. Direct spray toward the upper and lower leaf surfaces and small trunks, stems and twigs to the point of runoff where pests appear. Repeat applications as necessary up to a total of four times for trees and six times for ornamental, shrubs and flowers, but not more often than once every seven days.

ORNAMENTAL TREES, SHRUBS AND FLOWERS	PESTS
Ornamental Trees, Shrubs and Flowers	Ants (excluding harvester, carpenter, and pharoah ants), Apple aphid, Armyworm, Azalea leafminer, Bagworms, Birch leafminer, Blister beetle, Boxelder bug, Boxwood leafminer, Browntail moth, Cankerworms, Catalpa sphinx, Chiggers*, Cooley spruce gall, Elm leaf aphid, Elm spanworm, Eriophyid mites, European pine- shoot moth, Fall armyworm, Flea beetles, Fuller rose beetles, Gall midges, Gall wasps, Grasshoppers, Green striped- Mapleworm, Gypsy moth, Hackberry nipplegall makers, Holly budmoth, Holly leafminer, Jackpine budworm, Japanese beetle, Jeffrey pine needleminer, June beetles, Plant bugs, Lace bugs, Leafhoppers, Leafrollers, Locust borer, Maple leafcutter, Mealy bugs, Mimosa webworm, Nantucket pine tip moth, Oak leafminers, Oak leaf skeletonizer, Oakworm complex, Oleander caterpillar, Olive ash borer, Orange-striped oakworm, Orange tortrix, Periodical cicada, Pine Sawfly, Pine spittlebug, Pitch pine tip moth, Poinsettia horn worm, Psyllids, Puss caterpillar, Redhumped oakworm, Rose aphid, Rose chafer, Roseslug, Saddled prominent, Sawflies (exposed), Scale insects, Sowbugs, Spiny elm caterpillar, Springtails, Spruce budworm, Spruce needleminer, Subtropical pine tip moth, Tent caterpillar, Thorn bug, Thrips (exposed), Ticks*, Walnut caterpillar-Webworms, Western hemlock loopers, Western spruce budworm, Willow leaf beetles, Yellow poplar-weevil

^{*}Will kill pests that are present only at the time of application. Pest must be directly contacted for product to be effective.

Plant Response Precautions

Do not use on Boston Ivy, Virginia creeper and Maidenhair fern as injury will result. During early season it may also injure Virginia and Sand pines.

PESTS AROUND OUTDOOR RESIDENTIAL AREAS

This product may be used as a spot treatment around the outdoor perimeter of residential buildings. Thoroughly wet the outside perimeter of dwellings where pests tend to congregate. Use on lawns is prohibited. Will kill target pests that are present only at the time of application. Pest must be directly contacted for product to be effective.

NOTE: Staining may occur on certain surfaces such as stucco, brick, cinder block and wood. Therefore, applications of this product to surfaces where a noticeable residue or discoloration is objectionable should be avoided.

PESTS

Ants (including imported red fire ants), Bees, Brown dog ticks, Centipedes, Cockroaches, Crickets, Earwigs, Firebrats, Fleas, Millipedes, Silverfish, Spiders

[NOTICE: Research and testing have determined that the "Directions For Use" are appropriate for the proper use of this product under expected conditions. To the extent consistent with applicable law, the Buyer assumes responsibility for lack of performance or safety if not used according to the directions].

[Money Back Guarantee: If you are not satisfied with this product, we will gladly refund your original purchase price.]

Sevin is a registered trademark of Bayer

[Optional language (only if required by the State of CA):

WARNING: This product contains carbaryl, a chemical known to the State of California to cause [cancer] and [reproductive harm] in laboratory animals.]



Bayer Environmental Science

A Business Group of Bayer CropScience LP P.O. Box 12014, 2 T.W. Alexander Drive Research Triangle Park, North Carolina 27709 1-866-99BAYER (1-866-992-2937)

CP CARBARYL Insecticide Spray (MASTER) Approved 03/04/10, Notification 11/08/10