432-1213

19/31/3002

| 1 | 115 |
|------|-----|
| 2-28 | 95 |

| Please read instructions or | reversa before completing (| отт. | | Form Appro | ved. OMB No. | 2070-00 | 80. Approvel expires 2-28-9 |
|--|---|---|---|---|--|------------------------------------|---|
| \$EPA | United States Environmental Protection Agency Washington, DC 20460 | | | | Registra Amend Other | | OPP Identifier Number |
| | App | dication fo | or Pestici | de - Sectio | n I | | |
| 1. Company/Product Numb 432-1212 | or | | 2. EPA I John H | Product Manage lebert | r | 3. P | roposed Classification |
| 4. Company/Product (Name Sevin Grub Killer Grai | | | PM# 7 | _ | | | |
| Bayer Environmenta 95 Chestnut Ridge F Montvale, NJ 07645 | | | (b)(i), rr to: EPA R | | imilar or iden | tical in co | n FIFRA Section 3(c)(3) pmposition and labeling |
| | ······································ | S | ection - I | | | | |
| Explanation: Use addition Label Notification per PR Notification is consister labeling or the confidential sepa. I further understand the second second separation is separational separation. | ponse to Agency letter dated | or section I and ting Claim to the tice 98-10 and E duct. I understa | e label. EPA regulation and that it is a v terms of PR N | riolation of 18 U.3 lotice 98-10 and | dated lication. below. 46, and no oth S.C. Sec. 1001 | er changes to willfully | make any false statement to |
| | | S | ection - II | I | <u> </u> | . | |
| 1. Material This Product W | ill Be Packaged in: | | | · · · · · · · · · · · · · · · · · · · | | | |
| Child-Resistent Packaging Yes No * Certification must be submitted | Unit Packaging Yes No If "Yes" No | per if | Yes Yes No Yes" ckage wgt | No. per container | 2. Type of | Metal Plastic Glass Paper | Specify) |
| 3. Location of Net Contents Label 6. Manner in Which Label is | Container | Lithograph Paper glued Stenciled | ntainer | 5. | Location of Lab | pel Directio | ons |
| | <u></u> | | ction - IV | , _ | | | |
| 1. Contact Point (Complete | items directly below for ide | | | | ecessary to o | ocaes this | application ! |
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Name NORMA C. PANGILINAN Title Manager, Registrations Telephone No. (Include Area Manager) 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | | | | | e No. (Include Area Code) | |
| • | ements I have made on this f ny knowlingliy false or misle: | | | | | • | 6. Date Application Received (Stamped) |
| 2. Signature | | 3. Title | 3. Title | | | | |
| Harzil | ira | Mana | Manager, Registrations | | | | |
| 4. Typed Name | _ | 5. Date | _ | | | | |
| Norma C. Pangilinan, Ph.D. | | | December 6, 2005 | | | | |

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

White - EPA File Copy (original)

Yellow - Applicant Copy

NOTIFICATION

DEC 2 1 2005

SEVIN GRUB KILLER GRANULES (2% SEVIN)

For Use on Vegetables and Fruits
For Home Lawns and Ornamental/Recreational Turf
To Kill and Control Fire Ants, Ticks (including those that may vector Lyme Disease)

* Kills Ticks! Lyme Disease

Kills: Ants, Fleas, Ticks, Grubs....and many more.

- * Kills Japanese Beetles * Slow Release Granules * Bug Killer
- Kills Numerous [Over 30] Pests on Lawns and Ornamentals * Covers Up to _____Square Feet
 *Easy to Use Granules *Use on Lawns, Ornamentals and Around Homes *Special Value! 20%
 Bonus 12 lbs. for the price of 10 lbs.

ACTIVE INGREDIENT:

Keep Out of Reach of Children CAUTION

See back panel for additional precautionary statements

Net Weight: 5 pounds [Covers up to 2,146 sq. ft.] Net Weight: 10 pounds [Covers up to 4,292 sq. ft.] Net Weight: 12 pounds [Covers up to 5,150 sq. ft.] Net Weight: 20 pounds [Covers up to 8,584 sq. ft.]

EPA Reg. No.432-1212

EPA Est.___

Bayer Environmental Science, A Business Group of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

|Back Panel|

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed or inhaled. Avoid breathing avoid dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks and household or latex rubber gloves. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instruction for washable are provided, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product. Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking and using the toilet.
- Do not allow children or pets to contact treated area until granules have been absorbed into the soil.

| | FIRST AID |
|--|---|
| This product | is a N-methyl carbamate and is a cholinesterase inhibitor |
| If Swallowed | Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If Inhaled | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. |
| If in Eyes | Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. |
| If on Skin or Clothing | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. HOTLINE NUMBER |
| Have the product container or label You may also co | with you when calling a poison control center or doctor, or going for treatment. ntact 1-800-334-7577 for emergency treatment information. |
| | NOTE TO PHYSICIAN |
| | nate insecticide which is a cholinesterase inhibitor. Specific antidote consists of ves should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are not recommended. |

Environmental Hazards

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for commercial seed production, research purposes, sale or other commercial use such as sod.

Read Entire Label Before Using This Product
Strictly observe label directions and cautions. Do not apply this product in a way that will contact applicators or other persons, either directly or through drift. Unprotected persons must vacate the area being treated.

{For product packaged in containers of 5 pounds or less:}

Storage and Disposal

Storage: Store in original container in a cool, dry place away from children and animals. Avoid excess heat. Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

{For product packaged in containers greater than five pounds:}

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container in a cool, dry place away from children and animals. Avoid excess heat. In case of spill, confine spill and avoid contact; isolate and keep away from animals and unprotected persons. Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Container Disposal: Plastic: Triple rinse or equivalent. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Bags: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL DIRECTIONS FOR LAWNS, ORNAMENTAL AND RECREATIONAL TURF

Use when insects first appear and/or plant damage is imminent. For best results, mow lawn first and then apply material uniformly to dry turfgrass. Irrigate ¼" to wash insecticide into the grass. Do not allow use of treated areas during application. Keep children and pets off treated areas until grass is dry. Apply when air is calm to avoid drift and contact with skin and eyes. Repeat applications as necessary up to a total of 4 times per year but not more often than once every 7 days.

| Treatment | | Pounds of This Prod | uet |
|--|---|---------------------|---|
| Areas | Pests | Per 1000 Sq. Ft. | Comments |
| Lawns Ornamental And Recreational Turf Areas | Ants Lucerne Moth Armyworms Millipedes Centipedes Mosquitoes (Adults) Chiggers Scarab beetle adults: Chinch bugs Such as May beetle, Cutworms June beetle, Japane Earwigs beetle, Green June Essex skipper beetle European Sod Webworm Chafer (lawn moths) Fall army- Sowbugs Worm (lawn moths) Fiery Skipper Spittlebugs Grass- Springtails hoppers Ticks Green june Yellowstriped beetle larvae Leafhoppers | | For effective control, irrigate after application to remove chemical from granules. For heavy infestations of Chinch bugs, increase rate to 5lbs per 1000 sq. ft. and and treat the entire area. |



Black 9 For effective control of Billbag larvae turfgrass subsurface pests, apply European crane ataenius beetle when insects are fly larvae larvae actively feeding near Fleas the soil surface, White grubs: Hyperodes weevil usually during late Such as Japanese March through May, (bluegrass Beetle larvae, Chafer beetle Weevil larvae) or July through early September, or as re-Larvae. commended by local Phyllophaga spp. agricultural extension Larvae, Annual service agents. White grubs irrigate thoroughly after application to remove chemical from granules and move it down into the thatch.

Control of Ticks That May Vector Lyme Disease

For control of juvenile and adult Ixodes spp. ticks (Deer Ticks and Black Legged Tick) and ticks (Lone Start Tick) which may vector Lyme Disease: Apply granules at the rate of 7 to 9 pounds of product per 1000 sq. ft. with appropriate equipment to ensure uniform coverage. Target applications to control juvenile ticks (larvae and nymphs) in late spring or early summer (May/June). Control of adult ticks can be achieved with late summer and fall applications (August/September)> Make applications to lawns, shrub beds, ornamental plantings, wooded areas and around outside perimeters of home and other buildings. Do not make spot applications. Treat entire lawn and perimeter wooded areas and property boundaries where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove chemical from the granules. Ticks may be reintroduced from surrounding areas or host animals. Reapply as necessary to maintain adequate control, but not more often than once every 7 days.

Imported Fire Ants

For use as a mound treatment to control imported fire ants occurring in lawns, turf and premise areas, apply ½ cup of granules over and around each mound (Approximately 2'x 2'). Thoroughly irrigate mound immediately after treatment to remove to chemical from the granules and flush it down into the mound. A minimum of 1 gallon of water should be sprinkled over each mound, but do not use more than 2 gallons of water as flooding of the mound may result. Apply water gently to avoid disturbing ants and cause migration before chemical can be effective. For best results apply in cool weather (650F to 800F), early morning or late afternoon. Treat new mounds as they appear.

For Flowering Plants and Shrubs (Ornamental Plantings) in Landscape and Foundation Plantings.

| Pests | | Pounds of This Product Per 1000 Sq. Ft. | Specific Directions |
|---|---|--|--|
| Ants Armyworms Centipedes Crickets | Firebrats Fleas Millipedes Scorpions | 2.25 to 5 | Application to wet foliage periods of high humidity may cause injury to tender foliage. |
| Cutworms Earwigs | Silverfish Spiders Ticks | | Apply granules in a 6-foot band arounthe building, to shrub and flowerbeds, foundations, ornamental plantings and Lawn or soil areas immediately adjaces. To the building. For most effective Control, treated area should be lightly Watered after applications to remove The chemical from the granules. |
| | | | Repeat as needed for control up to 3 times per year but do not apply More often than once every 7 days. |
| | | | Do not use on plants above chest heigh |

Nuisance Pest Control Around the Outside of Homes, Porches, Patios and Buildings For control of Ants, Armyworms, Centipedes, Crickets, Cutworms, Earwigs, Firebrats, Fleas, Millipedes, Scorpions, Silverfish, Sowbugs, Spiders and Ticks: Apply granules uniformly in a 8 foot band around the building. Apply granules at the rate of 2 1/3 lbs. per 1000 sq. ft. to shrub and flower beds, foundations, ornamental plantings and lawn and soil areas immediately adjacent to the building. For most effective control, treated area should be lightly watered after application to remove the chemical from the granules. Repeat application as necessary to control new infestations.

Nuisance Pest Control in Vegetables and in Small Fruits and Berries (Not for Use in California)

Not to be applied to plants being grown for commercial purposes. For control of ants, centipedes, crickets, cutworms, earwigs, grasshoppers, millipedes, sowbugs (pillbugs), springtails and ticks. Water lightly after application to release the chemical from the granules. Applications may be made at any time before, during or after the growing season but it is recommended to begin applications when insect or damage is first observed and repeat as directed below. Retreatment intervals (days between treatments) are found listed in the table below.

Preharvest Use Limitation: Observe the listed preharvest intervals (number of days between the last application and harvest) for application during the growing season.

Do not use crops other than those listed in the following table.

| Crop | Maximum Single Application Rate (lbs./1000 sq. ft.) | Maximum Seasonal Application Rate (lbs./1000 sq. ft.) | Retreatment Interval (days) | PHI (days) | Comments |
|---|---|---|-----------------------------------|---------------|--|
| Asparagus (spears) | 2.25 | 6.88 | 3 | 1 | Total maximum |
| Asparagus (ferns*) | 2.25 | 4.6 | 7 | 1 | seasonal application rate is 23 lbs. per 1000 sq. ft. per crop. The total number of applications is 3 prior to harvest with 2 additional applications only to ferns. *after harvest of spears |
| Beans (fresh & dried): | 1.7 | 6.88 | 3 | 3(S) | S = succulents |
| [snap, navy & kidney], | | | | 21(D |) D = dried |
| Peas, (fresh & dried), | | | | 14(F) |) F = forage |
| Cowpeas, Lentils | | - | | 21(F | H = hay |
| Garden beets (roots), 2 Carrots, Horseradish, Radishes, Parsnips, | 2.25 | 6.88 | 7 | 7 | Total number of applications per crop can be 6 (some of |
| Rutabaga, Salsify, | | | | | which must be lower than the maximum single application |
| Turnips (roots) | | | | | rate) |
| Sugar beets | 1.7 | 3,45 | 14 | 28(R | • |
| Mustard greens | 2.25 | 6.88 | 7 | 14 | Total number of applications per crop can be 4 (some of which must be at lower than maximum single application rate) |

| Brussels sprouts, Broccoli, Cabbage Cauliflower Chinese Cabbage, Collards, Kale, Kohlrabi | 2.25 | 6.88 | 7 | 3(A) A = Brussels 14(B) sprouts, broccol cabbage, kohlra and cauliflower B = Chinese cabbage, kale an collards Total number of applications per crop can be 4 (some of which must be at a lower than maximum single |
|--|------|-------|----|---|
| Corn (field and pop) | 2.25 | 9.2 | 14 | application <u>rate)</u> 14(Fr) Fr=forage, silag 48(Fd) Fd=fodder, grai |
| Corn (sweet) | 2.25 | 18.35 | 3 | 2 E-ears 14(Fr) Fr=forage 48(Fd) Fd=fodder |
| Cucumber, melon Squash | 1.15 | 6.88 | 7 | 3 |
| Lettuce (head & Leaf) Endive, Celcery, Dandeliion, Parsley, Spinach, Swiss chard, Hanover salad, Rhubarb, Garden Beets (top), Turnips (tops) | 2.25 | 6.88 | 7 | Total number of applications per crop can be 5 (some of which must be at lower than maximum single applicatio rate) |

| Crop | Maximum Single Application Rate (lbs./1000 sq. ft.) | Maximum "Seasonal Application Rate (lbs./1000 sq. ft.) | Retreatment Interval (days) | PHI (days) | Comments |
|-----------------------------|--|--|-----------------------------------|--------------------|---|
| Potatoes | 2.25 | 6.88 | 7 | 7 | Total number of applications per crop can be 6 (some of which must be at lower than maximum single application rate) |
| Grain sorghum (not sweet) | 2.25 | 6.88 | 7 | 14 (Fr) 21 (Fd) | Fr = forage, grazing Fd = forage, grain Total number of applications per crop can be 4 (some of which must be at lower than single application rate). |
| Strawberry | 2.25 | 11.5 | 7 | 7 | |
| Tomato, Pepper, Eggplant | 2.25 | 9.2 | 7 | 3 | Total number of application per crop can be 7 (some of which must be at lower than maximum single application rate) |
| Wheat | 1.7 | 3.45 | 14 | 21(G) 7 (F) | G=grain or straw F=forage |

Do not plant rotational crops not listed on this or other carbaryl labels in carbaryl-treated soil.

Sevin Grub Killer granules are applied using a commercial spreader. Spreaders and settings are found in the table below. Do not repeat applications more than once a week.

| | | | Spreader S | Settings |
|-----------------------------------|-------------|---------------------|------------|----------------------|
| | | Pounds/1000 sq. Ft. | | Pounds / 500 Sq. Ft. |
| Spreader | 2/ 1/3 lbs. | 4 2/3 lbs. | 3.5 lbs. | 4.5 lbs. |
| Scots Even Green 18" | 15 | 16* | 15 | 16 |
| Scotts Speedy Broadca | st 14 | 15* | 14 | 15 |
| Red Devil Broadcast | 12 | 13* | 12 | 13 |
| Red Devil Drop 20" | 12 | 13* | 12 | 13 |
| Republic EZ Drop 18" | 8 | 10 | 8 | 10 |
| Republic EZ Broadcas | t 9 | 13* | 9 | 13 |
| Republic EZ Tractor Broadcast | 9 | 13* | 9 | 13 |
| Other Spreaders As Appropriate | TBD | TBD | TBD | TBD |


Alternative Spreader Settings for Untreated Peanut Hulls:

| Spreader | Type | Rate (lbs/1000 ft2) | Setting |
|------------------------|----------|---------------------|---------|
| Scotts Precision Green | Drop | 2.25 | 8 |
| P-4 | | 4.6 | 11 |
| | | 6.8 | . 14 |
| | | 9.2 | 16.5 |
| | | 11.5 | NA |
| | | 18.35 | NA |
| Cyclone Rotary | Rotary | 2.25 | 5 |
| | | 4.6 | 6 |
| | | 6.8 | 8 |
| | | 9.2 | · NA |
| | | 11.5 | NA |
| | <u> </u> | <u> 18.35</u> | NA |
| Earthway Rotary | Rotary | 2.25 | 14 |
| • | | 4.6 | 18 |
| | | 6.8 | 27 |
| | | 9.2 | : NA |
| | İ | 11.5 | · NA |
| | | 18.35 | NA |
| Earthways Ev-N-Spred | Rotary | 2.25 | 14 |
| • | | 4.6 | 18 |
| | | 6.8 | 27 |

| | | 9.2 | NA |
|--------------------------|--------|-------|-----------------|
| | | 11.5 | NA |
| | | 18.35 | NA |
| Republic EZ | Drop | 2.25 | 9 |
| • | • | 4.6 | 12 |
| | | 6.8 | 13 |
| | | 9.2 | 13 |
| | | 11.5 | 15 |
| | | 18.35 | NA |
| Scotts Speedygreen | Rotary | 2.25 | 8 |
| 3000 | • | 4.6 | NA |
| | • | 6.8 | NA |
| | 1 | 9.2 | NA |
| | I | 11.5 | NA |
| | | 18.35 | • NA |
| Craftsman Rotary (0195) | Rotary | 2.25 | 6 |
| | 1 | 4.6 | 10 |
| | | 6.8 | 15 |
| | | 9.2 | [†] NA |
| | | 11.5 | NA NA |
| | | 18.35 | NA |
| Precision EZ Spread | Rotary | 2.25 | 7 |
| SB550 | 1 | 4.6 | 9 |
| | | 6.8 | NA |
| | | 9.2 | NA |
| | | 11.5 | NA |
| · | + | 18.35 | NA |
| Pursell's Sta-Green | Rotary | 2.25 | 8 |
| SB4500 | | 4.6 | 9 |
| | | 6.8 | NA |
| | | 9.2 | NA |
| | | 11.5 | NA |
| | + | 18.35 | NA |
| Agri-Fab 85 lb. Spreader | Rotary | 2.25 | 6 |
| Model #4503123062 | | 4.6 | 7 |
| | | 6.8 | 8 |
| | | 9.2 | 10 |
| | | 11.5 | NA |
| | | 18.35 | NA. |

NA: Not applicable, no setting for this rate.

Example:



Buyer assumes all risks of use, storage or handling of this product not in strict accordance with this label.

If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call: (800)-420-9347

If you have questions relating to this product use or application, please call: (800) 969-7200 If you have a product spill or transportation emergency call: (800) 424-9300 For Material Safety Data Sheets please go to www.gardentech.com, or call (800) 969-7200.

Ak.111504

Bayer Environmental Science



VIA COURIER

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
1801 Bell Street
Arlington, VA 22202

NOTIFICATION

DEC 2 1 2005

Attn:

Ms. Sherada Hobgood

Subject:

Sevin Grub Killer Granules (2% Sevin) (EPA Reg. No. 432-1212)

-Label Notification per PR Notice 98-10:

Addition of Marketing Claim

Dear Ms. Hobgood:

Pursuant to PR Notice 98-10, Bayer Environmental Science wishes to notify the Agency of a minor labeling change for Sevin Grub Killer Granules (2% Sevin) EPA Reg No. 432-1212. The label change included addition of the marketing claims:

- "Kills Ticks! Lyme Disease"
- "Kills: Ants, Fleas, Ticks, Grubs.... and many more.".

In support of this notification, the following documents are being submitted:

- 1. Completed Application for Registration (EPA Form 8570-1).
- 2. Three copies of the revised product label plus one copy highlighting changes,

This notification is consistent with the provisions of PR Notice 98-10 and EPA Regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec.1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

December 6, 2005

Bayer Environmental Science 2 T.W. Alexander Drive Research Triangle Park NC 27709

Tel. No. 919-549-2000 Fax. No. 919-549-3937 Page 2 of 2

Please do not hesitate to call (919-549-2428) or e-mail me (norma.pangilinan@bayercropscience.com), if you have any questions.

Thank you very much for your assistance in this mater.

Sincerely,

BAYER ENVIRONMENTAL SCIENCE A Business Group of Bayer CropScience

NORMA C. PANGILINAN, Ph.D,

Manager, Registrations

cc: Reg. Book w/ attachments
K. Shearer w/o attachments