
		2	,		۱/s
		3.14.200			
lease read instructions on reverse bet	ore completing form. United States				0060, Approval expires 05-31-9 OPP Identifier Numb
			Registr		
	vironmental Protection	•••	Amend	ment	288964
	Washington, DC 2046	.0	Other:		
	Applicatio		ide - Sectior	1	
1. Company/Product Number 432-1259		2. EPA Produ J. Tavano	ct Manager	- un	3. Proposed Classification
4. Company/Product (Name) Maxforce RBF5		PM# 10			None 🗌 Restric
5. Name and Address of Applicant (Bayer Environmental Scien of Bayer CropScience LP 95 Chestnut Ridge Road Montvale, N.J. 07645 <u>PLEASE SEND ALL COR</u> <u>"CONTACT POINT" I</u>	ce, A business Group <u>RESPONDENCE TO</u> LISTED BELOW	(b)(I), my pr to: EPA Reg. N	oduct is simila		
Check if this is a new a	ddress				
		Section -			
Amendment – Explain below.					Ageneration
Resubmission in response to	Agency letter dated		"Me Too" Applica		MAR 1 4 2003
Notification - Explain below. Explanation: Use additionation	1 (-) :6 (Other - Explain		14 2003
	<u></u>	Section - I			····
		Section -	· · · · · · · · · · · · · · · · · · ·		
 Material This Product Will Be Pro	Unit Packaging	w	ater Soluble Pack	aging	2. Type of Container
Yes*	Yes		Yes		Metal
No No	<u>No</u>		No		Plastic
		j ···	81/		
*Certification must			"Yes" ackage wol	No. per	Glass
hocubaittod		_	ackage wgt.	No. per container	Glass Paper
be submitted	ation 4 Size(s) B(container	Glass Paper Other (Specifiy)
3. Location of Net Contents Inform	lation 4. Size(s) Re	etail Container		container 5. Location of	Glass Paper
3. Location of Net Contents Inform				5. Location of	Glass Paper Other (Specifiy)
3. Location of Net Contents Inform	ntainer	etail Container		5. Location of	Glass Glass Glass Cher (Specifiy) Label Directions
3. Location of Net Contents Inform	ntainer	etail Container ph lued d	ackage wgt.	5. Location of	Glass Glass Glass Cher (Specifiy) Label Directions
3. Location of Net Contents Inform Label Co 6. Manner in Which Label is Affixe	ntainer d to Product Dithogra Paper g Stencile	etail Container ph lued d Section - I	ackage wgt.	Container 5. Location of On Label	Glass
 3. Location of Net Contents Inform Label Co 6. Manner in Which Label is Affixe 1. Contact Point (Complete items of the second second	ntainer d to Product Lithogra Paper g Stencile directly below for identification	etail Container ph lued d Section - I	ackage wgt.	container 5. Location of On Label On labelin On labelin	Glass
3. Location of Net Contents Inform Label Co 6. Manner in Which Label is Affixe	ntainer d to Product Dithogra Paper g Stencile	etail Container ph lued d Section - I of individual to be	ackage wgt.	container 5. Location of On Labelin On labelin essary, to process	Glass
3. Location of Net Contents Inform Label Co 6. Manner in Which Label is Affixe 1. Contact Point (Complete items of Name	Intainer	etail Container ph lued d Section - I of individual to be Man chments thereto a lay be punishable	Other	container 5. Location of On Label On labelin assary, to process	Glass
3. Location of Net Contents Inform Label Co 6. Manner in Which Label is Affixe 1. Contact Point (Complete items of Name Adrian Krygsman I certify that the statements I have acknowledge that any knowingly fa	Intainer	etail Container ph lued d Section - I of individual to be Man chments thereto a lay be punishable . Title	Other	container 5. Location of On Label On labelin assary, to process	Glass

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete

.....


i

.



ł

2/8





RCACH KILLER BAIT GEL (3-Pack)

FOR USE IN COMMERCIAL . INDUSTRIAL AND RESIDENTIAL AREAS

ACTIVE INGREDIENT:

Fipronil*: [5-Amino-1-(2,6-dichloro-4-(trifluoromethyl)phenyl)-4-(1,R,S)-(trifluoromethyl) sulfinyl)-1H-pyrazole-3-carbonitrile]0.01%

*CAS NO. 120068-37-3

Contains: 3 plastic dispensers

KEEP OUT OF REACH OF CHILDREN **CAUTION:** SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Product may stain textiles and clothing

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep gel away from foods and food contact surfaces. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS: Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area out of reach of children

DISPOSAL: Wrap used pre-filled plastic reservoirs in several layers of newspaper and discard in trash.

If you have questions regarding the use of this product, please call 1-800-331-2867.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

MAXFORCE® FC Professional Insect Control® Roach Killer Bait Gel may be used as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structures) control of cockroaches in residential areas and the non-food/non-feed areas of institutional warehousing and commercial establishments, including warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, schools, laboratories, computer facilities, sewers, "aircraft," buses, boats/ships, trains, pet shops and zoos. Non-food/non-feed areas are areas such as garbage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

GENERAL DIRECTIONS

For roach gel: Application rate is dependent on level of infestation and species to be controlled. Bait can be applied as spots or as a bead in cracks and crevices. The lower application rate should be used for low to moderate infestations or re-treatments, and the higher rate for more severe infestations or where the population occurs in inaccessible areas, such as in walls. Gel placements should be at or near harborages or aggregation areas, such as corners, areas of movement or cracks and crevices. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control.

Do not treat food preparation surfaces. Do not apply bait get to areas which have been recently sprayed with insecticide, and do not spray insecticide over gel as it may cause the bail to become repellent.

For outdoor use (adjacent to home -or- structure): Place bait in areas adjacent to structures where roaches may nest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints. Apply bait in protected areas whenever possible. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness.

Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushing.

Retillable stations: This product may be used in the Maxforce Retillable Bail Station or B&G Perimeter Patrol Station. Place 0.5 to 5 grams of product per station. Do not exceed 5 grams per station (or up to 1 tsp). Refillable stations should be placed every 20 to 30 teet around the perimeter of a structure. Use approximately 2 stations per side of a typical single family home For best results, place stations near: air conditioning units, water faucets, foundation vents and weep holes,

SER EN LORDER MESSERT

decks, utility entryways (e.g. electrical, cable TV) and wherever target pests are frequently found. Regularly inspect all stations and replace bait as needed for continual control.

Position Refillable Bait Stations adjacent to structure in areas where they will not be damaged by lawn equipment such as mowers and trimmers. When placing the station in turf remove sod layer and place

station base plate against the soil. If the station is not equipped with a sticker to identify the contents, apply a sticker to the outside or inside of the station.

Applications in Food/Feed Handling Establishments: Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only. If gel contacts an exposed surface where food is

MARKAMI LOGINISADOLI INC ALTA HENRA DIMINE REFESTIVE RES in Print Contraction La Salar Of Street En Victor Contraction 2...... 2.014.8 SIVE DE LIS REALT? REESTRETENS FOR DENNA DE REDAK BAREEDERE PER 21,560 Abort Br STREET, MANUEL 121-113 F 114 MILE #GOINS SEE A GARGERAN MENNY PROVIDE GELENNE A BAR SEE STREE LA. HELMMENDER XVI SCHENNES

KERDEA DI GURK 15 が住ったの

handled, remove gel and wash exposed surface with an effective cleaning compound followed by a potable water rinse prior to use. Apply with syringe applicator directly into cracks and crevices. Place applicator tip 1/2 inch into cracks, crevices, holes and other small openings where roaches may be a problem. Apply small amounts of gel (up to 1/2 gram) per spot. Do not apply bait to areas where food/leed, utensils or processing surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation and prepared food holding surfaces as balt may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 130°F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, fume hoods, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a tood/feed area when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves and other equipment where cockroaches hide. During follow-up visits inspect bait placements and reapply when necessary. Application of this product in food/teed areas of food/teed handling establishments other than as a crack and crevice treatment is not permitted.

Contains: 3 plastic dispensers NET WT 2.1 OZ (60g) each TOTAL NET WT 6.3 OZ (180q)

© Baver AG 2002 Bayer Environmental Science MAXFORCE

A Business Group of Bayer CropScience LP 95 Chestnut Ridge Road, Montvale, NJ 07645

EPA Reg. No. 432-1259 EPA Est. No. 5813-MS-01 Maxforce is a registered trademark of Bayer AG

12/02



ROACH KILLER BAIT GEL (Reservoir)

FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

ACTIVE INGREDIENT:

*CAS NO. 120068-37-3

Contains: 4 plastic reservoirs

KEEP OUT OF REACH OF CHILDREN

CAUTION: SEE BELOW FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Product may stain textiles and clothing.

PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep gel away from foods and food contact surfaces. Harmful it absorbed through skin Causes moderate eye irritation. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

PHYSICAL AND CHEMICAL HAZARDS: Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area out of reach of children.

DISPOSAL: Wrap used pre-filled plastic reservoirs in several layers of newspaper and discard in trash.

If you have questions regarding the use of this product, please call 1-800-331-2867.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

MAXFORCE® FC Professional Insect Control® Roach Killer Bait Gel may be used as a spot or crack and crevice treatment for the indoor and outdoor (adjacent to homes & structires)control of cockroaches in residential areas and the non-food/non-feed areas of institutional, warehousing and commercial establishments, including warehouses, restaurants, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, schools, laboratories, computer facilities, sewers, "aircraft," buses, boats/ships, trains, pet shops and zoos. Non-food/non-feed areas are areas such as garbage rooms, lavatories, entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or botting). This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

GENERAL DIRECTIONS:

For roach get: Application rate is dependent on level of infestation and species to be controlled. Bait can be applied as spots or as a bead in cracks and crevices. The lower application rate should be used for low to moderate infestations or re-treatments, and the higher rate for more severe infestations or where the population occurs in inaccessible areas, such as in walls. Gel placements should be at or near harborages or aggregation areas, such as corners, areas of movement or cracks and crevices. Numerous smaller placements will provide faster control than fewer larger spots, especially for German cockroach control. Do not treat tood preparation surfaces. Do not apply bait gel to areas which have been recently sprayed with insecticide, and do not spray insecticide over gel as it may cause the bait to become repellent.

For outdoor use (adjacent to home -or- structure): Place bait in areas adjacent to structures where roaches may nest or breed. Apply bait to potential points of insect entry, such as eaves, sills, and expansion joints. Apply bait in protected areas whenever possible. Environmental stresses such as direct sunlight and water will reduce the residual effectiveness.

Priority treatment areas may be identified by visual inspection of such areas listed above or aided by trapping techniques or spot flushing.

Refiltable stations: This product may be used in the Maxforce Refiltable Bait Station or B&G Perimeter Patrol Station. Place 0.5 to 5 grams of product per station. Do not exceed 5 grams per station (or up to 1 tsp). Refiltable stations should be placed every 20 to 30 feet around the perimeter of a structure. Use approximately 2 stations per side of a typical single family home. For best results, place stations near: air conditioning units, water faucets, foundation vents and weep holes, decks, utility entryways (e.g. electrical, cable TV) and wherever target pests are frequently found. Regularly inspect all stations and replace bait as needed for continual control. Position Refliable Bait Stations adjacent to structure in areas where they will not be damaged by lawn

equipment such as mowers and trimmers. When placing the station in turf remove sod layer and place station base plate against the soil.

If the station is not equipped with a sticker to identify the contents, apply a sticker to the outside or inside of the station Applications in Food/Feed Handling Establishments: Application within food/feed areas of lood/leed handling establishments is limited to crack and crevice treatment only. If gel contacts an exposed surface where food is handled, remove get and wash exposed surface with an effective cleaning compound followed by a potable water rinse prior to use. Apply with Maxforce Bait Injector directly into cracks and crevices. Place applicator tip 1/2 inch into cracks, crevices, holes and other small openings where roaches may be a problem. Apply small amounts of gel (up to 1/2 oram) per spot. Do not apply bait to areas where food/feed, utensils or processing surfaces may

RECIES/LEVEL OF Nestation	*PPLIX*TONR*TE (PERSO.YD.)
IODERATE IFESTATIONS DREETRETSFOR SEMANOR SROWNERNOED ROMINES:	1-25P0 TS (DINE S2ED) OR 142": 5" BEAD
EVERE ORHEAVY IFESTATIOIIS FOR Servan or Brown Banded Roaches:	24 SPOTS (OU WATTER STZD.) OR 57 67 BESD
. ANGE ROX CHES SUCH AS KNERICKN, SNOKEY: BROWN DROKENTRI:	redoninipaded) spot Szz ¹ a(+) _s ² g

become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/leed preparation and prepared food holding surfaces as bait may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 130°F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, tume hoods, heat lamps, coffee urns, steam tables, toasters, tryers, dishwashers and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation. Apply bain in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves and other equipment where cockroaches hide. During follow-up visits inspect bait placements and reapply when necessary. Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted.

Break-Ilp Reservoir: Prior to use, break the snip-tip closed end of the gel reservoir at the scored line. After use, the reservoir may be resealed with the enclosed black outlet cap. The black outlet cap is reusable.

Diractions for changing reservoirs: With the arrow on the drive rod pointing up, pull the drive rod out as far as it will go. Remove the empty reservoir, wrap, and place in trash can. Reservoirs that still contain gel can be sealed with the end cap and outlet cap for future use. To attach a new reservoir follow the above directions.

Contains: 4 plastic reservoirs NET WT 1.05 OZ (30g) each, TOTAL NET WT 4.2 OZ (120g)

© Bayer AG 2002 Bayer Environmental Science A Business Group of Bayer CropScience LP 95 Chestnut Ridge Road, Montvale, NJ 07645 12/02 EPA Reg. No. 432-1259 EPA Est. No. 5813-MS-01





: -

.

5

4/8

SUPPLEMENTAL LABELING MAXFORCE® FC PROFESSIONAL INSECT CONTROL ROACH KILLER BAIT GEL

EPA Reg. No. 432-1259

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Refillable Stations:

1

this product may be used in the Maxforce Refillable Bait Station or B&G Perimeter Patrol Station. Do not exceed 1 tsp. per refillable bait station. Refillable stations should be placed every 20to 30 feet around the perimeter of a structure, use approximately 2 stations per side of a typical single family home.

For best results, place stations near; air conditioning units, water faucets, foundation vents and weep holes, decks, utility entryways (e.g. electrical cable TV) and wherever target pest are frequently found.


position Refillable Bait Stations (adjacent to home or structure) in areas where they will not be damaged by lawn equipment such as mowers and trimmers. When placing the station in turf, remove sod layer and place station base plate against the soil.

If the station is not equipped with a sticker to identify the contents, apply a sticker to the outside or inside of the station. Regularly insect all stations and replace bait as needed for continual control.

THIS LABEL MUST BE IN THE POSSESSION OF THE USER. REFER TO THE MAIN LABEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

Bayer Environmental Science A Business Group of Bayer CropScience LP 95 Chestnut Ridge Road Montvale, NJ 07645

14-1-supp



ID's 9¹³/16 x 9¹¹/16 x 8⁹/16 CDO #115255