



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

April 5, 2016

Kristian Paul
Regulatory Affairs Manager
Bayer Environmental Science; a division of
Bayer CropScience
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Subject: Label Amendment – Additional marketing claims; alternate brand name Savate
Product Name: BSN 2060 480SC Greenhouse and Nursery Ornamental
Insecticide/Miticide
EPA Registration Number: 432-1280
Application Date: 1/21/2016
Decision Number: 513412

Dear Mr. Paul:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

Alternate brand Savate is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 432-1280
Decision No. 513412

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Rita Kumar by phone at (703) 308-8291, or via email at kumar.rita@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to be 'MS' with a long horizontal stroke extending to the right.


Mark Suarez
Product Manager 07
Invertebrate & Vertebrate Branch 3
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

BSN 2060 480SC

Greenhouse and Nursery Ornamental Insecticide/Miticide

ABN: Savate



ACTIVE INGREDIENT:
 Spiromesifen: 2-oxo-3-(2,4,6-trimethylphenyl)-1-oxaspiro[4.4]non-3-en-4-yl
 3, 3-dimethylbutanoate45.2%

OTHER INGREDIENTS:54.8%

TOTAL:100.0%

BSN 2060 480SC contains 4 pounds of spiromesifen per US gallon (480 grams ai per liter).

EPA Reg. No. 432-1280 **EPA Est. No. 432-TX-01**

STOP - Read the label before use
Keep out of reach of children

CAUTION

See [Back][Side] Panel for First Aid Instructions and [Leaflet][Booklet] for Complete Precautionary Statements and Directions for Use. (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-334-9745

FIRST AID	
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration. • Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Note To Physician: No specific antidote is known. Treat symptomatically. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, loaders and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves (such as natural rubber, EPA selection category A)
- Shoes plus socks

Applicators must wear:

- Long-sleeved shirt and long pants

- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove Personal Protective Equipment immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Avoid contamination of surface water through runoff or spray drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater.

SURFACE WATER ADVISORY

This product may contaminate water through runoff or through drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read entire label and Conditions of Sale before using this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (**REI**) of **12 hours** following application. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are:

- Coveralls worn over short sleeved shirt and short pants
- Chemical resistant gloves (such as natural rubber, EPA selection category A)
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Offer for recycling, if available or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. If burned, stay out of smoke.

GENERAL INFORMATION

BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide is a Flowable (Suspension Concentrate) formulation that is intended for control of mites and whiteflies on ornamental plants, flowers and foliage plants in the greenhouse and field and container nurseries. The active ingredient contained in BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide is active on all mite development stages. However, mite juvenile stages are often more susceptible than adults. Application should be timed to coincide with early threshold levels in a developing mite population.

BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide is also highly effective against whitefly nymphs, plus it has a significant effect on the otherwise difficult to control pupal stage. For best results against whitefly, applications should target these life stages.

BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide can be applied by ground equipment. Thorough coverage of all plant parts is required for optimum performance. Performance evaluations should be made 4 – 10 days after application.

Apply specific dosage of BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide as needed for control. For best results good coverage of the upper and lower leaf surfaces is recommended. An adjuvant may be used to improve coverage on hard to wet foliage. For best results the treatment should be made when whitefly or mite population begins to build and before a damaging population becomes established. Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply with ground application equipment in adequate water for uniform coverage.

PRODUCT FEATURES

- Controls all stages of whitefly (nymphs, pupae, and adults).
- Provides knockdown and residual control of all developmental stages of mites

PESTS CONTROLLED	USE RATE
Spider mites (including twospotted spider mite, southern red mite, lewis mite, tumid mite, maple spider mite, spruce spider mite, honeylocust spider mite, euonymus mite, boxwood spider mite) Tarsonemid mites (including broad mite, cyclamen mite) Tenuipalpid mites (including false spider mite) Rust and blister mites (family <i>Eriophyidae</i>)	1 to 4 fl oz (30 – 120 mL)/100 gallons of spray solution or 1.5 – 6 mL/5 gallons of spray solution
Whiteflies (Including greenhouse, silverleaf and sweetpotato whitefly)	2 to 4 fl oz (60-120 mL)/100 gallons of spray solution or 3-6 mL/5 gallons of spray solution

<p>WHERE TO APPLY</p>	<ul style="list-style-type: none"> • To shrubs, trees (including non-bearing fruit and nut trees), flowers and foliage plants in greenhouses, shade houses and plants grown in field and container nurseries.
<p>HOW TO APPLY</p>	<ul style="list-style-type: none"> • Apply specific dosage of BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide as needed for control. For best results, apply to achieve uniform coverage of the upper and lower leaf surfaces but avoid excess application that results in runoff from plant foliage. • An adjuvant may be used to improve coverage on hard to hard-to-wet foliage. For best results the treatment should be made when the whitefly or mite population begins to build and before a damaging population becomes established. • BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide is very effective against the egg and nymphal stages of whiteflies and mites. Control should be directed at these stages. • Apply as a full coverage spray to the foliage using a minimum volume of 1 - 2 qts of final solution per 100 sq ft (or 100 to 200 gal per acre). Actual spray volume will vary depending on the size of plants. • For maximum efficacy against adult whiteflies, use the 4 fl oz rate of BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide. For heavy infestations, two applications may be necessary to control adult whiteflies. • Rate range is provided and is generally dependent on size of the plant and density of the foliage. Apply when pests first appear and prior to leaf damage or discoloration. Apply with ground application equipment in adequate water for uniform coverage. • Do not apply via chemigation. <p>Do not apply BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide more than three times per season.</p>

<p>RESISTANCE MANAGEMENT</p>	<ul style="list-style-type: none"> • BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide contains an active ingredient with a novel mode of action. Studies to determine cross-resistance with BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide linked to other commercial insecticide/miticides have demonstrated no cross-resistance. However, repeated use of any plant protection product may increase the development of resistant strains of pests, including insects and mites. • Using BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide in successive miticide applications during the same cropping cycle is not recommended. Use BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide as a part of a sound resistance management program that includes rotation with other treatments with different modes of action.
<p>PLANT TOLERANCE</p>	<ul style="list-style-type: none"> • BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide has been evaluated for phytotoxicity on a wide range of ornamental plants. However, due to the large number of species and varieties of ornamental plants, it is impossible to test every one for tolerance to BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide. The professional user should determine if BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide could be used safely prior to commercial use. In a small area, test the recommended rates on a small number of plants for phytotoxicity prior to widespread use. Before using BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide in tank mixture with other products and adjuvants, test the mixture on a small number of plants for phytotoxicity prior to widespread use. • Do not use BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide on geraniums (<i>Pelargonium</i> sp.), <i>Peperomia</i>, <i>Dracaena</i>, and 'Classy', 'Attache', or 'Vogue' varieties of rose. • Insufficient information on tolerance is available for the following ornamentals: lily-of-the-Incas (<i>Alstromoeria</i>), New Guinea impatiens, <i>Bacopa</i>, ferns, phlox, English ivy (<i>Hedera helix</i>), cyclamen, orchids, hoyo, <i>Agyranthemum</i>, hydrangea, schefflera, <i>Matthiola</i> (stock), Mexican heather, <i>Lobelia</i>, fuschia, <i>Cordylina</i>, croton, neanthebella palm, and primula. It is recommended that BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide not be used on these plants. • Do not apply more than 2 fl oz per 100 gal per application or 4 fl oz per cropping cycle to chrysanthemum, Shasta daisy, snapdragon, impatiens, verbena, lantana, gerbera daisy, or marigolds.
<p>SPRAY DRIFT</p>	<ul style="list-style-type: none"> • Avoid drift or runoff into lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds. Do not apply when winds are greater than 15 mph and avoid gusty conditions.
<p>IRRIGATION TIMING</p>	<ul style="list-style-type: none"> • If irrigation is used, conduct irrigation efficiently to prevent excessive loss of irrigation water through runoff. Time applications to allow sprays to dry prior to rain or sprinkler irrigation. Allow at least 24 hours between applications of product and any irrigation that results in surface water runoff into lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries, or commercial fish farm ponds.
<p>ENDANGERED SPECIES</p>	<ul style="list-style-type: none"> • The use of any pesticide in a manner that may kill or otherwise harm endangered species or adversely modify their habitat is a violation of Federal law.
<p>RESTRICTIONS</p>	<ul style="list-style-type: none"> • Do not use through irrigation systems. • Do not use on vegetable gardens. • Do not use on plants intended for use as feed or forage. • Do not use on bearing fruit or nut trees.
<p>PACKAGE HANDLING</p>	<ul style="list-style-type: none"> • BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide is packaged in polyethylene containers. Do not allow product or containers to freeze.

COMPATIBILITY / MIXING / ORDER-OF-MIXING

- BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide is physically and biologically compatible with many registered pesticides and fertilizers or micronutrients. When considering mixing BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide with other pesticides, or other additives, first contact your supplier for advice. For further information, contact your local Bayer representative. If your supplier and Bayer representative have no experience with the combination you are considering, you should conduct a test to determine physical compatibility. To determine physical compatibility, add the recommended proportions of each chemical with the same proportion of water as will be present in the chemical supply tank, into a suitable container, mix thoroughly and allow to stand for five minutes. If the combination remains mixed, or can be readily re-mixed, the mixture is considered physically compatible.
- BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide may be used with other recommended pesticides, fertilizers and micronutrients. The proper mixing procedure for BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide alone or in tank mix combinations with other pesticides is:
 - 1) Fill the spray tank 1/4 to 1/3 full with clean water;
 - 2) While recirculating and with the agitator running, add any products in PVA bags (**See Note**). Allow time for thorough mixing;
 - 3) Continue to fill spray tank with water until 1/2 full;
 - 4) Add any other wettable powder (WP) or wettable granules (WG) products;
 - 5) Add the required amount of BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide, and any other "flowable" (FL or SC) type products;
 - 6) Allow enough time for thorough mixing of each product added to tank;
 - 7) If applicable, add any remaining tank mix components: emulsifiable concentrates (EC), fertilizers and micronutrients.
 - 8) Fill spray tank to desired level and maintain constant agitation to ensure uniformity of spray mixture.
- Mix pesticides in areas not prone to runoff such as concrete mixing/loading pads, disked soil in flat terrain or graveled mix pads, or use a suitable method to contain spills and/or rinsate. Properly empty and triple-rinse pesticide containers at time of use.

For Use in California:

Allow growth of a vegetative filter strip within 25-ft (on which this product should not be applied) along lakes, reservoirs, rivers, permanent streams, marshes, potholes, vernal pools, natural ponds, estuaries or commercial fish farm ponds. This requirement does not apply to retention or holding ponds used to collect and recycle water for irrigation.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Bayer CropScience LP's control it is impossible for Bayer CropScience LP to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer CropScience LP disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 8 oz, 1 Gal

PRODUCED FOR



Bayer Environmental Science

A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

BSN 2060 480SC Greenhouse and Nursery Insecticide/Miticide (PENDING) 01/06/2016, 01/21/2016, 02/26/2016

[Marketing Claims]

- For Greenhouse and Nursery Use
- Miticide/Insecticide