



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

January 20, 2015

Norma C. Pangilinan, Ph.D.
Registration Manager
Bayer Environmental Science
Bayer CropScience
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Subject: Label Amendment – Add pest image, mode of action box, rate range for surface feeding insects, stewardship language under Environmental Hazards and Directions for Use, and clarify restriction on research use
Product Name: Dylox[®] 6.2 Granular Insecticide
EPA Registration Number: 432-1308
Application Date: October 21, 2014
Decision Number: 497683

Dear Dr. Pangilinan:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Page 2 of 2
EPA Reg. No. 432-1308
Decision No. 497683

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Eric Bohnenblust by phone at (703) 347-0426, or via email at Bohnenblust.eric@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Gebken", written in a cursive style.

Richard Gebken
Product Manager 10
Invertebrate & Vertebrate Branch 2
Office of Pesticide Programs

Enclosure

ACCEPTED

01/20/2015

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under
EPA Reg. No. 432-1308

GROUP

1B

INSECTICIDE

Dylox[®] 6.2 Granular Insecticide

For Control of White Grubs, Mole Crickets, Sod Webworms, Cutworms, and Other listed Pests of Turfgrass
Controls damaging insects below the soil surface

Provides control of Japanese beetle grubs

[Coleoptera: Scarabaeidae Image]

Editorial Note – [Bracketed text] is optional language

ACTIVE INGREDIENT:

Trichlorfon, Dimethyl (2,2,2-trichloro-1-hydroxy-ethyl)phosphonate..... 6.2%

OTHER INGREDIENTS:..... 93.8%

TOTAL:..... 100.0%

STORE IN A DRY PLACE

Treats 10,000 sq ft for White Grubs and Mole Crickets

Treats 15,000 sq ft for Sod Webworms

EPA Reg. No. 432-1308

EPA Est. No. ____

**STOP - READ THE LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN.**

CAUTION

PARA EL USUARIO: Si usted no lee o entiende inglés, no use este producto hasta que le hayan explicado completamente las instrucciones que figuran en la etiqueta.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

FIRST AID

Contains an organophosphate that inhibits cholinesterase.

If swallowed:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything to an unconscious person.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577.

Note To Physician: Prolonged exposure will cause cholinesterase depression. Atropine Sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Causes moderate eye irritation. Do not take internally. Do not breathe dust. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some people.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, Loaders, Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when turf grass areas are water-logged or the soil is saturated with water (i.e. will not allow penetration into the root zone of the plant). Avoid run-off or puddling of irrigation water following application.

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters.

Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

Use only as directed on this label. Do not use on crops used for food or forage.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use before using DYLOX 6.2 Granular Insecticide. **Do not use on turf being grown for sale or other commercial use as sod, or for commercial seed production or for research purposes except for research involving use of this product that is consistent with this label.**

ENTRY RESTRICTIONS

Do not enter or allow others to enter the treated area (except those persons involved in the incorporation) until the incorporation is complete. Incorporation is to be accomplished by watering-in. Do not enter or allow others to enter the treated area until the surface is dry after the watering-in. For home lawns, notify the property owner that he or she must water-in the product promptly or within the same day.

RESTRICTIONS

- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Only protected handlers may be in the area during application.
- Aerial application is prohibited.
- Use this product only on the turf sites permitted by this label.
- Do not apply this product using hand-held or chest-mounted spreaders.
- The applicator must water-in the product after application, or notify the property owner that he or she must water-in after application on the same day.
- Limit the number of applications to turf to 3 per calendar year with minimum retreatment intervals of 7 days.
- Limit the total of all applications to 24.5 lbs of active ingredient (395 lbs of product) per acre per year for controlling mole crickets and grubs and to 16.2 lbs of active ingredient (261 lbs of product) per acre per year for controlling surface feeding insects.
- Apply this product directly to the lawn or garden area. Water treated area as directed on this label.
- Do not water to the point of run-off. Do not make applications during rain.
- Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Additional Restrictions For Turf on Golf Courses	
<ul style="list-style-type: none"> • Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, natural ponds, or estuaries. • Broadcast use is limited to tees and greens. Use on fairways is limited to spot treatments. • Post-application watering-in is mandatory. 	

PRODUCT SPECIFICATIONS	
DYLOX® 6.2 GRANULAR INSECTICIDE	<ul style="list-style-type: none"> • Granular Insecticide containing 6.2% trichlorfon
CONTROLS	<ul style="list-style-type: none"> • White Grub Larvae (such as larvae of Asiatic garden beetle, black turfgrass ataenius, European chafer, green June beetle, Japanese beetle, <i>Aphodius</i> spp., May or June beetles (<i>Phyllophaga</i> spp.), northern and southern masked chafers (<i>Cyclocephala</i> spp.), oriental beetle and sugarcane grubs (<i>Tomarus</i> spp.)) • Mole crickets • Annual bluegrass weevil • Billbug larvae • Sod webworms and cranberry girdler • Cutworms and armyworms • Chinch bugs
WHERE TO APPLY	<ul style="list-style-type: none"> • All types of turf where pests have been seen or found excluding turf where application is not allowed. See IMPORTANT: above.
COVERAGE	<ul style="list-style-type: none"> • One bag (30 lbs) will treat 10,000 sq ft for white grubs, annual bluegrass weevil, chinch bugs, cranberry girdler, and mole crickets. • One bag (30 lbs) will treat 15,000 sq ft for sod webworms, armyworms, and cutworms.
HOW TO APPLY	<ul style="list-style-type: none"> • Do not apply with hand-held or chest-mounted spreader. • Apply granules using a broadcast or drop type spreader.
QUESTIONS	For questions or comments, call toll free 1 (800) 331-2867.

SURFACE FEEDING INSECTS	
Sod webworm, cranberry girdler, cutworm, and armyworm	
Use Rate	<ul style="list-style-type: none"> • 2-3 pounds per 1,000 sq ft of turf. • Refer to chart for spreader settings (Make sure your spreader is calibrated for this product before application).
How to Apply	<ul style="list-style-type: none"> • For properly managed turf remove excessive thatch, which encourages insects. Rake or de-thatch the area and remove the debris to expose insect hiding places. • Water the turf and cut to proper height. • Apply granules with a broadcast or drop type spreader. • To release the active ingredient from the granule, the applicator must water-in after application or notify the property owner that he or she must water-in, on the same day as application, with sufficient water to thoroughly wet the remaining thatch. If sufficient rainfall to thoroughly wet the thatch occurs after application but before watering-in, additional watering-in is not required.
When to Apply	<ul style="list-style-type: none"> • Treat when the insects are present. • To control multiple generations of sod webworm larvae, additional treatments at 30- to 40-day intervals during the summer months may be required. Do not make more than 3 applications per calendar year.

Apply Dylox 6.2G Insecticide uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply Dylox 6.2G Insecticide. Do not use spreaders that would apply the material in concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

Spreader Settings 2 pounds per 1,000 sq ft	
SPREADER	SETTING
Cyclone Pro	4
Cyclone-Roto	3
Lely Broadcast	2.5
Lesco Rotary	11
Prize CB85, CB T85	4.5
Scott's Drop	4
Scott's R7X	H
Spyker Pro	4
Greenview (Model 105)	6.5
Earthway (Model 2600)	8
Sears Broadcast	3.5
Sears Drop	6
(30 pounds will treat 15,000 sq ft at this rate)	

Spreader Settings 3 lb/1,000 sq ft	
Spreader	Setting
Cyclone Pro	4.75
Cyclone-Roto	3.5
Lely Broadcast	2.9
Lesco Rotary	14
Prize CB85, CB T85	5
Scott's Drop	4.5
Scott's R7X	J
Spyker Pro	4.7
Greenview (Model 105)	7.5
Earthway (Model 2600)	10
Sears Broadcast	4
Sears Drop	7
(30 pounds will treat 10,000 sq ft at this rate)	

ROOT FEEDING INSECTS

annual bluegrass weevil (larvae)
 Billbugs (larvae)
 cranberry girdler
 mole crickets
 chinch bugs
white grub larvae: larvae of Asiatic garden beetle, black turfgrass atenius, European chafer, green June beetle, Japanese beetle, *Aphodius spp.*, May or June beetles (*Phyllophaga spp.*), northern & southern masked chafers (*Cyclocephala spp.*), Oriental beetle, and sugarcane grubs (*Tomarus spp.*)

Use Rate	<ul style="list-style-type: none"> • 3 pounds per 1,000 sq ft of turf. • Refer to chart for spreader settings (Make sure your spreader is calibrated for this product before application).
-----------------	--

How to Apply	<ul style="list-style-type: none"> • Ensure that there is no more than 1/2 inch of thatch present at time of application. Under dry conditions, prewater prior to application for control of white grub and weevil larvae. • Water the turf and cut to proper height. • Apply granules with a broadcast or drop type spreader. • The applicator must water-in after application or notify the property owner that he or she must water-in, on the same day as application, with sufficient water to thoroughly wet the remaining thatch and move the product to the root zone. If sufficient rainfall to thoroughly wet the thatch occurs after application but before watering-in, additional watering-in is not required.
---------------------	---

When to Apply	<ul style="list-style-type: none"> • Apply when pests are present. • Consult your Agricultural Extension Service for the best time to treat in your area. • A second treatment may be needed for mature large sized grubs.
----------------------	---

STORAGE AND DISPOSAL	
Do not contaminate water, food, or feed by storage or disposal.	
PESTICIDE STORAGE:	<ul style="list-style-type: none"> • Store in its original container in a cool, dry, locked place out of reach of children.
PESTICIDE DISPOSAL:	<ul style="list-style-type: none"> • Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER HANDLING:	<ul style="list-style-type: none"> • Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling, if available or dispose of empty bag in a sanitary landfill.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Bayer CropScience LP's control it is impossible for Bayer CropScience LP to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, Bayer CropScience LP disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

NET CONTENTS: 30 lbs

PRODUCED FOR



Bayer Environmental Science

**A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709**

Dylox® 6.2G Granular Insecticide (PENDING) 04/01/2014, 05/02/2014, 06/13/2014, 10/01/2014