06/13/2005

S 1

Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval expires 2-28-95 OPP Identifier Number Registration **United States Amendment Environmental Protection Agency** Washington, DC 20460 Other Application for Pesticide - Section I 1. Company/Product Number 2. EPA Product Manager 3. Proposed Classification 432-1428 Dan Kenny None Restricted 4. Company/Product (Name) Allectus 0.15 GC Plus Turf Fertilizer Insecticide 5. Name and Address of Applicant (Include ZIP Code) 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: **Bayer Environmental Science** 95 Chestnut Ridge Road EPA Reg. No. Montvale, NJ 07645 Product Name Check if this is a new address Section II Final printed labels in response to NOTIFICATION Amendment - Explain Below Agency Letter dated Resubmission in response to Agency Letter dated "Me Too" Application. JUN 1 3 2005 Notification - Explain below. Other - explain below. Explanation: Use additional page(s) if necessary. (For section I and Section II.) This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcemnet action and penalties under sections 12 and 14 of FIFRA. - Notification to Add NY Restrictive Language Section III 1. Material This Product Will Be Packaged In: Child-Resistant Packaging Unit Packaging Water Soluble Packaging Type of Container Metal Yes' Yes Plastic Glass No No Paper If yes, No. per lf Yes. Other (Specify) No. per Unit Package wat. container Package wat. container Certification must be submitted. 3. Location of Net Contents Information 4. Size(s) of Retail Container 5. Location of Label Directions On Label Label Container On Labeling accompanying product 6. Manner in Which Label is Affixed to Product Other Lithograph Paper glued Stenciled Section IV 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) Telephone No. (Include Area Code) Richard M. Gorrell (201)-307-3376 Manager, Registrations 6. Date Application Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. Received I acknowledge that any knowingly false for misleading statement may be punishable by fine or imprisonment or (Stainped) both under applicable law. 2. Signature Manager, Registrations Typed Name 5. Date Richard M. Gorrell May 23, 2005

RESTRICTED USE PESTICIDE

Due To Toxicity To Fish and Aquatic Organisms

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

ALLECTUS 0.15 GC PLUS TURF FERTILIZER Insecticide

NOTIFICATION
JUN 1 3 2005

For fertilization and general insect control in turfgrass areas of golf courses and sod farms			
ACTIVE INGREDIENT:			
Imidacloprid*, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine	0.083%		
Bifenthrin**			
OTHER INGREDIENTS:			
	100.000%		
EPA Reg. No. 432-1428	EPA Est. No.		

^{*} Protected by U.S. Patent No. 4,742,060 & 5,783,203

Bifenthrin: U.S. Patent No. 4,238,505

STOP - Read the label before use. Keep out of reach of children.

CAUTION

DOINOLUSE LITIS PRODUCT ON COLT COURSES AND SOFT HARMS IN MASSAU COUNTY OR SUFFOUR COUNTY NEW YORK.

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-800-331-2867

FIRST AID				
lf on skin	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. 			
If in eyes	 Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. 			
Have the conductor	Call a poison control center or doctor for treatment advice. Ontainer or label with you when calling a poison control center or doctor or going for treatment. In case of			

emergency call toll free the Bayer Environmental Science Emergency Response Telephone No. 1-800-334-7577.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

This product contains a pyrethroid. If large amounts have been ingested, milk, cream and other digestible fats and oils may increase absorption and so should be avoided.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated dothing before reuse.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long sleeve shirt and pants
- Shoes plus socks

* Page 1 of 6

^{**}Cis Isomers 97% minimum, trans isomers 3% maximum.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendation:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- · Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Run-off may be hazardous to aquatic organisms in water adjacent to treated areas

This product contains a chemical with properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not formulate this product into other end-use products.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted –entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is

- Coveralls
- Waterproof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENT

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If material is spilled for any reason or cause, carefully contain any spilled material to prevent non-target contamination. Do not walk through spilled material and dispose of as directed for pesticides above. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Environmental Science Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577 or contact Chemtrec at 800-424-9300.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill, by incineration, or if allowed by the state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

APPLICATION TO TURFGRASS

The use of this product will provide plant nutrients as well as providing control of selected insect pests. ALLECTUS 0.15 GC can be used for the control of soil inhabiting pests of turfgrass, such as northern and southern masked chafers, *Cyclocephala borealis*, *C. immaculata*, and/or *C. lurida*; Asiatic garden beetle, *Maladera castanea*; black vine weevil, *Otiorhyncus sulcatus*, European chafer, *Rhizotroqus majalis*; green June beetle, *Cotinis nitida*; May or June beetles, *Phyllophaga* spp.; Japanese beetle, *Popillia japonica*; oriental beetle, *Anomala orientalis*; sugarcane grub, *Euetheola humulis rugiceps*; billbugs, *Spherophorus* spp.; annual bluegrass weevil, *Hyperodes* spp.; black turfgrass ataenius, *Ataenius spretulus*; *Aphodius* spp; European crane fly, *Tipula paludosa*; frit fly, *Oscinella frit*; chinch bugs, *Blissus* spp.; fire ants, *Solenopsis* spp.; cutworms, *Agrotis ipsilon, Peridroma saucia, Nephalodes menians*; armyworms, *Spodoptera* spp., *Pseudaletia* spp.; sod webworm, *Crambus* spp.; and mole crickets, *Scapteriscus* spp. ALLECTUS 0.15 GC can be used as directed on turfgrass in sites such as home lawns, business and office complexes, shopping complexes, multi-family residential complexes, airports, cemeteries, parks, playgrounds, and athletic fields.

The active ingredients ALLECTUS 0.15 GC have sufficient residual activity so that applications for control of subsurface feeders can be made preceding the egg laying activity. The need for an application for control of subsurface feeders can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods. Optimum control of subsurface feeders will be achieved when applications are made prior to egg hatch of the target pest, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

The active ingredients in ALLECTUS 0.15 GC have sufficient knockdown and residual activity to provide curative and residual control of surface feeding pests. Applications for control of surface feeding pests can be made when infestations are anticipated based on historical monitoring of the site, previous records or experience, current season adult trapping or presence of insects at economic thresholds as determined by scouting and/or recommendations of local State extension personnel or other qualified specialists.

Do not apply more than 300 lb. of ALLECTUS 0.15 GC (0.25 lb. of imidacloprid, 0.2 lb. bifenthrin) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 600 lb. of ALLECTUS 0.15 GC (0.5 lb. of imidacloprid, 0.4 lb. bifenthrin) per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Applications should not be made when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. Total amount applied cannot exceed a total of 600 lb (0.5 lb of imidacloprid, 0.4 lb of bifenthrin) per acre per year.

NOTE: Not for use in commercial greenhouses or on grass grown for seed or on golf courses and sod farms. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on lawns and grounds.

APPLICATION EQUIPMENT

Apply ALLECTUS 0.15 GC uniformly over the area being treated with normally used granular application equipment. Both drop-type and rotary-type spreaders may be used to apply ALLECTUS 0.15 GC. Avoid the use of spreaders that apply the material in narrow, concentrated bands. Calibrate application equipment prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly and accurately.

RECOMMENDED APPLICATIONS
(Be sure to read "APPLICATION EQUIPMENT" Section of this label)

CROP	PEST		DOSAGE ALLECTUS 0.15 GC
Turfgrasses	Larvae of subsurface feeders:	Green June beetle	240 to 300 lb per acre
	Asiatic garden beetle	Japanese beetle	(5.5 to 6.9 lb per 1000 ft sq)
	Black turfgrass ataenius	Northern masked chafer	
	Black vine weevil	Oriental beetle	
	European chafer	Phyllophaga spp.	
	European crane fly	Southern masked chafer	
		Sugarcane grub	
	Adults and/or larvae of subsurface and surface feeding pests: Centipedes Fleas Millipedes Ticks	Imported fire ants Mole crickets Pillbugs Sowbugs	192 to 300 lb per acre (4.4 to 6.9 lb per 1000 ft sq) or for heavy infestations, a single annual application of 400 to 500 lb per acre (11.1 to 13.8 lb per 1000 sq ft)
	Adults and/or larvae of surface feeders:		192 to 300 lb per acre
	Annual bluegrass weevil	Frit fly	(4.4 to 6.9 lb per 1000 ft sq)
	Billbugs	Leafhoppers	
	Chinch bugs	Mealybugs	
	Chillion bugs	Nuisance ants	
	Larvae of surface feeders:		120 to 300 lb per acre
	Armyworm	Cutworm	(2.75 to 6.9 lb per 1000 ft sq)
	_	Sod webworms	

Consult your local State Agricultural Experiment Station or State Extension Turf Specialists for more specific information regarding timing of application.

NOTE: For optimum control, irrigation or rainfall should occur within 24 hours after application to move the active ingredient through the thatch. Do not apply more than 600 lb product per acre (0.5 lb of imidacloprid active ingredient, 0.4 lb bifenthrin) per acre per year. Do not apply more than 300 lb product per acre (0.25 lb imidacloprid, 0.2 lb bifenthrin) per application except for mole cricket control. Avoid mowing turf or lawn area until after irrigation or rainfall has occurred so that uniformity of application will not be affected.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of ALLECTUS 0.15 GC if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

Imported Fire ants: ALLECTUS 0.15 GC may be applied as a broadcast treatment at 6.9 lb. per 1000 sq. ft.

Mole crickets: For maximum effectiveness, apply ALLECTUS 0.15 GC as a broadcast treatment at 6.9 lb. per 1000 sq. ft. prior to or during the peak egg hatch period. For application when both adults and/or large nymphs are present and actively tunneling, irrigate prior to application if soil is dry and apply 6.9 lb. per 1000 sq. ft. Rainfall or additional irrigation within 24 hours after application will improve effectiveness of ALLECTUS 0.15 GC. For heavily infested mole cricket areas, use a single seasonal application rate of 480 to 600 lb per acre.



Spreader settings for ALLECTUS 0.15 GC

The following chart can be used as a guide in establishing an accurate rate of ALLECTUS 0.15 GC. Application rates will vary depending upon speed, equipment wear and condition of spreader. Carefully calibrate application equipment prior to commercial use of this product.

SPREADER	SPEED (mph)	SETTINGS		
		120 lb/A	300 lb/A	
Cyclone Way	3			
Earthway 2400 (no information available)				
Gandy Turf Tender 24" (no information available)	3			
Gandy Turf Tender 42" (no information available)	3			
Lely	4.5			
Lesco Broadcast	3			
PSB Prizelawn CBR-III	3			
Scott's Proturf R-8A	3			
Scott's Proturf SR-2000	3			
Scott's Handy Green (obsolete)	3			
Scott's Accupro	3			
Spyker	3			
Vicon Model 402	4.5			

RESTRICTIONS

Avoid runoff or puddling of irrigation water following application. Avoid application of ALLECTUS 0.15 GC to areas which are waterlogged or saturated, which will not allow penetration into the root zone of the plant. Do not apply more than 600 lb (0.5 lb of imidacloprid and 0.4 lb of bifenthrin) per acre per year.

Treated areas may be replanted with any crop specified on an imidacloprid and a bifenthrin label, or with any crop for which a tolerance exists for the two active ingredients.

For crops not listed on any imidacloprid and a bifenthrin label, or for crops for which no tolerances for the active ingredients have been established, a 12-month plant-back interval should be observed.

Do not apply by air.

Not for use in nurseries or commercial greenhouses.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of liability before using this product.

If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Bayer Environmental Science's control it is impossible for Bayer Environmental Science to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages, resulting from the use or handling of this product.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

This Product contains imidacloprid, and it is specially formulated and sold under license from Bayer only for the uses set forth on this label. This active ingredient contained in this product, formulations containing this active ingredient, and/or methods of making or using this active ingredient or its formulations may be the subject of one or more Bayer patents, including but not limited to U.S. Patent No. 4,742,060 and 5,783,203. The purchase and/or use of this product constitutes acceptance of a limited license to use this product only in strict accordance with the instructions set forth on this label. Any use of this product other than as expressly set forth on this label is prohibited and is not licensed through the purchase and/or use of this product and may result in the infringement of one or more Bayer patents and/or other intellectual property rights.

©Bayer AG, 2004

Bayer Environmental Science A Business Group of Bayer CropScience LP 95 Chestnut Ridge Road Montvale, NJ 07645

ALLECTUS 0.15 GC Plus Turf Fert RMG EPA Notif NY Restr 5-23-05

Bayer Environmental Science



May 23, 2005

EPA05RMG0523Allectus0.15GCPlusTFNotifNYRestr Fedex

Mr. Dan Kenny U.S. Environmental Protection Agency Office of Pesticide Programs (7504C) Registration Division (7504C) **Document Processing Desk (Notification)** Room 266A 1801 Bell Street Arlington, VA 22202

RE: Allectus 0.15 GC Plus Turf Fertilizer Insecticide Label (432-1428): Notification to add **NY Restriction**

Dear Mr. Kenny:

Thank you for the help on this label and the other Allectus labels. As per our Science conversation, Bayer Environmental Science (BES) is submitting the revised Allectus GC 95 Chestnut Ridge Road labels as a notification. Since bifenthrin is restricted in the State of New York in Nassau Phone: 201.307.9700 and Suffolk counties, BES has revised the Allectus GC labels to include the exclusion of these counties for sale or use of Allectus GC insecticide products. The restrictive language is found under the Signal Word on each Allectus label.

Bayer Environmental

The following information is included with this letter:

- 8570-1 Application Form
- 5 copies of the revised label

If you have any questions or need anything else, please contact me at (201) 307-3376 or email me at mike.gorrell@bayercropscience.

Sincerely,

Richard M. Gorrell

Encl.

Cc: Joe Conti Mike Ruizzo

Norma Pangilinan

A Business Group of Bayer CropScience