

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (7505P) Ariel Rios Building 1200 Pennsylvania Ave., NW Washington, D.C. 20460		EPA Reg. Number: 432-1462	Date of Issuance: JAN 3 2007
	NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)		Term of Issuance: Conditional	Name of Pesticide Product: Prograss SC Herbicide
Name and Address of Registrant (include ZIP Code): Bayer Environmental Science 2 T.W. Alexander Drive Research Triangle Park, NC 27709				
Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.				
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) and (B) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data. 2. Make the following label change: <ol style="list-style-type: none"> a. A WPS Double Notification Statement must be added within the Agricultural Use Requirements box stating "Notify workers of the application by warning them orally and by posting warning signs at entrances to treated area." b. Revise the EPA Registration Number to read, EPA Reg. No. 432-1462. 3. Submit one copy of the revised final printed label for the record before you release the product for shipment. 				
Signature of Approving Official: Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)			Date: JAN 3 2007	

2 7 9

page 2
EPA Reg. No. 432-1462

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505P)

3 7 9

PROGRASS SC HERBICIDE

Not For Use On Commercial Sod Farms In Arizona
SUSPENSION CONCENTRATE HERBICIDE
For Sale and Use by Professional Applicators Only; Not For Sale or Use by Consumers or Homeowners

ACTIVE INGREDIENT:

ethofumesate: (2-ethoxy-2, 3-dihydro-3, 3-dimethyl-5-benzofuranyl methanesulfonate)*	42%
INERT INGREDIENTS**	58%
TOTAL:	100%

*This product contains 4 lb active ingredient per gallon.
Protected by U. S. Patent No. 4,188,202

EPA Reg. No. 432-XXXX

EPA Est. No.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
JAN 3 2007
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No

KEEP OUT OF REACH OF CHILDREN
CAUTION

432-1462

NET CONTENTS: 2.5 gallon

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

IN CASE OF MEDICAL, ENVIRONMENTAL, OR TRANSPORTATION EMERGENCIES OR INJURIES, CALL 1-800-334-7577 (24 HOURS/DAY)

For Product Information, Call Toll Free: 1-800-331-2867.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (6)].

Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

(except flaggers, or applicators in cockpits, and enclosed cabs)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

On-site Closed Mixing and Loading System Engineering Controls for Liquid Formulations (Dermal Protection Only): Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)] for dermal protection, and must:

- Wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders (this must consist of long-sleeve shirt, long pants, shoes, socks, chemical-resistant gloves, and chemical-resistant apron)
- Wear protective eyewear, if the system operates under pressure
- Be provided and must have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown: chemical-resistant footwear.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

STORAGE AND DISPOSAL

STORAGE: Protect Prograss SC Herbicide from freezing temperatures.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

DO NOT REUSE EMPTY CONTAINER

GENERAL INFORMATION

PROGRASS® SC Herbicide is a suspension concentrate containing 4 lbs. active ingredient per U.S. gallon for use on established perennial ryegrass, Kentucky bluegrass, creeping bentgrass, turf-type tall fescue, St. Augustinegrass, and dormant bermudagrass for the control and/or suppression of specific annual grasses and broadleaf weeds. It may also be used, as directed, after overseeding specific grasses.

PROGRASS SC is intended for professional use only on the ornamental turf of sites such as golf courses, sod farms, parks, cemeteries, lawns of commercial establishments and homeowner lawns (see Use Precautions below).

PROGRASS SC has both preemergent and early (two-leaf stage) postemergent activity; it is used most effectively in programs emphasizing both approaches.

WEED SPECIES CONTROLLED

PREEMERGENCE

Annual Grasses

Annual bluegrass.....Poa annua
 Large crabgrass.....Digitaria sanguinalis
 Smooth crabgrass.....Digitaria ischaemum
 Barnyardgrass.....Echinochloa crus-galli
 Green foxtail.....Setaria viridis
 Yellow foxtail.....Setaria glauca
 Canarygrass.....Phalaris canariensis

Annual Broadleaves

Burclover.....Medicago sp.
 Common chickweed.....Stellaria media
 Common purslane.....Portulaca oleracea
 Redroot pigweed.....Amaranthus retroflexus

PROGRASS will also reduce competition from:

Yellow nutsedge.....Cyperus esculentus
 Purple nutsedge.....Cyperus rotundus

POSTEMERGENCE

Annual Grasses

Annual bluegrass.....Poa annua

Annual Broadleaves

Common chickweed.....Stellaria media
 White clover.....Trifolium repens

PREEMERGENCE/EARLY POSTEMERGENCE

Annual Grasses

Smooth crabgrass.....Digitaria ischaemum
 Large crabgrass.....Digitaria sanguinalis

USE PRECAUTIONS

- PROGRASS SC should be used only for recommended purposes and at the recommended rates (DO NOT OVERAPPLY).
- Avoid spray overlap or turf injury may occur.
- Use of a spray colorant or indicator in the spray tank is recommended so that spray pattern overlapping can be avoided.
- Do not apply with flood jet nozzles and hand-held sprayers, since treatments may not be uniform.
- Do not apply this product through any type of irrigation system.
- PROGRASS SC application is most effective on healthy, vigorously growing turf.
- PROGRASS SC may be applied to homeowner lawns by licensed or certified applicators. PROGRASS SC is not intended for use by home-owners.
- Overseeding is recommended in conjunction with PROGRASS SC applications to achieve conversion to desired turfgrass species and to avoid stand thinning due to annual bluegrass loss.
- When overseeding, use the rate of PROGRASS SC recommended for the overseeded species. In mixed stands of established turfgrasses, use the rate recommended for the least tolerant species.
- Do not apply PROGRASS SC within 8 weeks following the application of a Plant Growth Regulator. A PROGRASS SC program may be initiated on creeping bentgrass 3 weeks after a single application of PRIMO® has been applied.
- PROGRASS SC application is recommended for golf course fairways, roughs, and tees but at fairway-height only. Do not apply to putting greens.
- Do not apply PROGRASS SC to zoysiagrass and hard or fine fescue; serious injury may result.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not graze livestock on treated turf. Do not feed treated grass clippings to livestock. Read entire Use Precautions, Directions for Use, and Disclaimer of Warranties before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

For Sod Farm Turf, do not harvest treated sod for 16 days following application.

Spray Drift Management: This chemical can contaminate surface water through spray drift. A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph.

Temperature Inversions:

Do not make applications into areas of temperature inversion of stable atmospheric conditions.

Shake container well before using.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops except turf grown for sod. The REI for turf grown for sod is 9 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of treated area until sprays have dried.

Professional pesticide applicators applying to residential turf, including home lawns, parks, and recreation areas must inform their customers that all persons and pets must be kept off the treated turf until sprays have dried.

SPRAY EQUIPMENT, APPLICATION, AND PRECAUTIONS

Apply PROGRASS SC Herbicide using standard, low pressure (20 to 50 psi) spray equipment. Spray equipment should be carefully calibrated before use and checked frequently during application. Addition of a spray indicator may improve treatment uniformity. The tank and line should be cleaned and rinsed before and after using PROGRASS SC. Do not allow spray mixture to stand in tank overnight. Store unused spray mixture in tightly-sealed containers and protect from frost.

VARIETAL TOLERANCE

The following turfgrass cultivars have shown tolerance to PROGRASS SC Herbicide; however, all species and varieties have not been tested. PROGRASS SC has been used safely on the following turfgrass cultivars, but its use is not strictly limited to these. Before large scale use of PROGRASS SC on other cultivars, a tolerance test on a small area is recommended.

PERENNIAL RYEGRASS—Acclaim, Blazer, Dasker, Derby, Elka, Fiesta, Goalie, Hunter, Linn, Loretta, Manhattan II, Palmer, Pennfine, Regal, Yorktown

KENTUCKY BLUEGRASS—Adelphi, American, Aspen, Asset, Challenger, Classic, Emundi, Huntsville, Georgetown, Glade, Haga, Julic, Liberty, Merit, Midnight, Monopoly, Mystic, Parade, Rugby, Sydsport, Touchdown.

DO NOT APPLY to the following cultivars: Northstar, Explorer, Limousine, RAN I and Total Eclipse.

CREEPING BENTGRASS—Carmen, Cobra, Highland, Lopez, Mariner, National, Penncross, Providence, Putter, Southshore, SR1020, and Viper have shown good to excellent tolerance to PROGRASS SC Herbicide. Injury has been noted on Emerald, Penneagle, and Pennlinks occasionally. PROGRASS SC may cause serious injury to Cohansey, Colonial, South German varieties, Egmont, Bardot, Tracenta, Allure, Astoria, and SR 7100.

TURF-TYPE TALL FESCUE—America, Arid, Mustang

ST. AUGUSTINEGRASS—Raleigh

DORMANT BERMUDAGRASS—Tifgreen, Tidwarf. Common bermudagrass is more susceptible to PROGRASS injury than hybrid bermudagrass.

GENERAL RECOMMENDATIONS

Fall applications to turfgrass sites to control annual bluegrass are most effective if application is initiated during the period of maximum weed germination and ended as close to the first killing freeze as possible. Spring applications may be made during the period of maximum weed germination. Consult the university extension service or weed science specialist for these dates in your locality. Spring applications are most effective following fall applications.

SEE CHART FOR USE RATES AND TIMINGS. READTEXT BELOW FORSPECIFIC RECOMMENDATIONS FOR EACH TYPE OF GRASS!

Cool Season Turfgrasses

I. Perennial Ryegrass

Application to perennial ryegrass is recommended in fall and/or spring during the primary period of annual bluegrass germination, and up to 30 days after annual bluegrass emergence. Annual bluegrass is easiest to control when young and newly emerged. Spring applications should be used in conjunction with fall applications. Consult Extension Service or University weed specialists for date(s) annual bluegrass germinates in your area.

When overseeding with perennial ryegrass, apply PROGRASS SC Herbicide to the seeded area 1 to 2 weeks after emergence of the ryegrass seedlings, when the seedlings are approximately 1" tall or after the first cutting. For overseeding, the soil should be prepared according to recommended practice. Remove or thoroughly work thatch, as it can reduce the effectiveness of the PROGRASS SC treatment. Delay application of straw or other mulch until after PROGRASS SC application. Application to overseeded perennial ryegrass in the fall or spring will also reduce competition from selected broadleaf weeds and crabgrass.

II. Kentucky Bluegrass

Apply PROGRASS SC Herbicide to bluegrass not less than 8 weeks after emergence. Bluegrass overseeded with ryegrass may be treated one to two weeks after ryegrass emergence.

Do not reseed with Kentucky bluegrass until 6 weeks after the last PROGRASS SC application. For fall application programs of PROGRASS SC, alternatives are to seed dormant bluegrass in the late fall or to delay reseeding until the spring. Kentucky bluegrass sod quality during the spring may be reduced following fall applications of PROGRASS SC.

III. Creeping Bentgrass

PROGRASS SC Herbicide may be applied safely to bentgrass cut at fairway height. PROGRASS SC use should be avoided in poorly drained and/or heavily shaded turf areas which are more prone to herbicide injury. Tank mixing with soluble or controlled release nitrogen at (.1 to .25 lb N per 11000 ft²) with PROGRASS SC may improve creeping bentgrass turf tolerance and improve creeping bentgrass conversion.

A spring application of PROGRASS SC following fall applications is useful in further reducing populations of annual bluegrass. Apply PROGRASS SC in the spring after creeping bentgrass has resumed active growth and has achieved full green-up.

Do not apply PROGRASS SC to renovated or overseeded bentgrass fairway turf less than three (3) to four (4) weeks after seedling emergence.

Do not overseed with creeping bentgrass until three (3) to four (4) weeks after the last PROGRASS SC application.

IV. Turf-Type Tall Fescue

Apply PROGRASS SC Herbicide to turf-type tall fescue in the fall. If overseeding with turf-type tall fescue, PROGRASS SC application and the seeding may be simultaneous.

Warm Season Turfgrasses

I. St. Augustinegrass (Established Turf)

Apply PROGRASS SC Herbicide to St. Augustinegrass in the spring and early summer to suppress bermudagrass and Bermudagrass seedhead formation or control annual bluegrass. Best results are achieved when multiple applications are made.

PROGRASS SC suppresses the development of actively growing Bermudagrass in established St. Augustine Sod. Begin PROGRASS SC applications when Bermudagrass first breaks dormancy in the spring. Application timing is critical to achieve optimum results and will vary depending on location and temperature. A second application of PROGRASS SC should be made 21-28 days after the first application. For optimum suppression, a third application may be necessary 21-28 days after the second application. Temporary stunting and minor discoloration of St. Augustinegrass may occur after application. To improve suppression, PROGRASS SC may be tank-mixed with Atrazine in each application (Atrazine at up to 2 lb ai/A for first application and 0.75 lb – 1 lb ai/A atrazine for second or third application). DO NOT treat St Augustinegrass that has not been established for six months. DO NOT treat St. Augustinegrass that is under stress or injury may occur. If the St. Augustinegrass shows signs of severe yellowing or stunting, discontinue applications.

II. Overseeded Dormant Bermudagrass

Apply PROGRASS SC Herbicide to overseeded bermudagrass in late fall, one to two weeks after emergence of overseeded perennial ryegrass to control annual bluegrass. Treatment to turf that is not fully dormant may cause early injury and/or delayed spring green-up. Applications of 1 to 1.125 quarts of PROGRASS SC per acre have shown acceptable control of annual bluegrass with minimum injury to nondormant (or predormant) bermudagrass when applied in late November or early December. Lower rates have shown lesser annual bluegrass control and higher rate have shown more bermudagrass injury when the bermudagrass is not fully dormant. The onset of dormancy will vary according to location and yearly weather. One or two supplemental applications at 21 to 28 day intervals may be required to maintain control through the winter months. DO NOT APPLY more than two applications of PROGRASS SC when treating nondormant bermudagrass. The final PROGRASS SC application should be made to bermudagrass no later than February 1st or 4 weeks prior to breaking of winter dormancy. Late applications (within 4 weeks of breaking dormancy) to turf that will be breaking dormancy may temporarily delay resumption of active growth (spring green-up) of bermudagrass in the spring. The breaking of winter dormancy will vary according to location and yearly weather. Application of PROGRASS SC Herbicide to bermudagrass that is stressed due to shade, poorly drained soils, and high traffic may result in increased turf injury. Avoid use of PROGRASS SC or use lower rates under these conditions. Care should be taken to keep the PROGRASS SC in the overseeded areas during the applications (particularly when making applications to nondormant bermudagrass) so that the ryegrass will mask any early injury or late transition in the spring. Higher seeding rate may also be helpful to minimize a thin stand appearance since annual bluegrass will be removed. Common bermudagrass in fairways or roughs may be more susceptible to herbicide injury than hybrid bermudagrass.

RECOMMENDED RATES AND TIMINGS ¹								
Turf Type	Main Target Pests	Start	# Apps	Application INT (Days)	Use Rate ²		Overseeding Safety Interval ³ (Weeks After Treatment)	Application Safety Interval ⁵ (Weeks After Emergence)
					Qt/Acre	Oz/ 1000Sq.Ft.		
Ryegrass	Annual Bluegrass	Fall Spring ⁴	2-3 1-2	21 - 28 21 - 28	1-2	.7-1.5	1-2	1-2
Kentucky Bluegrass	Annual Bluegrass	Fall	2 - 3	21 - 28	.75	.6	6	8
Creeping Bentgrass	Annual Bluegrass	Fall Spring ⁴	2-3 1-2	21 - 28 21 - 28	.75 .75	.6 .6	4	4
Turf-Type Tall Fescue	Annual Bluegrass	Fall	2 - 3	21 - 28	.75-1.5	.6-1.1	0	2-3
St. Augustinegrass (Est. Turf)	Bermudagrass Suppression	Spring	2 - 3	21 - 28	1.5-3	1-2.2	N/A	N/A
Overseeded (ryegrass) Bermudagrass:								
Dormant Bermudagrass	Annual Bluegrass	Fall	2 - 3	21 - 28	.75-2	.6-1.5	N/A	N/A
Nondormant Bermudagrass	Annual Bluegrass	Fall	2	21 - 28	1	.7	N/A	N/A

¹Refer to label text for further instructions regarding each grass species.

²Apply recommended rate of PROGRASS SC in 20 to 60 gallons of water per acre or 1 to 3 gallons of water per 1000 Ft².

³Interval applies to overseeding specific grass on same type of grass after last PROGRASS SC treatment.

⁴Use spring application after previous fall treatments only.

⁵Refers to interval in which PROGRASS SC may be safely applied to specific grass following seed emergence.

NA - Not applicable.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer

DISCLAIMER OF WARRANTIES: BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT ALLOWED BY LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG, 2005

TRADEMARK INFORMATION

Chipco and Prograss are registered trademarks of Bayer.

Backed By Bayer is a trademark of Bayer.

Bayer Environmental Science
A Business Group of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709

PROGRASS SC HERBICIDE (PENDING) Revised as per EPA 12-21-06