

432-1462

05/15/2009

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Richard M. Gorrell  
Manager, Registration  
Bayer Advanced  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709

MAY 15 2009

Subject: Label Notification(s) for Pesticide Registration Notices 2007-4

Dear Mr. Gorrell:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 and 2008-1 dated April 23, 2009 for:

**EPA Registration 432-1462**

**Prograss SC Herbicide**

The Registration Division (RD) has conducted a review of this request for applicability under PR Notice 2007-4 and 2008-1 and finds that the label changes requested falls within the scope of PR Notice 2007-4 and 2008-1. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Nicole Williams of my staff at 703-308-5551.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs

2/11



United States  
**Environmental Protection Agency**  
 Washington, DC 20460

Registration  
 Amendment  
 Other

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number 432-1462	2. EPA Product Manager Ms. Joanne Miller	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Prograss SC Herbicide	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Bayer Environmental Science 2 T. W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. <u>NOTIFICATION</u> Product Name <u>MAY 15 2009</u>	

**Section - II**

Amendment - Explain below.  Final printed labels in response to Agency letter dated \_\_\_\_\_  
 Resubmission in response to Agency letter dated \_\_\_\_\_  "Me Too" Application.  
 Notification - Explain below.  Other - Explain below.

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance of PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and it may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.  
 - Notification to add the revised Storage and Disposal Statement

**Section - III**

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container

3. Location of Net Contents Information  
 Label  Container

4. Size(s) Retail Container  
1 gallon

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product  
 Lithograph Paper glued Stenciled  Other encased in plastic glued to jug; fold out/accordion

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Richard M. Gorrell	Title Manager, Registrations	Telephone No. (Include Area Code) (919) 549-2423
----------------------------	---------------------------------	---

**Certification**  
 I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature <i>Richard M. Gorrell</i>	3. Title Manager, Registrations	6. Date Application Received (Stamped)
4. Typed Name Richard M. Gorrell	5. Date April 23, 2009	

OK

3/11

# Bayer Environmental Science



April 23, 2009

EPA09RMG0423PrograssSCNotifStorDisp  
Fedex

Ms. Linda Arrington  
U.S. Environmental Protection Agency  
Office of Pesticide Programs (7504P)  
Registration Division (7504P)  
Document Processing Desk (Me-Too)  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202-4501

RE: Prograss SC Herbicide (432-1462): Notification to Add the EPA Mandated Storage and Disposal Statement

Dear Ms. Linda Arrington:

Bayer Environmental Science (BES) is notifying the Agency of the revision to the Storage and Disposal Statement as mandated by the EPA. The revised wording is the same as given by the EPA example and should be reviewed as a notification.

Bayer Environmental  
Science  
2 T.W. Alexander Drive  
RTP, NC 27709  
Phone : 919.549.2000

The following information is included with this letter:

- EPA Application Form 8570-1
- 3 copies of the Prograss SC Herbicide label

If you have any questions please contact me at (919) 549-2423 or email me at [mike.gorrell@bayercropscience](mailto:mike.gorrell@bayercropscience).

Sincerely,

Richard M. Gorrell

Incl.

Cc: Karen Shearer  
Matt Bradley

A Business Group of  
Bayer CropScience

4/11

# PROGRASS SC HERBICIDE

NOTIFICATION

MAY 15 2009

Not For Use On Commercial Sod Farms In Arizona  
SUSPENSION CONCENTRATE HERBICIDE

For Sale and Use By Professional Applicators Only; Not for Sale or Use by Consumers or Homeowners

**ACTIVE INGREDIENT:**

ethofumesate: (2-ethoxy-2, 3-dihydro-3, 3-dimethyl-5-benzofuranyl methanesulfonate)\*.....42%

INERT INGREDIENTS\*\*.....58%

TOTAL:.....100%

\*This product contains 4 lb active ingredient per gallon.  
Protected by U. S. Patent No. 4,188,202

EPA Reg. No. 432-1462

EPA Est. No.

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

NET CONTENTS: 2.5 gallon

FIRST AID	
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**IN CASE OF MEDICAL, ENVIRONMENTAL, OR TRANSPORTATION EMERGENCIES OR INJURIES, CALL 1-800-334-7577 (24 HOURS/DAY)**

For Product Information, Call Toll Free: 1-800-331-2867.

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPA) for agricultural pesticides [40 CFR 170.240 (d) (6)].

### Personal Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- Shoes plus socks

(except flaggers, or applicators in cockpits, and enclosed cabs)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### ENGINEERING CONTROLS STATEMENT

**On-site Closed Mixing and Loading System Engineering Controls for Liquid Formulations (Dermal Protection Only):** Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)] for dermal protection, and must:

- Wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders (this must consist of long-sleeve shirt, long pants, shoes, socks, chemical-resistant gloves, and chemical-resistant apron)
- Wear protective eyewear, if the system operates under pressure
- Be provided and must have immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown: chemical-resistant footwear.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

This pesticide may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

## STORAGE AND DISPOSAL

**STORAGE:** Protect Prograss SC Herbicide from freezing temperatures.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

### GENERAL INFORMATION

PROGRASS® SC Herbicide is a suspension concentrate containing 4 lbs. active ingredient per U.S. gallon for use on established perennial ryegrass, Kentucky bluegrass, creeping bentgrass, turf-type tall fescue, St. Augustinegrass, and dormant bermudagrass for the control and/or suppression of specific annual grasses and broadleaf weeds. It may also be used, as directed, after overseeding specific grasses.

PROGRASS SC is intended for professional use only on the ornamental turf of sites such as golf courses, sod farms, parks, cemeteries, lawns of commercial establishments and homeowner lawns (see Use Precautions below).

PROGRASS SC has both preemergent and early (two-leaf stage) postemergent activity; it is used most effectively in programs emphasizing both approaches.

### WEED SPECIES CONTROLLED

#### PREEMERGENCE

##### Annual Grasses

- Annual bluegrass.....*Poa annua*
- Large crabgrass.....*Digitaria sanguinalis*
- Smooth crabgrass.....*Digitaria ischaemum*
- Barnyardgrass.....*Echinochloa crus-galli*
- Green foxtail.....*Setaria viridis*
- Yellow foxtail.....*Setaria glauca*
- Canarygrass.....*Phalaris canariensis*

##### Annual Broadleaves

- Burclover.....*Medicago sp.*
- Common chickweed.....*Stellaria media*
- Common purslane.....*Portulaca oleracea*
- Redroot pigweed.....*Amaranthus retroflexus*

PROGRASS will also reduce competition from:

- Yellow nutsedge.....*Cyperus esculentus*
- Purple nutsedge.....*Cyperus rotundus*

#### POSTEMERGENCE

##### Annual Grasses

- Annual bluegrass.....*Poa annua*
- Annual Broadleaves
- Common chickweed.....*Stellaria media*
- White clover.....*Trifolium repens*

#### PREEMERGENCE/EARLY POSTEMERGENCE

##### Annual Grasses

- Smooth crabgrass.....*Digitaria ischaemum*
- Large crabgrass.....*Digitaria sanguinalis*

## USE PRECAUTIONS

- PROGRASS SC should be used only for recommended purposes and at the recommended rates (DO NOT OVERAPPLY).
- Avoid spray overlap or turf injury may occur.
- Use of a spray colorant or indicator in the spray tank is recommended so that spray pattern overlapping can be avoided.
- Do not apply with flood jet nozzles and hand-held sprayers, since treatments may not be uniform.
- Do not apply this product through any type of irrigation system.
- PROGRASS SC application is most effective on healthy, vigorously growing turf.
- PROGRASS SC may be applied to homeowner lawns by licensed or certified applicators. PROGRASS SC is not intended for use by home-owners.
- Overseeding is recommended in conjunction with PROGRASS SC applications to achieve conversion to desired turfgrass species and to avoid stand thinning due to annual bluegrass loss.
- When overseeding, use the rate of PROGRASS SC recommended for the overseeded species. In mixed stands of established turfgrasses, use the rate recommended for the least tolerant species.
- Do not apply PROGRASS SC within 8 weeks following the application of a Plant Growth Regulator. A PROGRASS SC program may be initiated on creeping bentgrass 3 weeks after a single application of PRIMO® has been applied.
- PROGRASS SC application is recommended for golf course fairways, roughs, and tees but at fairway-height only. Do not apply to putting greens.
- Do not apply PROGRASS SC to zoysiagrass and hard or fine fescue; serious injury may result.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not graze livestock on treated turf. Do not feed treated grass clippings to livestock. Read entire Use Precautions, Directions for Use, and Disclaimer of Warranties before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

For Sod Farm Turf, do not harvest treated sod for 16 days following application.

**Spray Drift Management:** This chemical can contaminate surface water through spray drift. A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

### Wind Speed

Do not apply at wind speeds greater than 15 mph.

### Temperature Inversions:

Do not make application into areas of temperature inversion of stable atmospheric conditions.

**Shake container well before using.**

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours for all crops except turf grown for sod. The REI for turf grown for sod is 9 days.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep unprotected persons out of treated area until sprays have dried.

Professional pesticide applicators applying to residential turf, including home lawns, parks, and recreation areas must inform their customers that all persons and pets must be kept off the treated turf until sprays have dried.

## SPRAY EQUIPMENT, APPLICATION, AND PRECAUTIONS

Apply PROGRASS SC Herbicide using standard, low pressure (20 to 50 psi) spray equipment. Spray equipment should be carefully calibrated before use and checked frequently during application. Addition of a spray indicator may improve treatment uniformity. The tank and line should be cleaned and rinsed before and after using PROGRASS SC. Do not allow spray mixture to stand in tank overnight. Store unused spray mixture in tightly-sealed containers and protect from frost.

## VARIETAL TOLERANCE

The following turfgrass cultivars have shown tolerance to PROGRASS SC Herbicide; however, all species and varieties have not been tested. PROGRASS SC has been used safely on the following turfgrass cultivars, but its use is not strictly limited to these. Before large scale use of PROGRASS SC on other cultivars, a tolerance test on a small area is recommended.

**PERENNIAL RYEGRASS**—Acclaim, Blazer, Dasker, Derby, Elka, Fiesta, Goalie, Hunter, Linn, Loretta, Manhattan II, Palmer, Pennfine, Regal, Yorktown

**KENTUCKY BLUEGRASS**—Adelphi, American, Aspen, Asset, Challenger, Classic, Emundi, Huntsville, Georgetown, Glade, Haga, Julic, Liberty, Merit, Midnight, Monopoly, Mystic, Parade, Rugby, Sydsport, Touchdown.

**DO NOT APPLY** to the following cultivars: Northstar, Explorer, Limousine, RAN I and Total Eclipse.

**CREEPING BENTGRASS**—Carmen, Cobra, Highland, Lopez, Mariner, National, Penncross, Providence, Putter, Southshore, SR1020, and Viper have shown good to excellent tolerance to PROGRASS SC Herbicide. Injury has been noted on Emerald, Penneagle, and Pennlinks occasionally. PROGRASS SC may cause serious injury to Cohansey, Colonial, South German varieties, Egmont, Bardot, Tracenta, Allure, Astoria, and SR 7100.

**TURF-TYPE TALL FESCUE**—America, Arid, Mustang

**ST. AUGUSTINEGRASS**—Raleigh

**DORMANT BERMUDAGRASS**—Tifgreen, Tidwarf. Common bermudagrass is more susceptible to PROGRASS injury than hybrid bermudagrass.

## GENERAL RECOMMENDATIONS

Fall applications to turfgrass sites to control annual bluegrass are most effective if application is initiated during the period of maximum weed germination and ended as close to the first killing freeze as possible. Spring applications may be made during the period of maximum weed germination. Consult the university extension service or weed science specialist for these dates in your locality. Spring applications are most effective following fall applications.

**SEE CHART FOR USE RATES AND TIMINGS. READ TEXT BELOW FOR SPECIFIC RECOMMENDATIONS FOR EACH TYPE OF GRASS!**

### Cool Season Turfgrasses

#### I. Perennial Ryegrass

Application to perennial ryegrass is recommended in fall and/or spring during the primary period of annual bluegrass germination, and up to 30 days after annual bluegrass emergence. Annual bluegrass is easiest to control when young and newly emerged. Spring applications should be used in conjunction with fall applications. Consult Extension Service or University weed specialists for date(s) annual bluegrass germinates in your area.

When overseeding with perennial ryegrass, apply PROGRASS SC Herbicide to the seeded area 1 to 2 weeks after emergence of the ryegrass seedlings, when the seedlings are approximately 1" tall or after the first cutting. For overseeding, the soil should be prepared according to recommended practice. Remove or thoroughly work thatch, as it can reduce the effectiveness of the PROGRASS SC treatment. Delay application of straw or other mulch until after PROGRASS SC application. Application to overseeded perennial ryegrass in the fall or spring will also reduce competition from selected broadleaf weeds and crabgrass.

**II. Kentucky Bluegrass**

Apply PROGRASS SC Herbicide to bluegrass not less than 8 weeks after emergence. Bluegrass overseeded with ryegrass may be treated one to two weeks after ryegrass emergence. Do not reseed with Kentucky bluegrass until 6 weeks after the last PROGRASS SC application. For fall application programs of PROGRASS SC, alternatives are to seed dormant bluegrass in the late fall or to delay reseeding until the spring. Kentucky bluegrass sod quality during the spring may be reduced following fall applications of PROGRASS SC.

**III. Creeping Bentgrass**

PROGRASS SC Herbicide may be applied safely to bentgrass cut at fairway height. PROGRASS SC use should be avoided in poorly drained and/or heavily shaded turf areas which are more prone to herbicide injury. Tank mixing with soluble or controlled release nitrogen at (0.1 to 0.25 lb N per 11,000 sq ft) with PROGRASS SC may improve creeping bentgrass turf tolerance and improve creeping bentgrass conversion. A spring application of PROGRASS SC following fall applications is useful in further reducing populations of annual bluegrass. Apply PROGRASS SC in the spring after creeping bentgrass has resumed active growth and has achieved full green-up. Do not apply PROGRASS SC to renovated or overseeded bentgrass fairway turf less than three (3) to four (4) weeks after seedling emergence. Do not overseed with creeping bentgrass until three (3) to four (4) weeks after the last PROGRASS SC application.

**IV. Turf-Type Tall Fescue**

Apply PROGRASS SC Herbicide to turf-type tall fescue in the fall. If overseeding with turf-type tall fescue, PROGRASS SC application and the seeding may be simultaneous.

**Warm Season Turfgrasses**

**I. St. Augustinegrass (Established Turf)**

Apply PROGRASS SC Herbicide to St. Augustinegrass in the spring and early summer to suppress bermudagrass and bermudagrass seedhead formation or control annual bluegrass. Best results are achieved when multiple applications are made. PROGRASS SC suppresses the development of actively growing bermudagrass in established St. Augustine sod. Begin PROGRASS SC applications when bermudagrass first breaks dormancy in the spring. Application timing is critical to achieve optimum results and will vary depending on location and temperature. A second application of PROGRASS SC should be made 21-28 days after the first application. For optimum suppression, a third application may be necessary 21-28 days after the second application. Temporary stunting and minor discoloration of St. Augustinegrass may occur after application. To improve suppression, PROGRASS SC may be tank-mixed with atrazine in each application (Atrazine at up to 2 lb ai/A for first application and 0.75 lb – 1 lb ai/A atrazine for second or third application). DO NOT treat St Augustinegrass that has not been established for six months. DO NOT treat St. Augustinegrass that is under stress or injury may occur. If the St. Augustinegrass shows signs of severe yellowing or stunting, discontinue applications.

**II. Overseeded Dormant Bermudagrass**

Apply PROGRASS SC Herbicide to overseeded bermudagrass in late fall, one to two weeks after emergence of overseeded perennial ryegrass to control annual bluegrass. Treatment to turf that is not fully dormant may cause early injury and/or delayed spring green-up. Applications of 1 to 1.125 qt of PROGRASS SC per acre have shown acceptable control of annual bluegrass with minimum injury to nondormant (or predormant) bermudagrass when applied in late November or early December. Lower rates have shown lesser annual bluegrass control and higher rate have shown more bermudagrass injury when the bermudagrass is not fully dormant. The onset of dormancy will vary according to location and yearly weather. One or two supplemental applications at 21 to 28 day intervals may be required to maintain control through the winter months. DO NOT APPLY more than two applications of PROGRASS SC when treating nondormant bermudagrass. The final PROGRASS SC application should be made to bermudagrass no later than February 1<sup>st</sup> or 4 weeks prior to breaking of winter dormancy. Late applications (within 4 weeks of breaking dormancy) to turf that will be breaking dormancy may temporarily delay resumption of active growth (spring green-up) of bermudagrass in the spring. The breaking of winter dormancy will vary according to location and yearly weather. Application of PROGRASS SC Herbicide to bermudagrass that is stressed due to shade, poorly drained soils, and high traffic may result in increased turf injury. Avoid use of PROGRASS SC or use lower rates under these conditions. Care should be taken to keep the PROGRASS SC in the overseeded areas during the applications (particularly when making applications to nondormant bermudagrass) so that the ryegrass will mask any early injury or late transition in the spring. Higher seeding rate may also be helpful to minimize a thin stand appearance since annual bluegrass will be removed. Common bermudagrass in fairways or roughs may be more susceptible to herbicide injury than hybrid bermudagrass.

**RECOMMENDED RATES AND TIMINGS<sup>1</sup>**

Turf Type	Main Target Pests	Start	# Apps	Application INT (Days)	Qt/A	Use Rate <sup>2</sup>	Overseeding Safety Interval <sup>3</sup> (Weeks After Treatment)	Application Safety Interval <sup>5</sup> (Weeks After Emergence)
						Oz/ 1,000 sq ft		
Ryegrass	Annual Bluegrass	Fall Spring <sup>4</sup>	2-3 1-2	21 - 28 21 - 28	1-2	0.7-1.5	1-2	1-2
Kentucky Bluegrass	Annual Bluegrass	Fall	2 - 3	21 - 28	0.75	0.6	6	8
Creeping Bentgrass	Annual Bluegrass	Fall Spring <sup>4</sup>	2-3 1-2	21 - 28 21 - 28	0.75 0.75	0.6 0.6	4	4
Turf-Type Tall Fescue	Annual Bluegrass	Fall	2 - 3	21 - 28	0.75-1.5	0.6-1.1	0	2-3
St. Augustinegrass (Est. Turf)	Bermudagrass Suppression	Spring	2 - 3	21 - 28	1.5-3	1-2.2	N/A	N/A
Overseeded (ryegrass) Bermudagrass:	Annual Bluegrass	Fall	2 - 3	21 - 28	0.75-2	0.6-1.5	N/A	N/A
Dormant Bermudagrass								
Nondormant Bermudagrass	Annual Bluegrass	Fall	2	21 - 28	1	0.7	N/A	N/A

<sup>1</sup>Refer to label text for further instructions regarding each grass species.  
<sup>2</sup>Apply recommended rate of PROGRASS SC in 20 to 60 gallons of water per acre or 1 to 3 gallons of water per 1,000 sq ft.  
<sup>3</sup>Interval applies to overseeding specific grass on same type of grass after last PROGRASS SC treatment.  
<sup>4</sup>Use spring application after previous fall treatments only.  
<sup>5</sup>Refers to interval in which PROGRASS SC may be safely applied to specific grass following seed emergence.  
 NA - Not applicable.

11/11

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

©2006 Bayer CropScience LP

**TRADEMARK INFORMATION**

Chipco and Prograss are registered trademarks of Bayer.

Backed By Bayer is a trademark of Bayer.

Produced for



**Bayer Environmental Science**

**Bayer Environmental Science  
A Business Group of Bayer CropScience LP  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709**

**PROGRASS SC HERBICIDE (MASTER) EPA APPROVED 01-03-07, Notification 04/16/09**