



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 12, 2018

Mr. Brad Glenn, Ph.D.
Regulator Manager
Bayer CropScience
Environmental Science Division
2 T.W. Alexander Drive
Research Triangle Park, NC 27709

Subject: Label Amendment – Removal of Residential and Park Turf grass
Product Name: ***Prograss SC Herbicide***
EPA Registration Number: 432-1462
Application Date: November 22, 2017
Decision Number: 536260

Dear Dr. Glenn:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 432-1462
Decision No. 536260

with FIFRA section 6. If you have any questions, please contact Eleanor Thornton by phone at 703-305-6799, or via email at Thornton.eleanor@epa.gov.

Sincerely,

A handwritten signature in dark ink, appearing to read "Erik Kraft", written in a cursive style.

Erik Kraft, Product Manager 24
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

PROGRASS® SC HERBICIDE

Not For Use on Commercial Sod Farms in Arizona
SUSPENSION CONCENTRATE HERBICIDE

ACTIVE INGREDIENT:

Ethofumesate: (2-ethoxy-2, 3-dihydro-3, 3-dimethyl-5-benzofuranyl methanesulfonate)*42%

OTHER INGREDIENTS:58%

TOTAL:100%

*This product contains 4 lb active ingredient per gallon.

EPA Reg. No. 432-1462

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

See [Back][Side] Panel for First Aid Instructions and [Leaflet][Booklet] for Complete Precautionary Statements and Directions for Use.
 (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

FIRST AID

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

IN CASE OF MEDICAL, ENVIRONMENTAL, OR TRANSPORTATION EMERGENCIES OR INJURIES, CALL 1-800-334-7577 (24 HOURS/DAY)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing.

Personal Protective Equipment (PPE)

All mixers, loaders, applicators and other handlers (except applicators in enclosed cabs) must wear:

- Long-sleeved shirt and long pants
 - Chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
 - Shoes plus socks
- See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

ACCEPTED

06/12/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 432-1462

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire Use Precautions, Directions for Use, and Disclaimer of Warranties before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency in your State responsible for pesticide regulation.

For Sod Farm Turf, do not harvest treated sod for 3 days following application.

Spray Drift Management: This chemical can contaminate surface water through spray drift. A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph.

Temperature Inversions:

Do not make application into areas of temperature inversion of stable atmospheric conditions.

Shake container well before using.

Resistance Management

For resistance management, PROGRASS® SC Herbicide is a Group 8 herbicide. Any weed population may contain or develop plants naturally resistant to PROGRASS SC Herbicide and other Group 8 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies should be followed.

To delay herbicide resistance take one or more of the following steps:

- Rotate the use of PROGRASS SC Herbicide or other Group 8 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use.
- Scout before and after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other areas by cleaning equipment when moving between treated areas.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific weed biotypes.

- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

Tank Mix

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until sprays have dried.

USE RESTRICTIONS

- Use PROGRASS SC Herbicide only as directed at the specified rates (DO NOT OVERAPPLY).
- Do not apply with flood jet nozzles and hand-held sprayers, since treatments may not be uniform.
- Do not apply this product through any type of irrigation system.
- PROGRASS SC Herbicide is not intended for use by home-owners.
- Do not apply PROGRASS SC Herbicide within 8 weeks following the application of a Plant Growth Regulator. A PROGRASS SC Herbicide program may be initiated on creeping bentgrass 3 weeks after a single application of PRIMO® has been applied.
- PROGRASS SC Herbicide application is specified for golf course fairways, roughs, and tees but at fairway-height only. Do not apply to putting greens.
- Do not apply PROGRASS SC Herbicide to zoysiagrass and hard or fine fescue; serious injury may result.
- For sod farm turf: Do not harvest treated sod for 3 days following application.
- Do not graze livestock on treated turf.
- Do not feed treated grass clippings to livestock.
- Do not move hand set irrigation equipment within 7 days of application except as permitted by the Worker Protection Standard [170.603(d)].
- Do not apply more than 1.7 fl oz product/1000 sq ft. (2.3 lb a.i./A) per application.
- Do not exceed 6 fl oz PROGRASS SC / 1000 sq ft. (equivalent to 8.2 lb a.i./A) per year.
- Do not exceed 3 applications per year using the 1.7 fl oz / 1000 sq ft. (2.3 lb a.i./A) application rate.
- Do not exceed 4 applications per year using the 1.5 fl oz/1000 sq ft. (2.04 lb a.i./A) application rate.
- Do not exceed 16 applications per year using the 0.37 fl oz/1000 sq ft. (0.5 lb a.i./A) application rate.
- The minimum re-treatment interval (RTI) is 14 days. See table for individual turf type rates and re-treatment timings.

USE PRECAUTIONS

- Avoid spray overlap or turf injury may occur.
- Use of a spray colorant or indicator in the spray tank is instructed so that spray pattern overlapping can be avoided.
- PROGRASS SC Herbicide application is most effective on healthy, vigorously growing turf.

- Overseeding is directed in conjunction with PROGRASS SC Herbicide applications to achieve conversion to desired turfgrass species and to avoid stand thinning due to annual bluegrass loss.
- When overseeding, use the rate of PROGRASS SC Herbicide specified for the overseeded species. In mixed stands of established turfgrasses, use the rate specified for the most sensitive species.

PRODUCT INFORMATION

PROGRASS SC Herbicide is a suspension concentrate containing 4 lbs. active ingredient per U.S. gallon for use on established perennial ryegrass, Kentucky bluegrass, creeping bentgrass, turf-type tall fescue, St. Augustinegrass, and dormant bermudagrass for the control and/or suppression of specific annual grasses and broadleaf weeds. It may also be used, as directed, after overseeding specific grasses.

PROGRASS SC Herbicide is intended for commercial use only on the ornamental turf of sites including golf courses, and sod farms (see Use Precautions).

PROGRASS SC Herbicide has both preemergent and early (two-leaf stage) postemergent activity; it is used most effectively in programs emphasizing both approaches.

WEED SPECIES CONTROLLED

PREEMERGENCE

Annual Grasses

Annual bluegrass.....	<i>Poa annua</i>
Large crabgrass.....	<i>Digitaria sanguinalis</i>
Smooth crabgrass.....	<i>Digitaria ischaemum</i>
Barnyardgrass.....	<i>Echinochloa crus-galli</i>
Green foxtail.....	<i>Setaria viridis</i>
Yellow foxtail.....	<i>Setaria glauca</i>
Canarygrass.....	<i>Phalaris canariensis</i>

Annual Broadleaves

Burclover.....	<i>Medicago sp.</i>
Common chickweed.....	<i>Stellaria media</i>
Common purslane.....	<i>Portulaca oleracea</i>
Redroot pigweed.....	<i>Amaranthus retroflexus</i>

PROGRASS SC Herbicide will also reduce competition from:

Yellow nutsedge.....	<i>Cyperus esculentus</i>
Purple nutsedge.....	<i>Cyperus rotundus</i>

POSTEMERGENCE

Annual Grasses

Annual bluegrass.....	<i>Poa annua</i>
-----------------------	------------------

Annual Broadleaves

Common chickweed.....	<i>Stellaria media</i>
White clover.....	<i>Trifolium repens</i>

PREEMERGENCE/EARLY POSTEMERGENCE

Annual Grasses

Smooth crabgrass.....	<i>Digitaria ischaemum</i>
Large crabgrass.....	<i>Digitaria sanguinalis</i>

SPRAY EQUIPMENT, APPLICATION, AND PRECAUTIONS

Apply PROGRASS SC Herbicide using standard, low pressure (20 to 50 psi) spray equipment. Spray equipment needs to be carefully calibrated before use and checked frequently during application. Addition of a spray indicator may improve treatment uniformity. The tank and line needs to be cleaned and rinsed before and after using PROGRASS SC Herbicide. Do not allow spray mixture to stand in tank overnight. Store unused spray mixture in tightly-sealed containers and protect from frost.

ACCEPTABLE VARIETIES

The following turfgrass cultivars have shown the inherent ability to endure applications of PROGRASS SC Herbicide; however, all species and varieties have not been tested. PROGRASS SC Herbicide has been used safely on the following turfgrass cultivars, but its use is not strictly limited to these. Before large scale use of PROGRASS SC Herbicide on other cultivars, testing for turf endurance on a small area is advised.

PERENNIAL RYEGRASS—Acclaim, Blazer, Dasker, Derby, Elka, Fiesta, Goalie, Hunter, Linn, Loretta, Manhattan II, Palmer, Pennfine, Regal, Yorktown

KENTUCKY BLUEGRASS—Adelphi, American, Aspen, Asset, Challenger, Classic, Emundi, Huntsville, Georgetown, Glade, Haga, Julic, Liberty, Merit, Midnight, Monopoly, Mystic, Parade, Rugby, Sydsport, Touchdown.

DO NOT APPLY to the following cultivars: Northstar, Explorer, Limousine, RAN I and Total Eclipse.

CREEPING BENTGRASS—Carmen, Cobra, Highland, Lopez, Mariner, National, Penncross, Providence, Putter, Southshore, SR1020, and Viper have shown good to excellent endurance to PROGRASS SC Herbicide. Injury has been noted on Emerald, Penneagle, and Pennlinks occasionally. PROGRASS SC Herbicide may cause serious injury to Cohansey, Colonial, South German varieties, Egmont, Bardot, Tracenta, Allure, Astoria, and SR 7100.

TURF-TYPE TALL FESCUE—America, Arid, Mustang

ST. AUGUSTINEGRASS—Raleigh

DORMANT BERMUDAGRASS—Tifgreen, Tidwarf. Common bermudagrass is more susceptible to PROGRASS SC Herbicide injury than hybrid bermudagrass.

DIRECTIONS FOR SPECIFIC GRASS TYPES

Fall applications to turfgrass sites to control annual bluegrass are most effective if application is initiated during the period of maximum weed germination and ended as close to the first killing freeze as possible. Spring applications may be made during the period of maximum weed germination. Consult the university extension service or weed science specialist for these dates in your locality. Spring applications are most effective following fall applications.

SEE CHART FOR USE RATES AND TIMINGS. READ TEXT BELOW FOR SPECIFIC DIRECTIONS FOR EACH TYPE OF GRASS!

Cool Season Turfgrasses

I. Perennial Ryegrass

Application to perennial ryegrass is advised in fall and/or spring during the primary period of annual bluegrass germination, and up to 30 days after annual bluegrass emergence. Annual bluegrass is easiest to control when young and newly emerged. Spring applications work best when used in conjunction with fall applications. Consult Extension Service or University weed specialists for date(s) annual bluegrass germinates in your area.

When overseeding with perennial ryegrass, apply PROGRASS SC Herbicide to the seeded area 1 to 2 weeks after emergence of the ryegrass seedlings, when the seedlings are approximately 1" tall or after the first cutting. For overseeding, the soil needs to be prepared according to advised practice. Remove or thoroughly work thatch, as it can reduce the effectiveness of the PROGRASS SC Herbicide treatment. Delay application of straw or other mulch until after PROGRASS SC Herbicide application. Application to overseeded perennial ryegrass in the fall or spring will also reduce competition from selected broadleaf weeds and crabgrass.

II. Kentucky Bluegrass

Apply PROGRASS SC Herbicide to bluegrass not less than 8 weeks after emergence. Bluegrass overseeded with ryegrass may be treated one to two weeks after ryegrass emergence.

Do not reseed with Kentucky bluegrass until 6 weeks after the last PROGRASS SC Herbicide application. For fall application programs of PROGRASS SC Herbicide, alternatives are to seed dormant bluegrass in the late fall or to delay reseeding until the spring. Kentucky bluegrass sod quality during the spring may be reduced following fall applications of PROGRASS SC Herbicide.

III. Creeping Bentgrass

PROGRASS SC Herbicide may be applied safely to bentgrass cut at fairway height. PROGRASS SC Herbicide use needs to be avoided in poorly drained and/or heavily shaded turf areas which are more prone to herbicide injury. Tank mixing with soluble or controlled release nitrogen at (0.1 to 0.25 lb N per 11,000 sq ft) with PROGRASS SC Herbicide may improve creeping bentgrass turf endurance and improve creeping bentgrass conversion.

A spring application of PROGRASS SC Herbicide following fall applications is useful in further reducing populations of annual bluegrass. Apply PROGRASS SC Herbicide in the spring after creeping bentgrass has resumed active growth and has achieved full green-up.

Do not apply PROGRASS SC Herbicide to renovated or overseeded bentgrass fairway turf less than three (3) to four (4) weeks after seedling emergence.

Do not overseed with creeping bentgrass until three (3) to four (4) weeks after the last PROGRASS SC Herbicide application.

IV. Turf-Type Tall Fescue

Apply PROGRASS SC Herbicide to turf-type tall fescue in the fall. If overseeding with turf-type tall fescue, PROGRASS SC Herbicide application and the seeding may be simultaneous.

Warm Season Turfgrasses

I. St. Augustinegrass (Established Turf)

Apply PROGRASS SC Herbicide to St. Augustinegrass in the spring and early summer to suppress bermudagrass and bermudagrass

seedhead formation or control annual bluegrass. Best results are achieved when multiple applications are made. PROGRASS SC Herbicide suppresses the development of actively growing bermudagrass in established St. Augustine Sod. Begin PROGRASS SC Herbicide applications when bermudagrass first breaks dormancy in the spring. Application timing is critical to achieve optimum results and will vary depending on location and temperature. A second application of PROGRASS SC Herbicide may be made 21-28 days after the first application. For optimum suppression, a third application may be necessary 21-28 days after the second application. Temporary stunting and minor discoloration of St. Augustinegrass may occur after application. To improve suppression, PROGRASS SC Herbicide may be tank-mixed with Atrazine in each application. DO NOT treat St. Augustinegrass that has not been established for six months. DO NOT treat St. Augustinegrass that is under stress or injury may occur. If the St. Augustinegrass shows signs of severe yellowing or stunting, discontinue applications.

II. Overseeded Dormant Bermudagrass

Apply PROGRASS SC Herbicide to overseeded bermudagrass in late fall, one to two weeks after emergence of overseeded perennial ryegrass to control annual bluegrass. Treatment to turf that is not fully dormant may cause early injury and/or delayed spring green-up. Applications of 1.0 to 1.125 quarts (1 to 1.125 lb a.i./A) of PROGRASS SC Herbicide per acre have shown acceptable control of annual bluegrass with minimum injury to non-dormant (or predormant) bermudagrass when applied in late November or early December. Lower rates have shown lesser annual bluegrass control and higher rate have shown more bermudagrass injury when the bermudagrass is not fully dormant. The onset of dormancy will vary according to location and yearly weather. One or two supplemental applications at 21 to 28 day intervals may be required to maintain control through the winter months. DO NOT APPLY more than two applications of PROGRASS SC Herbicide when treating non-dormant bermudagrass. The final PROGRASS SC Herbicide application may be made to bermudagrass no later than February 1st or 4 weeks prior to breaking of winter dormancy. Late applications (within 4 weeks of breaking dormancy) to turf that will be breaking dormancy may temporarily delay resumption of active growth (spring green-up) of bermudagrass in the spring. The breaking of winter dormancy will vary according to location and yearly weather. Application of PROGRASS SC Herbicide to bermudagrass that is stressed due to shade, poorly drained soils, and high traffic may result in increased turf injury. Avoid use of PROGRASS SC Herbicide or use lower rates under these conditions. Care needs to be taken to keep the PROGRASS SC Herbicide in the overseeded areas during the applications (particularly when making applications to non-dormant bermudagrass) so that the ryegrass will mask any early injury or late transition in the spring. Higher seeding rate may also be helpful to minimize a thin stand appearance since annual bluegrass will be removed. Common bermudagrass in fairways or roughs may be more susceptible to herbicide injury than hybrid bermudagrass.

APPLICATION RATES AND TIMINGS ¹								
Turf Type	Main Target Pests	Start	# Apps	Re-treatment Interval (Days)	Use Rate ²		Overseeding Safety Interval ³ (Weeks After Treatment)	Application Safety Interval ⁶ (Weeks After Emergence)
					lb a.i./A	Qt/A (fl oz/ 1,000 sq ft)		
Ryegrass ⁵	Annual Bluegrass	Fall Spring ⁴	2-3 1-2	21 - 28 21 - 28	0.95- 2.04	1.0 - 2.0 (0.7-1.5)	1-2	1-2
Kentucky Bluegrass	Annual Bluegrass	Fall	2 - 3	21 - 28	0.82	0.75 (0.6)	6	8
Creeping Bentgrass	Annual Bluegrass	Fall	2-3	21 - 28	0.82	0.75 (0.6)	4	4
		Spring ⁴	1-2	21 - 28	0.82	0.75 (0.6)		
Turf-Type Tall Fescue	Annual Bluegrass	Fall	2 - 3	21 - 28	0.82 -1.5	0.75 - 1.5 (0.6-1.1)	0	2-3
St. Augustinegrass (Est. Turf)	Bermudagrass Suppression	Spring	2 - 3	21 - 28	1.5-2.3	1.5 - 2.3 (1- 1.7)	N/A	N/A
Seashore Paspalum	Bermudagrass Suppression	Spring	Multiple ⁷	14	0.5	0.5 (0.37)	6	N/A
Overseeded (ryegrass)								
Bermudagrass:								
Dormant Bermudagrass	Annual Bluegrass	Fall	2 - 3	21 - 28	0.82- 2.04	0.75 - 2.0 (0.6-1.5)	N/A	N/A
Non-dormant Bermudagrass	Annual Bluegrass	Fall	2	21 - 28	0.95	1.0 (0.7)	N/A	N/A

¹Refer to label text for further instructions regarding each grass species. Do not apply more than 2.3 qt per acre (=1.7 fl oz/1000 sq ft = 2.3 lb a.i./A) per application. Do not exceed 6 fl oz/1000 sq ft. PROGRASS SC herbicide (equivalent to 8.2 lb a.i./A) per year.

²Apply specified rate of PROGRASS SC Herbicide in 20 to 60 gallons of water per acre or 1/2 to 1.5 gallons of water per 1,000 sq ft.

³Interval applies to overseeding specific grass on same type of grass after last PROGRASS SC Herbicide treatment.

⁴Use spring application after previous fall treatments only.

⁵Do not exceed 4 applications per year using the 1.5 fl oz/1000 sq ft. (2.04 lb a.i./A) rate.

⁶Refers to interval in which PROGRASS SC Herbicide may be safely applied to specific grass following seed emergence.

⁷Multiple applications not to exceed 16 applications per year, using the 0.37 fl oz product/1,000 sq ft. (0.5 lb a.i./A) rate. Do not apply to turf when foliage is wet or dew is present. Multiple applications are necessary for maximum suppression.

N/A – Not applicable

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Protect Prograss SC Herbicide from freezing temperatures.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Chipco and Prograss are registered trademarks of Bayer.
Backed By Bayer is a trademark of Bayer.

NET CONTENTS: 2.5 gallon

Produced for



Bayer Environmental Science

**A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709**

PROGRASS SC HERBICIDE (Pending) 11/17/2017, 04/11/2018, 05/02/2018, 05/11/2018, 5/14/2018, 5/31/2018, 6/08/2018, 6/11/2018