

## UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOYIC SUBSTANCES

FEB 2 1 2009

Mr. Richard M. Gorrel Bayer Environmental Science 2 T. W. Alexander Drive, Research Triangle, NC 27709

Subject: Label Notification(s) for Pesticide Registration Notice 2007-4

Dear Registrant:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated January 30, 2009 for:

#### EPA Registration 432-1476 Tiberon 2.8% SC (Plant Growth Regulator)

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

Please be reminded that 40 CFR Part 156.140(a)(4) requires that a batch code, lot number, or other code identifying the batch of the pesticide distributed and sold be placed on nonrefillable containers. The code may appear either on the label (and can be added by non-notification/PR Notice 98-10) or durably marked on the container itself.

If you have any questions, please contact me directly at 703-305-6249 or Banza Djapao of my staff at 703-305-7269.

Sincerely,

Linda Arrington

Notifications & Minor Formulations Team Leader Registration Division (7505P) Office of Pesticide Programs

Please read instructions on n	everse before complet	ting form.	ــــــــــــــــــــــــــــــــــــــ	Form Approved	d. OMB No. 2	070-0060	Approval expires 2-28-95
<b>⊗EPA</b>	Environmental	nited States Protectiongton, DC 204	• .	<b>✓</b>	Registra Amenda Other		OPP Identifier Number
		Application	n for Pesticio	le - Section	l		
1. Company/Product Number 432-1476			1	Product Manager ny Kish		3. Pro	posed Classification
4. Company/Product (Name) Tiberon 2.8% SC (plant	growth regulator)		PM# 22	::			
5. Name and Address of App	licant (Include ZIP Co	de)	6. Expe	edited Reveiw	. In accorda	nce with	FIFRA Section 3(c)(3)
Bayer Environmental 2 T. W. Alexander Dri Research Triangle Pa	ve		to:	ny product is sin	NO.	TUFICA	
	is a new address	,	Produ	ct Name	FE	B 212	2009
			Section - I				
Amendment - Explain Resubmission in resp	onse to Agency letter	dated		Final printed labe Agency letter da "Me Too" Applic Other - Explain b	ited eation.	to	
Explanation: Use addition This notification is consistent labeling or the confidential sta EPA. I further understand the FIFRA and I may be subject Notification of adding a revise	with the provisions of Fatement of formula of the at if this notification is not one action a	PR Notice 98-10 his product. I used consistent wand penalties un	and EPA regulation nderstand that it is a ith the terms of PR N nder sections 12 and	violation of 18 U.S lotice 98-10 and 4	S.C. Sec. 1001	o willfully i	make any false statement to
<del> </del>		· · · · · · · · · · · · · · · · · · ·	Section - I	11 :			
1. Material This Product Will	Be Packaged In:		<del></del>	· · · · · · · · · · · · · · · · · · ·			
Child-Resistant Packaging	Unit Packaging		Water Soluble P	ackaging	2. Type of	Container	· · · · · · · · · · · · · · · · · · ·
Yes	Yes		Yes			Metal	
√ No	✓ No		√ No			Plastic Glass	
Certification must be submitted	If "Yes" Unit Packaging wgt	No. per container	if "Yes" Package wgt	No. per container		Paper Other (S	Specify)
3. Location of Net Contents		4. Size(s) Re	tail Container	5. L	ocation of Lab	el Directio	ons
6. Manner in Which Label is	onteiner Affixed to Product	Lithog Peper Stenc	<del></del>	Other			
		Stenc	Section - I				
1. Contact Point (Complete	items directly below	for identification	r <del> </del>	a contacted, if ne	cessary, to pr	·····	
Name Richard M. Gorrell			Title Manager, Regist	rations		(919) 549	e No. (Include Area Code) 3-2423
I certify that the state I acknowledge that an both under applicable	y knowlingtly false or		all attachments th				6. Date Application Received (Stamped)
2. Signature	1 m. Horale		3. Title Manager, Registra	tions			
4. Typed Name Richard M. Gorrell			5. Date				
I		1	Jan	uary 30, 200	Ja		1

## **Bayer Environmental Science**



Bayer Environmental

RTP, NC 27709 Phone: 919.549.2000

2 T. W. Alexander Drive

Science

January 30, 2009

EPA09RMG0130TiberonSCNotifStorDisp Fedex

Ms. Linda Arrinaton U.S. Environmental Protection Agency Office of Pesticide Programs (7504P) Document Processing Desk (NOTIF) Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

RE:

Tiberon 2.8% SC plant growth regulator (EPA Reg. No. 432-1476): Notification to Add the Revised Storage Disposal Statement

Dear Ms. Arrington:

Bayer Environmental Science (BES) is submitting Tiberon 2.8% SC revised label to add the EPA mandated storage disposal statement. The changes are the same as the EPA example and therefore the label is being submitted by notification.

The following information is contained in this submission:

- 1. EPA Forms 8570-1
- 2. 4 copies of the label (one on electronic disc)

If you have any questions, contact me at (919) 549-2423 or email me at mike.gorrell@bayercropscience.com.

Sincerely,

Richard M. Gorrell

Manager, Registrations

incl.

Cc: Karen Shearer Scott Welge Don Myers

> A Business Group of **Bayer Cropscience**

#### NOTIFICATION

FEB 2 1 2009

## Tiberon 2.8% SC

FOR USE IN OUTDOOR NURSERY AND PLANT PRODUCTION FACILITIES	
DO NOT USE IN A GREENHOUSE	
ACTIVE INGREDIENT: Cyclanilide (1-[[(2,4-Dichlorophenyl)amino]carbonyl]cyclopropa	necarboxylic acid)2.8%
OTHER INGREDIENTS:	
TOTAL:	
*Contains 0.24 pounds cyclanilide per gallon.	·
EPA Reg. No. 432-1476	EPA Est. No.

# WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-334-9745

#### **FIRST AID**

IF IN EYES:	Hold eye open and rinse slowly and gently with water for 15-20 minutes.			
	Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.			
	Call a poison control center or doctor for treatment advice.			
IF SWALLOWED:	Call a poison control center or doctor immediately for treatment advice.			
	Have person sip a glass of water if able to swallow.			
	Do not induce vomiting unless told to do so by a poison control center or doctor.			
	Do not give anything by mouth to an unconscious person.			
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.			
	Rinse skin immediately with plenty of water for 15-20 minutes.			
	Call a poison control center or doctor for treatment advice.			

### PRECAUTIONARY STATEMENTS

## WARNING HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Harmful if swallowed. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear protective eyewear, long-sleeved shirt, long pants, shoes plus socks, and chemical resistant gloves made of a waterproof material.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product concentrate. Do not reuse them.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **User Safety Recommendations**

Users must wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Users must remove clothing immediately if pesticide gets inside. Then wash body thoroughly and change into clean clothing. The contaminated clothing must be washed before reuse.

Users must remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

#### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations. Read entire label before using this product.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

#### **PESTICIDE STORAGE**

Store in original container and keep tightly closed. Store in a cool dry place.

#### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

#### **CONTAINER DISPOSAL**

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses; and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry-interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls; chemical-resistant gloves made of a waterproof material; shoes plus socks; protective eyewear.

#### **GENERAL PRECAUTIONS AND RESTRICTIONS**

For local recommendations on rates, spray volumes (gallons of water per acre), and spray equipment under varying temperature and rainfall conditions consult Bayer Environmental Science Representative for his experience with this product in your area.

#### Use Directions for Tiberon Plant Growth Regulator General Information

Tiberon is a plant growth regulator that inhibits auxin activity, particularly in meristematic plant tissue. As a result, apical dominance is suppressed. This effect has resulted in increased branching of trees without the need to prune the central leader. Increased branching in shrubs and cut-flowers can also be achieved. Plants treated with Tiberon are often more compact than untreated plants.

Branching is typically induced near actively growing terminals. In some plants, such as California pepper tree and Chinese flame tree (*Koelreuteria bipinnata*), buds also break in blank wood below the terminal shoot apex. Effects on auxin activity are temporary, usually about 3 weeks. Additional treatments may be made to continue the effect. Treatment intervals for each specific use need to be evaluated on a case-by-case basis.

This product may replace the need to prune off the terminal growth to induce branching or can be used in conjunction with pruning or pinching.

Prior to the production-scale use of this product small-scale on-site testing should be implemented for the specific site and variety to determine appropriate rate to provide desirable performance. Tiberon may be used in a program approach with other plant growth regulators, but it is recommended that all uses in a program with other plant growth regulators be evaluated on a case-by-case basis.

#### **Application**

Tiberon is applied as a broadcast spray. Spray foliage to drip. Entire plants or specific plant parts may be treated. Use of a non ionic adjuvant may promote increased activity. Plants must be vigorously growing to achieve the branching response. This product is acidic. Application of Tiberon on leaves treated with copper can result in leaf burn.

Tiberon can be used in nursery sites on any non-bearing plant. However, testing on a small scale should be conducted on-site to determine performance and impact on plant quality prior to use on a large scale. Consider testing various rates, timing, and one vs. two applications for optimal response at each location. Results may vary from year to year.

#### Use on Ornamentals and Cut Flowers

The following are general guidelines. Plants must be in an active and vigorous growing condition at the time of application and for several weeks following the application. Any ornamental plant that has shown a desirable response to Tiberon may be considered as a candidate for application of this product. Consult the rate chart below for the correct amount of Tiberon per gallon or liter of water. Most woody ornamentals respond well to 1 to 2 applications of 50 to 200 ppm. Typically one application of 100 ppm is adequate. Many herbaceous ornamentals respond to applications between 5 and 20 ppm.

For Lilacs use 1-3 applications of 100-200 ppm one month apart to achieve the desired amount of branching.

Use 50 to 75 ppm rate range for Goldenrod (Solidago).

Some species respond to 5 to 20 ppm, including Bradford pears, and Myrtle (Myrtus cummunis).

Ornamentals that have shown a favorable response (increased branching) to Tiberon:

Common Name	Scientific Name
Abelia	Abelia gaucherie
Anise	Illicium parviflorum
Arizona redwood	Vuaquelinia californica
Azalea	Rhododendron sp
Barberry	Berberis thunbergii
Birch	Betula spp
Blueberry (non bearing ornamental)	Vaccinium spp
Bottle brush	Callistemon lanceolatus
California pepper tree	Schinus molle
Camellia	Camellia spp
Carissa	Carrisa macrocarpa tuttlei
Cherry	Prunus spp
Chinese flame tree	Koelreuteria bipinnata
Convolvulus	Convolvulus spp
Coneflower, Eastern purple	Coneflower, Eastern purple
Coreopsis	Coreopsis rosea
Crabapple ,	Malus spp:
Crape myrtle	Lagerstroemia sp
Croton	Codiaeum spp
Eastern redbud	Cercis canadensis
Euonymus	Euonymus alata
Ficus	Ficus nitida
Flowering quince	Chaenomeles spp
Gardenia	Gardenia jasminoides 'veitchii'
Common hawthorn	Crataegus spp
Hibiscus (tropical)	Hibiscus spp
Honey locust	Glidetsia triacanthos
Hydrangea	Hydrangea spp
Indian hawthorn	Raphiolepsis indica
Holly/inkberry	llex glabra

Common Name	Scientific Name		
Holly	Ilex crenata 'Sky Pencil'		
Holly/Burford	llex cornuta 'Bufordii'		
Japanese ternstroemia	Ternstroemia gymnanthera		
Little-leaf linden	Tilia cordata		
Loquat	Eriobotrya japonica		
Myrtle	Myrtus spp		
Oak	Quercus spp		
Oleander	Nerium oleander		
Phlox	Phlox paniculata		
Physocarpa	Asclepias physocarpa		
Privet	Ligustrum spp		
Purple coneflower	Echinacea purpurea		
Rose	Rosa spp		
Spirea	Spirea spp		
Tree lilac	Syringa reticulata		
Viburnum	Viburnum tinus		
Xylosma	Xylosma senticosum		

Ornamentals that have not shown a favorable response to Tiberon:

Common Name	Scientific Name
Acacia	Acacia redolens
Acacia	Acacia spp
Blue Girl holly	llex spp
Boronia	Boronia spp
Butterfly bush	Buddleia spp
Cape honeysuckle	Tecomaria capensis
Cistus	Cistus corbariensis
Common buckthorn	Rhamnus spp
Desert pine	Pinus eldarica
Dieffenbachia/dumb cane	Dieffenbachia seguine
Dwarf schefflera	Schefflera aboricola
Heller Japanese holly	llex crenata 'Helleri'
Hosta	Hosta spp
Hydrangea	Hydrangea macrophylla 'Geoffrey
	Chadbundt'
Impatiens	Impatiens capensis
Jacaranda	Jacaranda acutifolia
Japanese pieris	Pieris japonica 'Cavatine'
Juniper	Juniperus horizontalis 'Blue Rug'
Maple	Acer spp
Mock orange	Pittosporum tobira
Nandina/Heavenly bamboo	Nandina domestica
Olive	Olea europea
Photinia	Photina spp
Rhododendron	Rhododendron 'Golden Touch'
Schefflera	Schefflera spp
Star jasmine/confederate jasmine	Trachelospermum jasminoides
Strawberry tree	Arbutus unedo
Rosemary	Rosemarinus officinalis
Russian sage	Perovskia atriplicifolia
Variegated privet	Ligustrum sinensis 'variegata'
Viburnum	Viburnum nudum
Winter daphne	Daphne odora
Wisteria	Wisteria spp

#### Use on fruit trees (non-food only)

Sweet cherry and Apple varieties: 1 to 2 applications of 50 to 100 ppm. Typically one application at 100 ppm provides acceptable branching. If using a two-spray regime, consider making the second application about 1 week later. Some reduction in growth of the central leader may be observed with two applications.

Pear and Plum varieties: 1 application of 5 to 20 ppm. Typically 10 to 20 ppm provides acceptable branching.

Treatments to nectarine have not been effective.

#### Application timing for fruit trees (non-food only)

The following information is for initial guidelines for application timing. It is important to have a vigorously growing plant at the time of application and for several weeks following application to obtain satisfactory branching and good development of new branches.

Induced branching of cherries tends to occur 4 to 8 inches above the height of the terminal shoot apex at the time of application. Induced branching in apples occurs about 1 to 4 inches below the height of the terminal shoot apex at the time of application.

Tiberon Rate Chart					
Cyclanilide	Amount Tiberon per volume of water				
ppm final concentration	ounces per gallon water	ounces per 100 gallons water	mL per liter water	mL per 100 liter water	
1	0.0045	0.45	0.035	3.5	
5	0.0220	2.20	0.180	18	
10	0.0450	4.50	0.350	. 35	
20	0.0900	9.00	0.700	70	
50	0.2200	22.00	1.800	180	
100	0.4500	45.00	3.500	350	
200	0.9000	90.00	7.000	700	
500	2.2400	224.00	18.000	1800	
1000	4.5000	450.00	35.000	3500	
	l .	1	,		

Do not tank mix Tiberon SC with pesticides or liquid fertilizers containing micronutrients. Use Restrictions

- Do not apply more than 0.34 pound active ingredient per acre per crop per year
- Do not apply more than 3 applications per crop per year

#### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury or ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTANT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTANT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES, RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTANT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Net Contents:

NOTIFICATION

FEB 2 1 2009

PRODUCED FOR



## Bayer Environmental Science

A BUSINESS GROUP OF BAYER CROPSCIENCE LP P O BOX 12014 2 T. W. Alexander Drive Research Triangle Park, NC

Tiberon 2.8% SC (MASTER) 10/11/07, Notification 01/28/09