

432-1487

8/18/2010

111



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Mr. Richard M. Gorrell  
Product Registration  
Bayer Environmental Science  
2T.W Alexander Drive  
Research Triangle Park, NC 27709

AUG 18 2010

SUBJECT: Application for Pesticide Notification (PRN 98-10)  
Request General Label Change/Add Fairy Ring Disease  
EPA Reg. No. 432-1487  
Application Dated July 9, 2010

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 07/09/10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-5335 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

Paul J. Mastradone, Ph.D., Acting  
Notifications & Minor Formulations Team Leader  
Registration Division (7505P)  
Office of Pesticide Programs



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 432-1487	2. EPA Product Manager Ms. Shaja Joyner	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Triticonazole 3SC Fungicide (ABN Chipco Triton FLO Fung.)	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Bayer Environmental Science 2 T. W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**NOTIFICATION**

**AUG 18 2010**

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

This notification is consistent with the provisions of PR Notices 98-10 and 2007-4 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.  
- Fairy Ring Added

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	
* Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
				<input type="checkbox"/> Plastic	
				<input type="checkbox"/> Glass	
				<input type="checkbox"/> Paper	
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 4x1 gallon		5. Location of Label Directions <input checked="" type="checkbox"/>	
6. Manner in Which Label is Affixed to Product		<input checked="" type="checkbox"/> Lithograph Paper glued Stenciled		<input type="checkbox"/> Other _____	

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Richard M. Gorrell	Title Manager, Registrations	Telephone No. (include Area Code) (919) 549-2923
----------------------------	---------------------------------	---

#### Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, or both under applicable law.

6. Date Application Received  
(Stamped)

2. Signature <i>Richard M. Gorrell</i>	3. Title Manager, Registrations
4. Typed Name Richard M. Gorrell	5. Date July 9, 2010



# Bayer Environmental Science

July 9, 2010

EPA10RMG0709Triticonazole3SCFungNotifFairyRing  
Fedex

Ms. Linda Arrington  
U.S. Environmental Protection Agency  
Office of Pesticide Programs (7504P)  
Document Processing Desk  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202-4501

RE: Triticonazole 3SC Fungicide (432-1487) (ABN Chipco Triton FLO Fungicide)  
Label Notification: Fairy Ring Disease Control Added

Dear Ms. Arrington:

Bayer Environmental Science (BES) is submitting Triticonazole 3SC Fungicide notifying the Agency of the addition of control of fairy ring disease to the label.

The following information is enclosed with this letter:

1. 8570-1 Application Form
2. 5 copies of the revised label (one copy highlighted in yellow)

Bayer Environmental Science  
2 T.W. Alexander Drive  
RTP, NC 27709  
Phone : 919.549.2000


If you have any questions, contact me at (919) 549-2423 or email me at [mike.gorrell@bayercropscience.com](mailto:mike.gorrell@bayercropscience.com).

Sincerely,

Richard M. Gorrell  
Manager, Registrations

Incl.

Cc: Karen Shearer  
Jimmy Johnson  
Richard Rees



A Business Group of  
Bayer CropScience

# Triticonazole 3 SC Fungicide

*For control of diseases on turfgrasses of commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas, cemeteries and sports fields.*

**ACTIVE INGREDIENT:**

Triticonazole: ..... 30.1%

**OTHER INGREDIENTS:** ..... 69.9%

\* Contains 3 lbs triticonazole per US gallon

**NOTIFICATION**

AUG 18 2010      Total ..... 100.0%

**EPA Reg. No. 432-1487**

**EPA Est. No.**

## KEEP OUT OF REACH OF CHILDREN CAUTION

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call (1-800-331-2867)

**FIRST AID**

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<p>In case of emergency call toll free the Bayer CropScience Emergency Response Telephone No. 1-800-334-7577. Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	
<p><b>NOTE TO PHYSICIAN:</b> Treat Symptomatically.</p>	

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.


**Personal Protective Equipment (PPE):**

Applicators and other handlers must wear long sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves made of any waterproof material such as natural rubber  $\geq$  14 mils (if more options are needed, follow the instructions for category A on an EPA chemical-resistance category selection chart).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements:**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.



### User Safety Recommendations

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove and wash contaminated clothing before reuse.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

### DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of any of any waterproof material such as natural rubber ≥ 14 mils.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when the product is used to produce agricultural plants on farms, nurseries or greenhouses.

DO NOT enter or allow others to enter treated areas until sprays have dried.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross contamination with other pesticides.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture, or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Disposal:** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple-rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. After triple rinsing procedure dispose of container.

## GENERAL INFORMATION

TRITICONAZOLE 3 SC FUNGICIDE is a broad spectrum fungicide with preventive, systemic, residual and curative properties recommended for the control or suppression of certain turf diseases. Disease control is optimal when TRITICONAZOLE 3 SC FUNGICIDE is applied in a preventive and scheduled spray program and used in rotation with other fungicides.

Failure to follow these label instructions may result in turfgrass injury and inferior turfgrass quality. This product may cause staining. Avoid off-target sites such as sidewalks, patios, driveways pavers or similar materials. Avoid entry to areas used for recreation and sports until spray deposits are completely dry after application to prevent staining of clothing and shoes.

TRITICONAZOLE 3 SC FUNGICIDE can be used on all commercial, institutional and residential turfgrass, such as golf courses, sod farms, commercial lawns, rights of way, parks, recreation areas, cemeteries and sports fields. Not for use by homeowners.

## APPLICATION INFORMATION

TRITICONAZOLE 3 SC FUNGICIDE may be used alone to control important diseases of turfgrass or alternatively used in tank mix or sequentially with other registered turfgrass fungicides.

Applications using sufficient water volume to provide thorough and uniform coverage generally provide the most effective disease control. Do not make applications when conditions favor drift beyond the target application area. All applications of TRITICONAZOLE 3 SC FUNGICIDE must be made in accordance with the use directions of this label.

TRITICONAZOLE 3 SC FUNGICIDE, when used in conjunction with best turfgrass management practice, will control the diseases listed under **TURF-SPECIFIC USE DIRECTIONS**.

TRITICONAZOLE 3 SC FUNGICIDE, when combined with other products such as **Chipco® Signature™**, will relieve the symptoms of summer stress syndrome/decline.

### Turfgrass Species:

- Colonial and Creeping Bentgrass
- Common and Hybrid Bermudagrass
- Annual, Rough and Kentucky Bluegrass
- Buffalograss
- Centipedegrass
- Dichondra
- Fine and Tall Fescue
- Seashore Paspalum
- Annual and Perennial Ryegrass
- St. Augustinegrass
- Zoysiagrass
- Kikuyugrass

**DO NOT** use on ultra-dwarf bermudagrass varieties.

**AVOID** the shorter application intervals if the turf is under excessive stress in the summer or under an intensive growth control program.

### Ground Application

For ground application equipment use the appropriate product quantities described in this label in 1 to 5 gallons of water per 1,000 sq ft for turf (44 to 220 gallons per acre).

## SPRAY MIXING AND COMPATIBILITY

In general, begin with clean spray equipment and add one-half of the required quantity of water to the spray or mixing tank and start agitation. Refer to the product amounts tabulated under **PRODUCT QUANTITY (FL OZ) by SPRAY VOLUME and TANK CAPACITY** and add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank-mixing with other products. In general, follow the order beginning first with water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates and emulsifiable concentrates last. Maintain agitation throughout spraying. Do not allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

The turf safety of all potential tank-mixes with TRITICONAZOLE 3 SC FUNGICIDE including additives and other pesticides has not been tested on all turf type variants. Before applying any tank-mixture not specifically recommended on this label, safety to the target turf should be confirmed. TRITICONAZOLE 3 SC FUNGICIDE is compatible with most commonly used fungicide, herbicide, insecticide, growth regulator and foliar nutrient products. However, the physical compatibility of TRITICONAZOLE 3 SC FUNGICIDE with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

- The use of a soil penetrating adjuvant may improve the movement into the soil when irrigation after treatment is required for disease control.
- Combinations of TRITICONAZOLE 3 SC FUNGICIDE at high labeled rates with PGRs may impact turf quality and reduce turf growth particularly during period of heat stress and high humidity.
- When preparing spray mixtures of TRITICONAZOLE 3 SC FUNGICIDE with Bayer Products add the products in the following order: 1) Water dispersible products , 2) TRITICONAZOLE 3 SC FUNGICIDE followed by other suspension concentrates such as Compass™, 26GT™, and lastly 3) Chipco® Signature™.


**RESISTANCE MANAGEMENT**

The active ingredient in TRITICONAZOLE 3 SC FUNGICIDE belongs to the Sterol Inhibitors or Demethylation Inhibitors class of chemistry (Group 3). To maintain long term effectiveness of this fungicide, follow the specific resistance management guidance from a local extension specialist.

To limit the potential for development of disease resistance to this fungicide class do not make more than 3 sequential applications of TRITICONAZOLE 3 SC FUNGICIDE or any Group 3 containing fungicide before rotating with a fungicide from a different group.

**GENERAL RESTRICTIONS AND LIMITATIONS**

- DO NOT** apply more than the maximum seasonal rate from any combination of products containing triticonazole.
- DO NOT** apply more than 3.3 fl oz of TRITICONAZOLE 3 SC FUNGICIDE per 1,000 sq ft (3.451 lb ai/A) per year.
- Follow the **TURF-SPECIFIC USE DIRECTIONS** for repeat application intervals.
- For golf courses only: DO NOT** apply to turf cut higher than 1 inch on golf holes where water bodies are present.
- This product cannot be used to formulate or reformulate any other pesticide product.**
- DO NOT** apply this product to crops other than turfgrass.
- DO NOT** apply to areas used for grazing.
- DO NOT** feed clippings to livestock or poultry.
- DO NOT** apply through irrigation equipment of any type.
- DO NOT** use on ultra-dwarf bermudagrass varieties.



## TURF-SPECIFIC USE DIRECTIONS

TRITICONAZOLE 3 SC FUNGICIDE controls the diseases listed below.

Disease Control	Rate fl oz Product /1,000 sq ft	Use Information
Anthracnose ( <i>Colletotrichum cereale</i> )	0.41-1.1	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.
Brown Patch ( <i>Rhizoctonia solani</i> )	0.41-1.1	Begin fungicide applications preventively when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.
Brown Patch, Cool weather/Yellow patch ( <i>Rhizoctonia cerealis</i> )	0.55 -1.1	Make 1 to 2 applications in the fall using a 21-28 day interval or when conditions are favorable for disease development.
Brown Ring (Waitea) Patch ( <i>Waitea circinata</i> var. <i>circinata</i> )	0.55 -1.1	Begin fungicide applications at the early stage of yellow ring development when conditions are favorable for disease development. Continue as needed, on a 14-28 day interval. Late curative applications will not be effective. <b>Irrigate to move the fungicide beneath the thatch.</b>
Dollar Spot ( <i>Sclerotinia homeocarpa</i> )	0.28 -1.1	Begin fungicide applications preventively. Continue as needed, on a 14-28 day interval. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval. Rotation with 26GT™ fungicide is suggested for resistance management.
Fairy Ring	0.55 - 1.1	The first application should be made in the spring when 5-day average daily soil temperatures taken at 2 inches are 55 – 65°F. In sandy soils, average air temperature can also be an adequate measure.  The second application should be made 28 days later, and applications should be made in 2 gallons H <sub>2</sub> O/1000 ft <sup>2</sup> . In addition, applications should be watered in with ½ to ¼ inch of irrigation. If possible, irrigation should be applied immediately. At the latest, irrigation should occur that night. TRITICONAZOLE 3 SC FUNGICIDE applications that are not watered in have resulted in unsatisfactory control.
Fusarium Patch ( <i>Microdochium nivale</i> )	0.28 -1.1	Begin fungicide applications preventively in the spring and early summer when the turf stays moist and temperatures are in the range 32-65° F. Repeat applications at 10-14 day intervals when heavy disease pressure is anticipated. <b>Under conditions for severe disease</b> use the higher rate and tank mix with 26GT fungicide.
Gray Snow Mold, Typhula Blight ( <i>Typhula</i> spp.)	0.28 -1.1	Begin fungicide applications preventively in the late fall. Make 1-2 applications, repeating at 14-28 day intervals when heavy disease pressure is anticipated. <b>Under conditions for severe disease</b> use the higher rate and tank mix with other fungicides with alternative modes of action that are effective on snow molds.
Necrotic Ring Spot ( <i>Ophiosphaerella korrae</i> )	0.28 -0.55	Begin fungicide applications preventively when soil temperatures reach 60° F early spring or summer and continue at 28 day intervals when conditions are favorable for development. <b>Irrigate or aerate to allow movement of fungicide into the root zone.</b>



Disease Control	Rate fl oz Product /1,000 sq ft	Use Information
Pink Patch ( <i>Limonomyces roseipellis</i> )	0.55 -1.1	Begin fungicide applications preventively when soil temperatures reach 60° F to 70° F under periods of high rainfall when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval.
Red Thread ( <i>Laetisaria fuciformis</i> )	0.41 -1.1	Begin fungicide applications preventively when night air temperatures reach 60° F to 70° F under periods of high rainfall when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated. <b>Under conditions for severe disease</b> use the higher rate and shorter interval. Continue as needed, on a 14-28 day interval.
Rust ( <i>Puccinia</i> spp.)	0.28 -0.55	Begin fungicide applications preventively when conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions for severe disease or for an early curative application</b> , use the higher rate and shorter interval. Continue as needed, on a 14-28 day interval.
Summer Patch ( <i>Magnaporthe poae</i> )	0.55 -1.1	Begin fungicide applications preventively when soil temperatures reach 60° F to 65° F at a 2-inch soil depth unless otherwise advised by local recommendations. Continue as needed, on a 14-28 day interval. Use the higher rate and shorter interval under severe disease pressure or if there is prior history.
Take-all Patch ( <i>Gaeumannomyces graminis</i> var. <i>avenae</i> )	0.55 -1.1	Begin fungicide applications preventively in the fall. Make 1-2 applications, repeating at 14-28 day intervals when local disease conditions are favorable for development. <b>Under preventive applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions or prior history for severe disease or for an early curative application</b> , use the higher rate and shorter interval.
Large Patch (aka Rhizoctonia Large Patch Zoysia Patch) ( <i>Rhizoctonia solani</i> )	0.55 -1.1	Begin fungicide applications preventatively in the fall. Make 1-2 applications, repeating at 14-28 day intervals when conditions are favorable for development. <b>Under preventative applications</b> where light disease pressure is anticipated, use the lower rate and longer interval. <b>Under conditions or prior history for severe disease or for an early curative application</b> , use the higher rate and shorter interval.

Additional Turfgrass Uses	Rate oz/1,000 sq ft	Use Information
Algae	0.55 -1.1	Use preventively the lower rate for the suppression of algae. Continue as needed, on a 14-28 day interval. <b>Tank mix with chlorothalonil or mancozeb to improve the curative control of algae.</b>
Summer stress syndrome / summer decline	0.55 -1.1	Begin applications prior to the onset of conditions for abiotic stress to reduce symptoms of summer stress/decline. Continue as needed, on a 14-28 day interval. <b>Tank mix Chipco® Signature™ to improve the relief in symptoms on turf under moderate to severe stress.</b>

<sup>1</sup>DO NOT apply more than 3.3 fl oz per 1,000 sq ft of TRITICONAZOLE 3 SC FUNGICIDE per year.

**PRODUCT QUANTITY (FL OZ) by SPRAY VOLUME and TANK CAPACITY**

TRITICONAZOLE 3 SC FUNGICIDE at 0.55 fl oz per 1,000 sq ft

		SPRAY VOLUME (gallons per 1,000 sq ft)				
		1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
<b>Spray Tank Capacity</b>	<b>25 Gallons</b>	14 fl oz	7 fl oz	4.7 fl oz	3.5 fl oz	2.8 fl oz
	<b>50 Gallons</b>	28 fl oz	14 fl oz	9.3 fl oz	7 fl oz	5.6 fl oz
	<b>100 Gallons</b>	56 fl oz	28 fl oz	18.7 fl oz	14 fl oz	11.2 fl oz
	<b>200 Gallons</b>	112 fl oz	56 fl oz	37.3 fl oz	28 fl oz	22.4 fl oz

TRITICONAZOLE 3 SC FUNGICIDE at 1.1 fl oz per 1,000 sq ft

		SPRAY VOLUME (gallons per 1,000 sq ft)				
		1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
<b>Spray Tank Capacity</b>	<b>25 Gallons</b>	28 fl oz	14 fl oz	9.3 fl oz	7 fl oz	5.6 fl oz
	<b>50 Gallons</b>	56 fl oz	28 fl oz	18.7 fl oz	14 fl oz	11.2 fl oz
	<b>100 Gallons</b>	112 fl oz	56 fl oz	37.3 fl oz	28 fl oz	22.4 fl oz
	<b>200 Gallons</b>	224 fl oz	112 fl oz	74.6 fl oz	56 fl oz	44.8 fl oz

11/11

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

Chipco is a registered trademark of Bayer.  
Signature is a registered trademark of Loveland Inc.  
26GT is a trademark of Bayer.

**NET CONTENTS:**

Produced for



**Bayer Environmental Science**

**A Division of Bayer CropScience LP  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709**

TRITICONAZOLE 3 SC FUNGICIDE (PENDING) 07/09/10