



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

July 13, 2023

Joshua Wood  
Registration Manager  
Bayer CropScience LP  
800 N. Lindbergh Blvd.  
St. Louis, MO 63167

Subject: Registration Review Label Mitigation for TRIFLOXYSTROBIN  
Product Name: CHIPCO 26019T FUNGICIDE  
EPA Registration Number: 432-1505  
Application Date: June 30, 2022  
Decision Number: 592607

Dear Joshua Wood:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Trifloxystrobin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2  
EPA Reg. No. 432-1505  
Decision No. 592607

If you have any questions about this letter, please contact DeMariah Koger by phone at (202)-566-2288, or via email at [koger.demariah@epa.gov](mailto:koger.demariah@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read "Linda Arrington".

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

Enclosure

IPRODIONE	GROUP	2	FUNGICIDE
TRIFLOXYSTROBIN	GROUP	11	FUNGICIDE

# CHIPCO® 26019T Fungicide

[ABN: INTERFACE® FUNGICIDE; INTERFACE StressGard]

- Editorial Note – Marketing claim positioned here

**A Fungicide for the enhancement of greener and more dense turfgrass and for the protection against winter stress and for the prevention and control of certain diseases of turfgrass.**

**Intended for use by professional applicators.**

Editorial Note – [Bracketed text] is optional

**ACTIVE INGREDIENT:**

Iprodione: 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidinecarboxamide).....**23.10%\***  
 Trifloxystrobin (CAS No. 141517-21-7)..... **1.44%\***

**OTHER INGREDIENTS:** ..... **75.46%**

**TOTAL:** ..... **100.00%**

\*Equivalent to 2.14 pounds Iprodione and 0.13 pounds Trifloxystrobin per gallon.

[Shake well before use]

**EPA Reg. No. 432-1505**

**EPA Est. No. 264-MO-02**

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

See [Back][Side] Panel for First Aid Instructions and [Leaflet][Booklet] for Complete Precautionary Statements and Directions for Use. (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

### FIRST AID

**If in eyes:**


- Hold eyes open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

**If swallowed:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of medical emergency for additional information call toll free 1-800-334-7577.

**Note to Physician:** If ingested, induce emesis or lavage stomach. Treat symptomatically.



# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Causes moderate eye irritation. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

### PERSONAL PROTECTIVE EQUIPMENT

#### Mixers, loaders, applicators and other handlers must wear:

Long-sleeved shirt and long pants,

Chemical-resistant shoes and socks,

and chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, Viton®  $\geq$  14 mils.

In addition, mixers and loaders must wear a chemical-resistant apron. See engineering controls for additional requirements.

#### For high-pressure hand wand and backpack sprayer application, mixers, loaders and applicators must wear:

Double-layer clothes,

Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, Viton®  $\geq$  14 mils,

Socks and chemical resistant shoes, and

A minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters

#### For other handling activities and in case of a spill or other emergency exposure, handlers must wear:

Coveralls over long-sleeved shirt and long pants,

Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, Viton®  $\geq$  14 mils,

Chemical-resistant footwear, and

Chemical-resistant apron when cleaning equipment.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate. Applying this product when rain is not predicted for the next 24 hours will help reduce potential risk to aquatic invertebrates by reducing pesticide runoff from the treatment area into water bodies.

### Surface Water Advisory:

Iprodione can contaminate surface water through ground spray applications. Under some conditions, it may also have a high potential for runoff into surface water after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

### Ground Water Advisory:

Several trifloxystrobin degradates have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

## DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water:

- Chemical-resistant apron worn over coveralls over long-sleeved shirt and long pants
- Chemical-resistant boots
- Chemical-resistant gloves made of barrier laminate, butyl rubber  $\geq$  14 mils, nitrile rubber  $\geq$  14 mils, neoprene rubber  $\geq$  14 mils, polyvinyl chloride (PVC)  $\geq$  14 mils, Viton®  $\geq$  14 mils

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to turfgrass uses (golf courses, landscape and institutional areas) of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter or allow others to enter the treated area until sprays have dried.

## PRODUCT INFORMATION

In order to assure maximum crop tolerance and disease control, follow directions on this label and all the precautions and limitations of the package label.

## PRODUCT USE RESTRICTIONS

Use of this product at residential sites is prohibited.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, reservoir, river, permanent stream, marsh or natural pond, estuary, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1 inch on golf holes where water bodies are present.

Do not apply this product when the wind direction is toward aquatic areas.

Do not apply by aerial application.

Do not apply by chemigation.

### MANDATORY SPRAY DRIFT MANAGEMENT

#### Ground Boom Applications:

- Apply with the nozzle height recommended by the manufacturer, but no more than 4 feet above the ground or plant canopy.
- Applicators are required to use a medium or coarser droplet size (ASABE S572.1).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

## IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

## BOOM HEIGHT

For ground equipment, the boom should remain level with the target plants and have minimal bounce.

## RELEASE HEIGHT

Higher release heights increase the potential for spray drift.

## SHIELDED SPRAYERS

Shielding the boom or individual nozzle can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## **TEMPERATURE INVERSIONS**

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## **WIND**

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### **Handheld Technology Applications:**

- Take precautions to minimize spray drift.

## **SPRAY MIXING AND COMPATIBILITY**

In general, begin with clean spray equipment and add one-half of the required amount of water to the spray or mixing tank and start agitation. Add the required quantity of fungicide and the tank-mix partner if applicable to the water and complete filling with water to the required total volume. Follow the recommendations of your State Cooperative Extension Service for tank-mixing with other products. In general, follow the order beginning first with water soluble packaging (wait for it to completely dissolve), wettable powders and water-dispersible granular products, liquid flowables and suspension concentrates and emulsifiable concentrates last. Maintain agitation throughout spraying. Do not allow spray mixture to remain in the tank overnight, or for long periods during the day without agitation.

The crop safety of all potential tank-mixes with Chipco® 26019T Fungicide including additives and other pesticides has not been tested on all crops. Follow more restrictive labeling of any tank mix partner. Do not tank mix with any product which contains a prohibition on tank mixing. Before applying any tank-mixture not specifically identified on this label, confirm safety to the target crop. Chipco 26019T Fungicide is compatible with most commonly used fungicide, herbicide, insecticide, and foliar nutrient products. However, the physical compatibility of Chipco 26019T Fungicide with all potential tank-mix partners has not been fully investigated. If tank mixing with other pesticides is desirable, conduct a jar test with the volumes and rates typically used in agricultural application. Using a small container of water, add the proportionate amounts of the products: wettable powders and water-dispersible granular products first, then liquid flowables and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

## **RESISTANCE MANAGEMENT**

The active ingredients in Chipco 26019T Fungicide belong to the dicarboxamides (Group 2) and the QoI inhibitor class of strobilurins (Group 11). Any fungal population may contain/develop individuals naturally resistant to Chipco 26019T and other Group 11 and/or Group 2 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same areas. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of CHIPCO or other Group 2 (iprodisone) and Group 11 (trifloxystrobin) fungicide within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and which considers host plant resistance, impact of

environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-866-99BAYER (1-866-992-2937). You can also contact your pesticide distributor or university extension specialist to report resistance.

## **TURF SPECIFIC USE DIRECTIONS**

### **PLANT HEALTH**

Chipco 26019T Fungicide used at the prescribed labeled rates protects the turf against winter stress resulting in faster spring green up. When applied during the spring, summer, or fall, Chipco 26019T Fungicide may result in healthier, denser, and greener turf and alleviates summer stress.

Chipco 26019T Fungicide is a foliar applied fungicide, specified for turfgrass disease control on golf courses, sod farms, and institutional areas where fine turf is grown. When used in conjunction with good turf management practices, Chipco 26019T Fungicide is effective in controlling the following diseases:

**Spring, Summer And Fall Diseases:** Dollar Spot, Brown Patch, Large Patch, Fusarium Blight and Necrotic Ring Spot\*, Leaf Spot and Melting Out, Microdochium Patch, Pink Patch, Rapid Blight, Corticum Red Thread, Curvularia Blight, Anthracnose (suppression) and Rust.

**Winter Diseases:** Gray Snow Mold

Apply the specified rates as indicated in the table in 0.5 to 10 gallons of water per 1,000 sq ft. Do not drench. Do not allow the spray mixture to stand for longer than 12 hours as some breakdown of the product may occur. Maintain agitation during spray operations. Apply with a properly calibrated sprayer.

\*Not registered for use in California to control this disease.



# TURF

## APPLICATION DIRECTIONS

Begin applications when conditions favor disease development or when the disease first appears unless otherwise noted.

DISEASE	INTERVAL OF APPLICATIONS	Fl oz of product/1,000 sq ft
<b>Dollar Spot</b> <b>Brown Patch</b> <i>(Rhizoctonia solani)</i> <b>Leaf Spot and Melting Out</b> caused by <i>Bipolaris</i> , <i>Exserohilum</i> , <i>Dreschlera</i> and <i>Marielliota</i> spp. (formerly grouped as Helminthosporium Leaf Spot diseases) <b>Rapid Blight</b> <b>Rust</b>	<b>Greens and Tees:</b> Repeat at 14 to 21 day intervals as long as required.  <b>Fairways and Other Turf Areas:</b> Repeat at 14 to 28 day intervals as long as required.	3 to 5   <b>NOTE:</b> On Fairways, for Dollar Spot Control use 2 to 4 fl oz/1,000 sq ft
<b>Large Patch</b> <i>(Rhizoctonia solani)</i>	Make first application in fall when conditions are favorable for disease development and there are no symptoms visible. Make repeat applications in spring as needed on a 14-21 day interval.	4
<b>Fusarium Blight</b> <i>(Fusarium spp.)</i> <b>Necrotic Ring Spot*</b> <i>(Ophiosphaerella korrae)</i>	Use only preventative foliar applications when conditions first become favorable for disease development. Make additional applications as necessary at 28 day intervals.	7
<b>Microdochium Patch</b> <i>(Microdochium nivale)</i> formerly called 'Fusarium Patch'	Repeat at 14 to 21 day intervals as long as required.	4 to 7
<b>Gray Snow Mold</b> <i>(Typhula spp.)</i> <b>Pink Snow Mold</b> <i>(Microdochium nivale)</i>	One application before lasting winter snow cover; a second application may be applied if there is an extended period of cold wet weather before lasting snow cover. Apply again in the spring after snow melt.	4 to 7
<b>Corticium Red Thread</b> <i>(Laetisaria fuciformis)</i> <b>Pink Patch</b> <b>Curvularia</b> <i>(Curvularia sp.)</i>	Use as a preventative every 14 days as long as required.	3 to 4 5 to 7
<b>Anthracnose (suppression only)</b> <i>(Colletotrichum cereale)</i>	Will provide suppression during periods of anthracnose pressure; mix Chipco 26019T Fungicide with CHIPCO SIGNATURE or other anthracnose control fungicide. Repeat at 14-day intervals as long as required.	4 to 7

Do not exceed a total of 7 fl oz product/1000 sq ft per application (5.1 lb iprodione and 0.31 lb trifloxystrobin per acre).

Do not exceed a total of 30 fl oz product/1,000 sq ft per year.

Do not make more than 6 applications per year.

Under severe conditions, use the higher rate and/or shorter interval of applications for all diseases. When disease pressure is light to moderate, use the lower rates and longer intervals.

Do not mow or irrigate treated areas until the foliage is completely dry, usually a 24 hour waiting period following treatment is preferred.

Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock or poultry.

\*Not registered for use in California to control this disease.

## **TANK MIXTURES**

### **ADDITIONAL DISEASE CONTROL**

If turf is threatened by additional diseases, Chipco 26019T Fungicide is compatible with most commonly used fungicides such as ProStar<sup>®</sup>, and Chipco Triton<sup>®</sup> FLO. If a tank mixture is used, follow label directions for the use of that product. When tank-mixing with other products, it is the responsibility of the end-user/applicator to ensure that the tank-mix partner is registered in the state where the application is being made.

Do not exceed a total of 30 fl oz Chipco 26019T Fungicide per 1,000 sq ft per year with a maximum of 6 applications.

Do not exceed the maximum labeled rate for any tank mix partner.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**Summer Stress Complex/Summer Decline:** For management of Summer Stress Complex/Summer Decline, apply Chipco<sup>®</sup> Signature<sup>™</sup> Fungicide at 4 to 8 oz of product per 1,000 sq ft with Chipco 26019T Fungicide at 3 to 7 fl oz of product per 1,000 sq ft.

**Pythium Blight:** Pythium blight will be controlled by the tank mixing of CHIPCO<sup>®</sup> Signature<sup>™</sup> Fungicide or ALIETTE<sup>®</sup> WDG brand Fungicide or Banol<sup>®</sup> Fungicide with Chipco 26019T Fungicide. If a tank mixture is used, follow label directions for the use of that product and apply at the rate labeled for control of the target disease organism.

**Gray Snow Mold:** Tank-mix Interface with a labeled rate of fungicide for the preventive control of Gray Snow Mold such as Chipco Triton<sup>®</sup> FLO (0.75-0.85 fl oz/1000 sq ft).

Application must be made in autumn before snow cover occurs. Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1,000 sq ft). Use the higher rate if the turf remains frozen prior to snow cover. For optimal control, reapply this treatment if a winter thaw and loss of snow cover occurs.

Not all products are registered in all states; please verify state registration of ProStar<sup>®</sup>, Chipco Triton<sup>®</sup> FLO, Chipco<sup>®</sup> Signature<sup>™</sup> Fungicide, ALIETTE<sup>®</sup> WDG brand Fungicide, and Banol Fungicide in your state before selling, distributing, or using.

## **STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

### **PESTICIDE STORAGE**

Store in a cool, dry place.

### **PESTICIDE DISPOSAL**

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste representative at the nearest EPA regional office for guidance.

### **CONTAINER HANDLING**

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

## **IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

This product contains a chemical known to the state of California to cause cancer.

Bayer (reg'd), the Bayer Cross (reg'd), Chipco®, ProStar®, Chipco Triton®, INTERFACE™ and Backed by Bayer™ are trademarks of Bayer.

## **NET CONTENT:**

PRODUCED FOR



**Bayer Environmental Science**

A Division of Bayer CropScience LP  
800 N. Lindbergh Bvd.  
St. Louis, MO 63137  
1-866-99BAYER (1-866-992-2937)

Chipco 26019T Fungicide (Pending) 04/23/2019, 04/25/2019, 06/30/2022, 07/06/2022, 07/07/2022,  
05/18/2023, 05/19/2023, 06/08/2023

[Optional Marketing Claims:]

Pictures:

