



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Mr. Richard M. Gorrell
Product Registration
Bayer Environmental Science
2T Alexander Drive
Research Triangle Park, NC 27709

JUL 11 2008

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Directions for Use
EPA Reg. No. 432-4877
Application Dated June 10, 2008

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 06/10/08 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.


If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

Bayer Environmental Science



June 10, 2008

EPA08RMG0610TritconazoleNotifMissingSentence
FEDEX

Ms. Linda Arrington
Document Processing Desk (7504P)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
Arlington, VA 22202-4501

RE: Notification of an Error Correction: Added Phrase Required by the EPA from the January 11, 2007 Label Approval for Tritconazole 70 WDG Fungicide Label (432-4877)

Dear Ms. Arrington:

Bayer Environmental Science (BES) is notifying the Agency that the sentence "Intended for use only by licensed, commercial applicators" for Tritconazole 70 WDG Fungicide was added. The sentence was inadvertently left-off the label notification submitted April 1, 2008. The EPA had requested we use this wording as a condition of registration for this end-use product back on January 11, 2007.

Bayer Environme
Science
2 T.W. Alexander
RTP, NC 27709
Phone : 919.549.

The following documents are enclosed with this letter:


- Application Form 8570-1
- 1 highlighted and 2 plain copies of the revised label
- 1 electronic disc copy

If you have any questions, contact me at (919) 549-2423 or email me at mike.gorrell@bayercropscience.com.


Sincerely,

Richard M. Gorrell
Manager, Registrations

Cc: Karen Shearer
Scott Welge
Richard Rees



A Business Group of
Bayer CropScience



<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

1. Company/Product Number 432-4877	2. EPA Product Manager Mr. Tony Kish	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Triticonazole 70 WDG	PM# 22	
5. Name and Address of Applicant (Include ZIP Code) Bayer Environmental Science 2 T. W. Alexander Drive Research Triangle Park, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ <div style="text-align: center;">NOTIFICATION</div> <div style="text-align: right;">JUL 11 2008</div> Product Name _____	

<input type="checkbox"/>	Amendment - Explain below.	<input checked="" type="checkbox"/>	Final printed labels in response to Agency letter dated _____
<input type="checkbox"/>	Resubmission in response to Agency letter dated _____	<input type="checkbox"/>	"Me Too" Application.
<input checked="" type="checkbox"/>	Notification - Explain below.	<input type="checkbox"/>	Other - Explain below.

This notification is consistent with the provisions of PR Notices 98-10 and 2007-4 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

- Notification to add the mandated EPA labeling statement from the label approval dated January 11, 2007.


1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container		If "Yes" Package wgt No. per container			
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container			4. Size(s) Retail Container 6x2 pounds plastic jugs			5. Location of Label Directions <input type="checkbox"/> On labelling accompanying product	
6. Manner in Which Label is Affixed to Product				<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled <input type="checkbox"/> Other _____			

1. Contact Point *(Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)*

Name Richard M. Gorrell	Title Manager, Registrations	Telephone No. (Include Area Code) (919) 549-2420
----------------------------	---------------------------------	---

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature 	3. Title Manager, Registrations
4. Typed Name Richard M. Gorrell	5. Date <div style="text-align: right;">June 10, 2008</div>



June 10, 2008

EPA08RMG0610TritconazoleNotifMissingSentence
FEDEX

Ms. Linda Arrington
Document Processing Desk (7504P)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
Arlington, VA 22202-4501

RE: Notification of an Error Correction: Added Phrase Required by the EPA from the January 11, 2007 Label Approval for Triticonazole 70 WDG Fungicide Label (432-4877)

Dear Ms. Arrington:

Bayer Environmental Science (BES) is notifying the Agency that the sentence "Intended for use only by licensed, commercial applicators" for Triticonazole 70 WDG Fungicide was added. The sentence was inadvertently left-off the label notification submitted April 1, 2008. The EPA had requested we use this wording as a condition of registration for this end-use product back on January 11, 2007.

Bayer Environmental
Science
2 T.W. Alexander
RTP, NC 27709
Phone : 919.549.2

The following documents are enclosed with this letter:

- Application Form 8570-1
- 1 highlighted and 2 plain copies of the revised label
- 1 electronic disc copy

If you have any questions, contact me at (919) 549-2423 or email me at mike.gorrell@bayercropscience.com.

Sincerely,

Richard M. Gorrell
Manager, Registrations

Cc: Karen Shearer
Scott Welge
Richard Rees

E-LABEL Notice

The electronic copy of the label (PDF on a CD) submitted with this application has been removed by the In-processing Team for entry into the Electronic Label Library* (ELL) to make it available to OPP staff.

You should be using this e-label to conduct your label review.

Note: If the e-label is not found in the ELL then it was probably not named correctly and could not be entered into the ELL. However, the file can be retrieved from the CD which is retained by the In-processing Team.

If the e-label is not found in the ELL, contact Tom Harris, RD (308-9423), Willie Abney, AD (308-1689) or Jim Downing, BPPD (308-9071) for further information and assistance.

*** A Lotus Notes database**

$\frac{6}{15}$

ACTIVE INGREDIENT:

Triticonazole:.....70.0%

OTHER INGREDIENTS:..... 30.0%

TOTAL:.....100.0%

EPA Reg. No. 432-4877

EPA Est. No.

CAUTION

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor for treatment advice. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Have person sip a glass of water if able to swallow. • Do not give anything by mouth to an unconscious person.

For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

reatment.

NOTIFICATION

JUL 11 2008

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers of this product must wear long-sleeved shirt and long pants, chemical-resistant gloves made out of any waterproof material, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water. Drift or runoff may be hazardous to aquatic invertebrates in neighboring areas.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical-resistant gloves made out of any waterproof material, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE

Store in original container. Keep container closed. Store in an area that is out of reach of children and animals, away from food, feedstuffs, fertilizers and seed. Protect from frost and freezing.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Offer for reconditioning, if appropriate. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then puncture and dispose of in a sanitary landfill or by other procedures approved by State and local authorities.

TURF

GENERAL INFORMATION

TURF (Golf courses, Commercial turfgrass, Sod Farms and Residential Turf). Intended for use only by licensed, commercial applicators.

Triticonazole 70 WDG Fungicide is a foliar applied fungicide, recommended for turfgrass disease control on golf courses, commercial turfgrass, sod farms and residential turf where fine turf is grown. Intended for use only by licensed, commercial applicators. When used in conjunction with good turf management practices, Triticonazole 70 WDG Fungicide is effective in controlling the following diseases:

Dollar Spot	Anthracnose	Red Leafspot	Red Thread
Brown Patch	Take-All Patch	Gray Snow Mold	Large Brown Patch
Summer Patch	Rust	Pink Snow Mold	

Triticonazole 70 WDG brand Fungicide is also effective in controlling the following diseases when combined with other products:

Pythium Blight	Summer Stress Complex / Decline
----------------	---------------------------------

Begin applications when local environmental conditions favor disease development unless otherwise noted. Apply with sufficient water to obtain adequate coverage (1-5 gallons of spray solution per 1,000 sq ft).

Recommended Turf Grass Species

Kentucky Bluegrass

*Poa annua**

Tall Fescue


Fine Fescue

Bentgrass (Creeping, Colonial)

Perennial Ryegrass

Zoysia

* Allow 28 days between Triticonazole 70 WDG Fungicide applications on *Poa annua*.



Triticonazole 70 WDG Fungicide on Turf Treated with Plant Growth Regulators

Triticonazole 70 WDG Fungicide has been tested and shown no adverse effects when applied to recommended turf types with the following Plant Growth Regulators (PGR's):

PGR

CHIPCO® Proxy™ (Ethephon)

Acceptable Turf Types

Kentucky Bluegrass
Perennial Rye
Bentgrass
Tall Fescue
*Poa annua**

Primo® (Trinexapac-ethyl)

Kentucky Bluegrass
Perennial Rye
Bentgrass
Tall Fescue
*Poa annua**

Embark® (Mefluidide)

Bentgrass
Tall Fescue
*Poa annua**

*DO NOT tank mix Triticonazole 70 WDG Fungicide with a growth regulator on *Poa annua*. Apply separately. Reduce growth regulator rates or remove an application when Triticonazole 70 WDG Fungicide is applied to *Poa annua* when turf is stressed from heat, drought or overwatering.

APPLICATION DIRECTIONS

DISEASE	USE DIRECTIONS	RATE OUNCE PRODUCT PER 1,000 sq ft		NOTES
		Preventative	Curative	
Dollar Spot (<i>Lanzia</i> spp. and <i>Moellerodiscus</i> spp.)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.	0.15-0.225 ounce	0.225-0.30 ounce	
Brown Patch (<i>Rhizoctonia solani</i>)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions use the curative rate and shorter interval.	0.15-0.30 ounce	0.30 – 0.60* ounce *0.60 ounce rate may be applied if needed in Transition areas of the South under heavy disease pressure.	In conditions favoring moderate to heavy disease pressure, Triticonazole 70 WDG can be combined with other fungicides labeled for Brown Patch including ProStar® or 26GT® at labeled rates.

DISEASE	USE DIRECTIONS	RATE OUNCE PRODUCT PER 1,000 sq ft		NOTES
		Preventative	Curative	
Summer Patch (<i>Magnaporthe poae</i>)	Begin applications in spring (April-May) and repeat at 14-28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.30-0.60 ounce		
Anthracnose (<i>Colletotrichum</i> sp.)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.	0.15-0.225 ounce	0.225-0.30 ounce	
Take-All Patch (<i>Gaeumannomyces graminis</i>)	Make 1-2 fall applications (September-October) and 1-2 spring applications (April-May) depending on local conditions. Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions use the curative rate and shorter interval.	0.15-0.30 ounce	0.30 ounce	
Large Brown Patch (<i>Rhizoctonia solani</i>)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under active disease conditions use the curative rate and shorter interval.	0.15-0.30 ounce	0.30 ounce	
RED LEAFSPOT (<i>Drechslera erythrosipila</i>)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.	0.15-0.225 ounce	0.225-0.30 ounce	

11/15

DISEASE	USE DIRECTIONS	RATE OUNCE PRODUCT PER 1,000 sq ft		NOTES
		Preventative	Curative	
ALGAE	For algae suppression repeat applications at 14-28 day intervals per the following situation: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.15-0.30 ounce		
Rust (<i>Puccinia</i> spp.)	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Use lower rate at recommended intervals. Curative Treatment: Under severe disease conditions and for active disease use higher rate and shorter interval.	0.15-0.225 ounce	0.225-0.30 ounce	
GRAY SNOW MOLD (<i>Typhula</i> spp)	Make 1-2 applications late in the fall before snow cover occurs. Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.15-0.30 ounce		Triticonazole 70 WDG can be combined with the following fungicides for greater efficacy: ProStar, Chlorothalonil, PCNB, Chipco Signature or 26 GT. Use only at labeled rates.
PINK SNOW MOLD (<i>Microdochium</i>)	Make 1-2 applications late in the fall before snow cover occurs. Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals.	0.15-0.30 ounce	0.30 ounce	Triticonazole 70 WDG can be combined with the following fungicides for greater efficacy: Chlorothalonil, PCNB, Chipco Signature or 26 GT. Use only at labeled rates.
RED THREAD	Repeat applications at 14-28 day intervals per the following situations: Preventative Treatment: Where light disease pressure is anticipated, use the lower preventative rate at recommended intervals. When conditions favor heavy disease pressure, or prior history of strong disease pressure is anticipated, use the higher preventative rate at recommended intervals. Curative Treatment: Under light disease pressure use the lower curative rate and shorter interval. Under severe disease conditions use the higher rate and shorter interval.	0.15-0.225 ounce	0.225-0.30 ounce	
Do not apply more than 1.8 ounces (50.4 grams) per 1,000 sq ft of Triticonazole 70 WDG annually				

Additional Tank Mixtures

DISEASE	USE DIRECTIONS
PYTHIUM BLIGHT	Combine Triticonazole 70 WDG with either CHIPCO® Signature™ or Banol® brand fungicide for control of Pythium Blight. Apply at labeled rates.
SUMMER STRESS COMPLEX/SUMMER DECLINE	Combine Triticonazole 70 WDG with CHIPCO® Signature™ for diseases associated with Summer Decline. Apply at labeled rates.
Do not apply more than 1.8 ounces per 1,000 sq ft of Triticonazole 70 WDG annually.	

MIXING RATES

Triticonazole 70 WDG at 0.15 ounce per 1,000 sq ft

SPRAY VOLUME (Gallons per 1,000 sq ft)

Spray Tank Capacity

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
25 Gallons	3.8 Ounces	1.9 Ounces	1.3 Ounces	0.94 Ounce	0.75 Ounce
50 Gallons	7.5 Ounces	3.8 Ounces	2.5 Ounces	1.9 Ounces	1.5 Ounces
100 Gallons	15 Ounces	7.5 Ounces	5 Ounces	3.8 Ounces	3 Ounces
200 Gallons	30 Ounces	15 Ounces	10 Ounces	7.5 Ounces	6 Ounces

Triticonazole 70 WDG at 0.23 ounce per 1,000 sq ft

SPRAY VOLUME (Gallons per 1,000 sq ft)

Spray Tank Capacity

	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
25 Gallons	5.6 Ounces	2.8 Ounces	1.9 Ounces	1.4 Ounces	1.1 Ounces
50 Gallons	11.3 Ounces	5.6 Ounces	3.8 Ounces	2.8 Ounces	2.2 Ounces
100 Gallons	22.5 Ounces	11.3 Ounces	7.5 Ounces	5.6 Ounces	4.5 Ounces
200 Gallons	45 Ounces	22.5 Ounces	15 Ounces	11.3 Ounces	9 Ounces

Triticonazole 70 WDG at 0.30 ounce per 1,000 sq ft

SPRAY VOLUME (Gallons per 1,000 sq ft)

Spray Tank Capacity


	1 Gallon	2 Gallons	3 Gallons	4 Gallons	5 Gallons
25 Gallons	7.5 Ounces	3.8 Ounces	2.5 Ounces	1.9 Ounces	1.5 Ounces
50 Gallons	15 Ounces	7.5 Ounces	5 Ounces	3.8 Ounces	3 Ounces
100 Gallons	30 Ounces	15 Ounces	10 Ounces	7.5 Ounces	6 Ounces
200 Gallons	60 Ounces	30 Ounces	20 Ounces	15 Ounces	12 Ounces

NOTES

- Apply with a properly calibrated sprayer and maintain agitation during spray operations.
- Apply with sufficient water to obtain adequate coverage (1 to 5 gallons of spray solution per 1,000 sq ft).

RESTRICTIONS AND LIMITATIONS

- For golf courses only, do not apply to turf cut higher than 1 inch on golf holes where water bodies are present.
- Do not apply more than 1.8 ounces (50.4 grams) per 1,000 sq ft annually.
- Do not drench. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against mixing.
- Do not irrigate treated areas until the spray on foliage is completely dry.
- This product is compatible with most commonly used fungicides, however, check compatibility before tank mixing.
- Mixture with stickers, extenders, or wetting agents is not necessary.
- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock or poultry.
- Do not apply Triconazole 70 WDG Fungicide through any type of irrigation system.



IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.


Net Content:

Banol[®], Compass[®], Chipco[®], ProStar[®], Proxy[®], and 26 GT[®] are registered trademarks of Bayer and Signature[™] is a trademark of Bayer. Heritage[®] and Primo[®] are registered trademarks of Syngenta. Embark[®] is a registered trademark of PBI/GORDON CORPORATION.

Produced for

Bayer Environmental Science
A Business Group of Bayer CropScience LP
P.O. Box 12014, 2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Triticonazole 70 WDG Fungicide (MASTER) 01/11/07, Notification 04/01/08, Notification 06/06/08



Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic file identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

PROPOSED LABEL		
EPA Registration #	Date Submitted to EPA	Electronic file name
432-4877	06/06/08	000432-04877.20080606.Triticonazole70W DGFungicide.pdf

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.


Signature

06/06/2008

Date

Anna B. Wilkins

Name (typed)

Label Development Specialist

Title

