

STANDARD'S PASTURE MINERAL MEDICATED

DIRECTIONS FOR CONTINUOUS SELF-FEEDING PROGRAM

WHEN TO FEED: For hornfly control on pastured cattle start feeding STANDARD'S PASTURE MINERAL, MEDICATED in the spring before flies appear. Continue feeding all summer and into the fall until cold weather restricts fly development. For grub control, feed continuously for a minimum of 75 days.

HOW MUCH TO FEED: Recommended consumption of STANDARD'S PASTURE MINERAL, MEDICATED containing 6% Ronnel is 0.2 lb. per 100 lb. of body weight per head per month.

HOW TO FEED: Feed STANDARD'S PASTURE MINERAL, MEDICATED continuously free-choice to cattle in covered mineral-feeder troughs or boxes. Provide one mineral feeder box or feeding space for each 15 to 20 head of cattle. Locate mineral-feeder boxes and troughs near watering areas. To encourage uniform and normal consumption, put out only a 7-day supply at one time, and keep feeder boxes and troughs clean. For adequate insect control, each animal must eat its share of the mineral. Small, late calves may require an additional 7- or 14-day feeding program after weaning for grub control.

WARNING

During continuous feeding program, do not allow cattle to run out of STANDARD'S PASTURE MINERAL, MEDICATED and do not feed any additional salt or mineral.

DIRECTIONS FOR 7-DAY OR 14-DAY FEEDING PROGRAMS

Mix STANDARD'S PASTURE MINERAL, MEDICATED with ground grain for top dressing, or mix with the complete ration to make an insecticidal feed. Feed continuously for 7 or 14 days and make certain that all cattle are eating. Adequate feeding space should be available for each animal. Make certain that insecticidal feed is evenly distributed on, or mixed into, the total ration. Provide a total of 0.2 lb. of STANDARD'S PASTURE MINERAL, MEDICATED for each 100 lb. body weight (to be fed over the 7- or 14-day feeding period).

NOTE: Insecticidal feed made from STANDARD'S PASTURE MINERAL, MEDICATED should be used within 4 weeks of mixing.

WHEN TO FEED: After end of adult heel fly season and before grubs reach the back. Ronnel is not effective for grub control during later stages of life cycle, when grubs are in the backs of cattle. Reduction of cattle lice will be obtained when fed as directed for the 7-day program.

7-DAY TREATMENT

WHEN TO FEED: Southern area cattle — July through September.
Northern area cattle — August through October.

HOW TO FEED: Mix thoroughly 2 lb. of STANDARD'S PASTURE MINERAL, MEDICATED containing 6% Ronnel with 19 lb. feed to produce an insecticidal feed. Prepare only enough for the 7-day feeding program. Feed 0.3 lb. of insecticidal feed per 100 lb. body weight daily for 7 continuous days.

14-DAY TREATMENT

WHEN TO FEED: Southern area cattle — July through September.
Northern area cattle — August through DECEMBER.

HOW TO FEED: Mix thoroughly 1 lb. of STANDARD'S PASTURE MINERAL, MEDICATED containing 6% Ronnel with 20 lb. of feed to produce an insecticidal feed. Prepare only enough for the 14-day feeding program. Feed 0.3 lb. of insecticidal feed per 100 lb. body weight daily for 14 continuous days.

10 APR 1974

EDWARDSVILLE
PESTICIDE DIVISION

STANDARD
CHEMICAL
MFG. CO., OMAHA, NEBR.
ESTABLISHED 1904



