

Product Name
Trade Name
Date of Issue

Not constitute a part of the product
Date 7/2/55

Active Ingredients
Warnings
Substance Name
Net Content

Net Content

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Protect pets and domestic animals from bait. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. If swallowed, intravenous and oral administration is advised. If combined with blood transfusions are indicated as in the case of hemorrhage caused by strokes or hemorrhagic conditions.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning or emptying. Dispose of wastes.

Labels must be present in areas not accessible to children, pets, wild birds or domestic animals. Do not tamper, feed bait to nest.

CONTAINER DISPOSAL: Do not reuse container. Destroy when empty.

DIRECTIONS FOR USE:

For Norway Rats and Roof Rats: Place 4 to 6 ounces of bait in one or two portions in a dark, dry, unobstructed area where rats are accustomed to feed. Bait should be placed in a bait station or other container which is not accessible to children, pets, wild birds or domestic animals. As soon as the bait is placed in the bait station, it should be checked to see that it is in place and that the bait is not accessible to children, pets, wild birds or domestic animals. Bait should be checked daily and replaced as needed.

For House Mice: Place the bait in small pieces in a dark, dry, unobstructed area where mice are accustomed to feed. Bait should be placed in a bait station or other container which is not accessible to children, pets, wild birds or domestic animals. As soon as the bait is placed in the bait station, it should be checked to see that it is in place and that the bait is not accessible to children, pets, wild birds or domestic animals. Bait should be checked daily and replaced as needed.

For Field Mice: Place the bait in small pieces in a dark, dry, unobstructed area where mice are accustomed to feed. Bait should be placed in a bait station or other container which is not accessible to children, pets, wild birds or domestic animals. As soon as the bait is placed in the bait station, it should be checked to see that it is in place and that the bait is not accessible to children, pets, wild birds or domestic animals. Bait should be checked daily and replaced as needed.

1955-1956

DIRECTIONS FOR USE

RATS:

Place 4 to 6 ounces of bait in dry containers next to burrows, in runways or where rats are accustomed to feed. Assure an uninterrupted supply of bait for a period of not less than 12 days and continue baiting until all signs of feeding have stopped.

MICE:

Place tablespoonful amounts of bait in shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8-12 feet are recommended. Assure uninterrupted supply of bait for a period of not less than 15 days.

Replace stale, damp and moldy baits, with fresh baits.

IMPORTANT

Baiting should be continued as long as evidence of feeding is observed. Except in extremely heavy infestations, this should be from five days to two weeks. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

CAUTION

Place bait in locations which are not accessible to children, pets and domestic animals, or in tamper proof bait boxes. If swallowed by humans, domestic animals, or pets, this material may reduce the clotting ability of the blood and cause bleeding, in such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

This rodenticide contains warfarin, an anticoagulant chemical that reduces the clotting ability of the blood, and causes fatal hemorrhage in rats upon repeated feedings, it also contains an antibacterial agent sulfaquinoxaline, which inhibits bacteria that produce vitamin K, an antidote for warfarin. Do not reuse container. Destroy when empty.

E.P.A. REGISTRATION NO. 445-19

ROSEEX
Contains PROLIN a better, more effective
WARFARIN rodenticide
Ready to Use

**KILLS RATS
and MICE**

ACTIVE INGREDIENTS:

Warfarin [3-(alpha-acetonylbenzyl)-4-hydroxycoumarin] 0.025%
N-(2-Quinoxaliny) Sulfanilamide 0.025%
Sulfaquinoxaline) 99.950%

INERT INGREDIENTS

Prolin (warfarin plus sulfaquinoxaline) is manufactured under license from the Wisconsin Alumni Research Foundation, U. S. Patent No. 2,427,578, other patent applications pending.

NET CONTENTS — 1 POUND

**CAUTION: Keep out of reach of children.
See other caution on back panel.**

PREPARED BY

ROSE EXTERMINATOR CO.

100 EAST COURT STREET
CINCINNATI, OHIO
PHONE 721-6682

ACCEPTED WITH COMMENTS