

FRUIT INSECTS — CHERRY FRUIT FLY: Apply 1/2 ounce per 100 to 125 sq. ft. of water. Repeat applications at 2 to 3 week intervals until harvest.

PLUM CURCULIO ON PEACHES, PLUMS & APPLES: Apply 1/2 ounce per 100 to 125 sq. ft. of water. Repeat applications at 2 to 3 week intervals until harvest.

GRAPEBERRY MOTH: Apply 1/2 ounce per 100 to 125 sq. ft. of water. Repeat applications at 2 to 3 week intervals until harvest.

CAUTION: Do not use on plants in flower. Do not use on plants in fruit. Do not use on plants in seed.

NOTICE: Do not use on plants in flower. Do not use on plants in fruit. Do not use on plants in seed.

Manufactured For
WOODBURY CHEMICAL COMPANY
 Division of TECHNE CORP
 ST. JOSEPH MO 64502 DENVER COLO

Woodbury's
 WOODBURY
 CHEMICAL COMPANY

ACCEPTED

METHOXYCHLOR "50"

HORN FLIES: FOR USE ON BEEF CATTLE: Use 1/2 ounce per 10 to 12 sq. ft. of water. Apply 2 quarts per animal. Repeat applications at 2 to 3 week intervals until harvest.

For dusting on Beef Cattle and Dairy Cattle: Use 1 heaping tablespoon of powder per 100 to 125 sq. ft. of water. Repeat applications at 2 to 3 week intervals until harvest.

Warning: Do not use on plants in flower.

STABLE FLIES IN BARN, ANIMAL PENS AND FARM BUILDINGS: Use 1/2 ounce per 100 to 125 sq. ft. of water. Apply to all surfaces where flies rest or crawl. Repeat applications at 2 to 3 week intervals. Care should be taken to avoid contamination of milk and other equipment.

BEEF CATTLE LICE, HOG LICE AND SHEEP TICKS: Use 1/2 ounce per 10 to 12 sq. ft. of water. Spray on all parts of the animal. Repeat applications at 2 to 3 week intervals until harvest.

VEGETABLE INSECTS: Apply 1/2 ounce per 100 to 125 sq. ft. of water. Repeat applications at 2 to 3 week intervals until harvest.

Apply to all parts of the plant. Repeat applications at 2 to 3 week intervals until harvest.