

[449-114]

**ALFALFA AND CLOVER INSECTS:** For control of spotted alfalfa aphid, pea aphid, leafhoppers, apply 1½ pints MALATHION per acre in sufficient water for coverage. Make no application within 7 days of pasturing or harvest, and do not apply to clover in bloom.

**INSECTS ON FRUITS:** For control of aphids, bud moth, codling moth, Forbes scale, plum curculio, red banded leaf roller and spider mites on apples, apricots, peaches and pears, apply MALATHION at rate of 1½ - 2 pints per 100 gal. water (2 tsp. per gal.). Do not apply within 7 days of picking.

**INSECTS ON VEGETABLES:** For control of aphids, cabbage looper, imported cabbage worm, leafhopper, Mexican bean beetle, spider mites on beans, beets, cabbage, celeris, eggplant, peas, peppers, potatoes, apply 1 pt. MALATHION per acre in sufficient water (1 tsp. per gal.). Do not apply within 7 days of harvest.

**INSECTS ON ORNAMENTALS:** For control of aphids, lace bug, leafhoppers, mealybugs, tarnished plant bug, thrips, whitefly, apply 1½ pts. MALATHION per 100 gal. water (1 tsp. per gal.). For control of bagworm, red spider mite, certain scale insects, use 2 pts. MALATHION per 100 gal. water (2 tsp. per gal.).

Avoid the use of MALATHION on certain ferns, including Boston, Maiden Hair, and Pteris and some species of Crassula.

**FOR CONTROL OF LICE ON CATTLE AND HOGS:** One gallon MALATHION per 100 gal. water. Cover completely. One application may be sufficient. Repeat only if needed. Spray hog houses, pens and litter.

**DO NOT APPLY TO LACTATING DAIRY ANIMALS.**

**DO NOT TREAT CALVES UNDER ONE MONTH OF AGE.**

**TICKS ON CATTLE, SHEEP AND GOATS:** 1 - 2 gallons MALATHION per 100 gal. water, apply as complete coverage. Repeat at two week intervals if needed. Do not apply to milk quats.

**FOR CONTROL OF NORTHERN FOWL MITE, POULTRY LICE:** Mix two tablespoonfuls MALATHION per gallon of water per 100 - 150 birds. Repeat 4 - 8 weeks or when necessary.

**AVOID CONTAMINATION OF FEED, FOODSTUFFS, FEEDING TROUGHS AND WATER FOUNTAINS.**

**FOR CONTROL OF CHICKEN RED MITE, POULTRY LICE:** Apply 4 tablespoonfuls MALATHION per gal. water liberally to walls, ceilings, roosts, nests, and adjacent areas. Force spray into all cracks and crevices.

**FOR CONTROL OF POULTRY TICKS:** Apply 6 - 7 ounces per gallon as above.

**ROOST PAINT:** For Chicken Red Mite and Poultry Lice mix 4 tablespoonfuls MALATHION per gal. of water. Brush on at rate of 1 pint per 150 feet of roost. Repeat as necessary.

## DIRECTIONS

**HORN FLIES ON BEEF CATTLE:** 1 - 1½ gal. MALATHION to 100 gal. water. Cover completely. Repeat at 2-week intervals if needed.

**FLY CONTROL: Straight SPRAY —** Mix MALATHION with water as follows: 1 oz. with 1½ qts., 1 pt. with 6 gal., 2 gal. with 100 gal., and apply spray at 1 gal. per 1000 sq. ft. on painted surfaces, and 2 gal. per 1000 sq. ft. on unpainted surfaces. Apply where flies light or congregate.

**BAIT SPRAY:** Mix MALATHION as follows: 1 oz. with 2 tbsp. sugar in 1 gal. water; 1 pt. with 3 cups sugar in 6 gal. water; 2 gal. with 20 lbs. sugar in 100 gal. water.

MALATHION may be used in and around all farm buildings EXCEPTING milk rooms. Do not apply to freshly white-washed surfaces — wait 14 days.

**FOR SPACE AND CONTACT SPRAY—** Outdoors: Flies, Mosquitoes and small flying insects: Use 2% to 5% Malathion fog, or space spray. For 2% spray, dilute 55% Malathion in 1 to 28 parts of water, kerosene oil, or diesel oil. When using kerosene type solvent as carrier, dilute 1 part of 55% Malathion in 28 parts of a mixture consisting of 4 parts kerosene-type solvent and 1 part aromatic hydrocarbon-type solvent. For 5% spray, dilute 1 to 11 using similar solvents.

NOTICE: Buyer assumes all risks of trial not in strict accordance with directions given herewith.


## CAUTION

Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Remove animals and poultry from buildings before treating.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning. (Keep out of Smoke.)

USDA Reg. No. 449-114



Buyer assumes all risks of  
not in strict accordance with  
instructions given herewith.

### CAUTION


If swallowed. Avoid prolonged  
contact with spray mist. Avoid prolonged  
contact with skin; wash  
after using. Avoid contamina-  
tion of foodstuffs. Keep out of  
reach of children. Remove animals and  
livestock from buildings before treating.

Do not use on or near  
water. Do not use in  
open body of water. Keep  
away from water supplies.  
Apply this  
product as specified on this label.

Equipment and containers and  
wastes by burying in non-  
away from water supplies.  
should be disposed of by punch-  
ing holes in them and burying with  
soil by burning. (Keep out of  
reach of children.)

Pat. No. 449-114

ACCE  
UNDER THE FEDERAL  
FUNGICIDE AND RO  
FOR ECONOMIC POIS  
E. UNDER NO  
TO ATTACHED COMM



CONTENTS  
GALLON

SURE DEATH BRAND

# MALATHION W-E

Containing 5 Pounds Actual MALATHION

For Control of Certain Insects Attacking Alfalfa and Clover,  
Vegetables and Ornamentals. For Control of Flies Resistant  
to DDT and Other Chlorinated Insecticides

#### ACTIVE INGREDIENTS:

Malathion*	55%
Xylol	35%
INERT INGREDIENTS	10%

\*O.O.s.m. (O.O.s.m. is a trademark of O.O.s.m. Manufacturing Co.)

**WOODBURY  
CHEMICAL  
COMPANY • St. Joseph, Mo.**

**CAUTION** KEEP OUT OF REACH OF CHILDREN SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY LABELING