


[449-173] 192



Powdery Standard

LEAD ARSENATE

LEAD ARSENATE

U. S. Patent 2,088,651

DANGER!!
KEEP OUT OF REACH OF CHILDREN



SEE SIDE PANEL FOR ANTIDOTE STATEMENT
AND ADDITIONAL PRECAUTIONARY LABELING

DIRECTIONS

TURF

SOIL TREATMENT FOR LAWN AND ORNAMENTAL AREAS: For Japanese Beetle Grubs, White Grubs and certain other grubs, apply 1 lb. per 100 sq. feet of lawn area, either as a spray in water, or dry. Spread or spray uniformly over the area to be treated, then wash insecticide (either dry or spray) off grass blades into soil by spraying lawn with clear water. When preparing lawn for seeding, the material may be raked into the soil to a depth of 1 or 2 inches about 30 days before planting grass seed.

Keep children and pets off treated areas.

WARNING: May be fatal if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing, natural synthetic rubber gloves, and goggles. If spilled on skin, remove contaminated clothing and wash skin immediately with soap and warm water. Do not breathe fumes, dust, spray mist. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for protection against lead arsenate. Information as to suitable types of masks or respirators is available from the U. S. Department of Agriculture. Do not contaminate feed or foodstuffs. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Vacated areas should not be reentered until the insecticide has dissipated. Keep out of reach of children. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning. (Keep out of smoke.)

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Stomach pains. Vomiting and diarrhea.

ANTIDOTE: Internal: Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician immediately. External: If the skin is contaminated, wash immediately with soap and warm water.

DIRECTIONS

MIXING: Partly fill the spray tank with water, start the engine and gradually add the required amount of Lead Arsenate as the tank is filled.

APPLICATION: Thoroughly wet all exposed plant parts with spray mixture.

APPLES AND PEARS: For Codling Moth, Plum Curculio, Tent Caterpillar, and certain other chewing insects, use 1 pound together with 1 pound Hydrated Lime (Spray Grade) in 33 gallons water. For small sprayers, use 2 level tablespoonfuls with an equal amount of Hydrated Lime in 1 gallon water. Apply sprays at petal fall and repeat every 7 to 14 days as long as necessary. Do not apply within 40 days of harvest. NOTE: Residues remaining at harvest should be removed by washing, wiping, or other effective means.

FOR DUSTING: Thoroughly mix 1 part Lead Arsenate and 1 part Hydrated Lime. Apply uniformly as often as necessary to keep plants protected. Dusts may be used on tomatoes, tobacco, and ornamental plants. Precautions given under the above crop headings should be observed. NOTE: RESIDUES remaining at harvest should be removed by washing, wiping, or other effective means.

This application will control same insects listed under Tomatoes, and Tobacco mentioned previously. Follow same cautions.

Do not graze livestock in treated groves or orchards.

TOMATOES: For Hornworm, Colorado Potato Beetle and certain other chewing insects, use 1 pound in 33 gallons water, or 2 level tablespoonfuls in 1 gallon water. Apply when the insects first appear and repeat as necessary to keep plants protected. Do not apply to tomatoes within 40 days of harvest. NOTE: Residues remaining at harvest should be removed by washing, wiping, or other effective means.

FOR HORNWORM: Use 1 pound in 12 gallons water and apply to 1/2 acre. For large plants more than 12 gallons of spray mixture may be needed. Apply spray when insects first appear and repeat as necessary to keep plants protected. Use of insecticides may leave undesirable residues and every precaution should be taken in spraying tobacco to avoid excessive residues.

SHADE TREES AND ORNAMENTAL PLANTS: For Japanese Beetle and certain other chewing insects, use 1/2 to 1 pound in 16 gallons water, or 2 to 4 level tablespoonfuls in 1 gallon water, with an equal amount of Hydrated Lime (Spray Grade). Apply spray when insects first appear and repeat as necessary to keep plants protected.

ACTIVE INGREDIENTS: DRY LEAD ARSENATE..... 98%
INERT INGREDIENTS..... 2%
Total Arsenic Expressed as Metallic Arsenic (As)..... 20.50%
Total Lead Expressed as Metallic Lead (Pb)..... 58.50%
Arsenic in Water-Soluble Form Expressed as
 Metallic Arsenic (As)..... .25%

[442-173]282

CONDITIONS OF SALE AND WARRANTY. The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application even though the product is used in accordance with Directions. All such risks shall be assumed by the Buyer.

Woodbury Chemical Co. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the inherent risks referred to above. Woodbury makes no express or implied warranty of fitness or merchantability or any other express or implied warranty. In no case shall Woodbury or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Woodbury and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty which may be varied only by agreement in writing signed by a duly authorized representative of Woodbury Chemical Company.

Manufactured by

WOODBURY CHEMICAL COMPANY

St. Joseph, Mo.

Denver, Colo.

Orlando, Fla.

Lubbock, Texas

USDA Reg. No. 449-173