


CONTENTS

1 QUART



# LIME SULPHUR

## INSECTICIDE and FUNGICIDE

### FOR SHRUBS, ROSES, FRUIT AND SHADE TREES

#### ACTIVE INGREDIENTS

Calcium Polysulfide 29  
(Total Combined sulphur not less than 23%) 71

#### INERT INGREDIENTS

Total 100

#### DIRECTIONS

see mixing table for proper dilutions

- (1) **SUMMER FOLIAGE STRENGTH.** Roses For control of Black Spot, Powdery Mildew, Rust and Red Spider, apply at 10 to 14 day intervals in growing season.
- (2) Grapes For control of Powdery Mildew and Anthracnose, apply when shoots are 6 to 8 inches long and again after bloom. Existing foliage may be somewhat damaged.
- (3) Peaches For control of Brown Rot and Scab prevention, make one application at petal fall stage. Do not use on Apricots.
- (4) Currants and Gooseberries For control of Powdery Mildew, apply just before and after blooming.
- (5) **EARLY SPRING STRENGTH.** As growth starts, Rose For control of Rose and San Jose Scales, Case Bearers and Powdery Mildew, apply in Fall after leaves drop and again in Spring before the buds swell.

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY LABELING

**DO NOT STORE WHERE PRODUCT WILL FREEZE!!!**

- (6) Raspberries and Blackberries: For control of Red Berry Mites, Anthracnose, Powdery Mildew, and Rust, make one application when new canes are 8 to 12 inches tall. Repeat in Fall for Red Berry Mites.
- (7) **DORMANT STRENGTH.** Apples, Pears, Plums and Cherries: For San Jose Scale control, use before buds swell in Spring. For Pear Leaf Blister Mites, delay until buds start to swell as mites overwinter in the bud.
- (8) Peaches For control of Peach Leaf Curl, San Jose Scale, Brown Rot and Twig Borer, apply once in late Fall, repeat in Spring just before buds swell. Do not use on Apricots.
- (9) Shade Trees and Shrubs: For control of San Jose Scale, Powdery Mildew, Canker, Anthracnose, Maple Gall, Leaf Blotch, and Olive Scale, apply once in early Spring before buds swell. Do not apply to Euonymus, Euonymus, or Rhododendron or allow to drift on these susceptible species.

#### MIXING TABLE AND DIRECTIONS

Amount of Spray Desired	Dormant	Early Spring	Summer Foliage
1 Quart	6 tablespoons	4 tablespoons	1/2 teaspoon
1 Gallon	3/4 pint	5 tablespoons	3 teaspoons
3 Gallons	2 1/2 pints	1/2 pint	3 tablespoons
	(5 cups)		

#### CAUTION

Avoid contact with hands and face. If splashed into eyes, wash at once with pure water. Do not use on foliage when temperatures exceed 85°. Do not use an oil spray within three weeks following a Lime Sulphur treatment. Do not spray in the heat of the day. Take care spray drift does not contact light-painted surfaces as dark stains or spots may result. Rinse all equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning. (Keep out of Smoke.)

NOTICE: Seller's responsibility is limited to accuracy of contents and confirmation to the chemical description. Buyer assumes all risk and liability for use, storage, and handling, and accepts and uses this product under these conditions.

Manufactured by  
**WOODBURY CHEMICAL COMPANY**  
St. Joseph, Mo. Denver, Colo. Orlando, Fla.

USDA Reg. No. 449-221

ACCEPTED WITH COMMENTS

may 25 1964

Comment  
449-221