

NET WEIGHT

30 POUNDS




COVERS 2,000 SQUARE FEET
CONTROLS CRABGRASS
KILLS SOIL INSECTS

FERTILIZES LAWNS IN ONE APPLICATION
Contains DACTHAL: U.S. Patent No. 2,923,634

LAWN GUARD

WITH 11-11-11 FERTILIZER
GUARANTEED 90-100% CRABGRASS CONTROL
when used as directed.



ACTIVE INGREDIENTS
Dacthal 100% 1.00%
Fertilizer 11-11-11 1.00%
INERT INGREDIENTS 98.00%

CAUTION - KEEP OUT OF REACH OF CHILDREN

Controls crabgrass and certain other weeds if applied in time to kill before the germination of their seeds. Kills many insects present in the soil, and provides winter spring application of excellent fertilizer. Will not harm established bluegrass.

WEEDS CONTROLLED: Carpetweed, Hairy crabgrass, Smooth crabgrass, Green and yellow foxtails, Florida Pigeon, or hen foxtail, Johnson grass from seed, Lambsquarters, Livegrass, Purslane, Witchgrass, and hen foxtail.

INSECTS CONTROLLED: Ants, Chiggers, Chinch bugs, Japanese beetle larvae, and webworms. Will injure White grubs. Water lawn before application and do not water again for 2-3 days.

FERTILIZER: Guaranteed Analysis—Nitrogen 11% Available Phosphoric Acid 11% Available Potash 11%

DIRECTIONS FOR USE

LAWN GUARD must be applied before weed seeds germinate. Application may be made in late winter or early spring generally after the first Forsythia blooms. Apply LAWN GUARD with a fertilizer spreader at 5-6 pounds per 1,000 square feet. A spot application of 7.5 pounds per 1,000 square feet in one direction and a second at the same rate at right angles will give more even coverage. If some crabgrass is still present, it is recommended that grass be mowed as early as possible and application of LAWN GUARD be made in time to kill the seeds of the newly sprouting grass. LAWN GUARD may be applied any time of the year for fertilizer. At any time after application, this delay may reduce the survival percentage of the crabgrass seeds. Do not apply to wet soil. Water thoroughly after application. Do not allow children to start on soil surface after application.

PRECAUTIONS: Insects and weeds germinating after application of LAWN GUARD in the spring should not be applied until they are 60 days old. Do not use on lawns until after the first of July.

SETTINGS FOR POPULAR FERTILIZER SPREADERS

Model	Rate (lb./1,000 sq. ft.)	HALE RATE (lb./1,000 sq. ft.)
John Deere 1000	3	7 1/2
John Deere 1000	4	10
John Deere 1000	5	12 1/2
John Deere 1000	6	15
John Deere 1000	7	17 1/2
John Deere 1000	8	20
John Deere 1000	9	22 1/2
John Deere 1000	10	25
John Deere 1000	11	27 1/2
John Deere 1000	12	30
John Deere 1000	13	32 1/2
John Deere 1000	14	35
John Deere 1000	15	37 1/2
John Deere 1000	16	40
John Deere 1000	17	42 1/2
John Deere 1000	18	45
John Deere 1000	19	47 1/2
John Deere 1000	20	50
John Deere 1000	21	52 1/2
John Deere 1000	22	55
John Deere 1000	23	57 1/2
John Deere 1000	24	60
John Deere 1000	25	62 1/2
John Deere 1000	26	65
John Deere 1000	27	67 1/2
John Deere 1000	28	70
John Deere 1000	29	72 1/2
John Deere 1000	30	75
John Deere 1000	31	77 1/2
John Deere 1000	32	80
John Deere 1000	33	82 1/2
John Deere 1000	34	85
John Deere 1000	35	87 1/2
John Deere 1000	36	90
John Deere 1000	37	92 1/2
John Deere 1000	38	95
John Deere 1000	39	97 1/2
John Deere 1000	40	100

CAUTION: Avoid breathing dust. Do not use on lawns with children. Do not use on lawns with children. Avoid contact with eyes. Wash after handling or using. Keep out of reach of children. Do not use on lawns with children. Wash after handling or using. Do not use on lawns with children. Wash after handling or using.

Do not apply to newly seeded lawns, or to dandelions at any time. Some thinning of fescue grasses has occasionally been observed. Birds feeding on treated areas may be killed. Rinse equipment and containers and dispose of waste by burning.

ACCEPTED
 Feb 20 1969
 UNDER THE FEDERAL INSECTICIDE
 ACT AND THE FERTILIZER ACT
 REGISTERED IN THE FERTILIZER REGISTER
 FOR THE PESTICIDE SUBJECT
 TO ATTACHED COMMENTS.

449-29

COVERS 2,000 SQUARE FEET
 CONTROLS CRABGRASS
 KILLS SOIL INSECTS

FERTILIZES LAWNS IN ONE APPLICATION
 Contains DACTHAL: U.S. Patent No. 2,923,634

LAWN GUARD

WITH 11-11-11 FERTILIZER
 GUARANTEED 90-100% CRABGRASS CONTROL
 when used as directed.

ACTIVE INGREDIENT
 DACTHAL (1,1-DIBROMO-2,2-DICHLOROETHANE) 1.5%
 FERTILIZER 11-11-11 11.0%
 INERT INGREDIENT 87.5%
 TOTAL 100.0%

CAUTION KEEP OUT OF REACH OF CHILDREN

Controls crabgrass and certain other weeds if applied in the spring before the germination of these seeds has occurred. Also kills many insects present in the soil, and provides a full spring application of balanced fertilizer. Will not injure established bluegrass.

WEEDS CONTROLLED: Carpetweed, Hairy crabgrass, Smooth crabgrass, Common chickweed, fall panicum, Florida Pusley, green foxtail, Johnson grass from seed, Lambsquarters, Lovegrass, Purslane, Witchgrass, Yellow foxtail.

INSECTS CONTROLLED: Ants, Chiggers, Chinch bugs, Japanese beetle larvae, sod wireworms, White grubs, Wireworms. Water lawn before application and do not water again for 2-3 days.

FERTILIZER: Guaranteed Analysis—Nitrogen 11%, Available Phosphoric Acid 11%, Available Potash 11%

DIRECTIONS FOR USE

LAWN GUARD must be applied before weed seeds germinate. Application may be made in late winter or early spring, generally no later than the time Forsythia blooms. Apply LAWN GUARD with a fertilizer spreader at 15 lbs. per 1,000 square feet. A split application of 7.5 pounds per 1,000 square feet in one direction, and a second at the same rate, applied at right angles, will give more even coverage. If spring seeding is desired, it is recommended that grass be seeded as early as possible, and application of LAWN GUARD delayed until there is greening of the newly sprouted grass. Or, LAWN GUARD may be applied early, and seeding delayed for at least 60 days after application. This delay improves the survival percentage of the desirable grasses. Do not apply when grass is wet. Water thoroughly after application. Do not rake or disturb the soil surface after application.

ROSE BEDS: In rose beds crabgrass may be controlled by application of LAWN GUARD in the spring at one pound to approximately 60 sq. ft. Do not hoe or otherwise disturb soil surface until after June 15th.

SETTINGS FOR POPULAR FERTILIZER SPREADERS

	Full Rate	HALF RATE (twice over)
Central Imperial	9	7 1/2
Cyclone (7 ft. spreader)	5.0	4.5
Jackson	8	6 1/2
Lawn Broom	10	6 1/2
Modern 340G	9	7 1/2
Modern 350G	7 1/2	6
Modern 360G	7 1/2	6
Scott	7	7 1/2
Smars Cross Country	8	6 1/2

CAUTION: Avoid breathing dust. Do not eat, drink, or use as foodstuffs. Do not take internally. Avoid contact with eyes. Wash after handling or using. Keep children away from all treated areas until insecticide is washed into soil and grass is thoroughly dry.

Do not apply to newly-seeded lawns, or to dicamba at any time. Some thinning of fescue grasses has occasionally been observed. Birds feeding on treated areas may be killed. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning (Keep out of smoke).

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured by

WOODBURY CHEMICAL COMPANY

St. Joseph, Mo. Denver, Colo. Orlando, Fla. Lubbock, Tex.