

CAUTION

Avoid inhalation of dust or spray mist. Avoid contact with skin. Avoid contamination of feed and foodstuffs. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

COMPATABILITY

Captan 50W can be combined effectively at recommended dosages with most commonly used fungicides and insecticides with the exception of oil and strongly alkaline materials including calcium arsenate. Observe all cautions and limitations on labeling of all products used in mixtures.

RECOMMENDATIONS

The dosage rates on this label indicate quantity of Captan 50W per gallon of dilute spray unless otherwise stated.

APPLE: Scab, Frogeye — Use 4 to 8 tsp. at 5 to 7 day intervals as needed to maintain cover in the pre-bloom, bloom, petal-fall, first and second cover sprays. Use at 8 tsp. rate if conditions favor scab or if active scab can be found; Scab, Brooks Fruit Spot, Sooty Blotch, Fly Spec, Black Rot, Black Pox, Bitter Rot — Use 4 to 8 tsp. at 10 to 14 day intervals in the third and later cover sprays. Use at the 8 tsp. rate if conditions favor fruit rots and sooty blotch. Continue applications to within 30 days of harvest to control sooty blotch; up to start of harvest to control fruit rots.

PEACH: Brown Rot, Scab — Use 8 tsp. in the full pink, bloom, petal-fall, shuck, cover and pre-harvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 10 day intervals as needed to maintain cover. Do not apply within one day of harvest.

Woodbury's
WOODBURY
CHEMICAL COMPANY

CONTENTS ONE POUND

CAPTAN 50-W

ACTIVE INGREDIENT:

CAPTAN (N-trichloromethylthio-4-cyano-1,4-dithiazole-2-dithiocarbamate) 50%

INERT INGREDIENTS 50%

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Precautions

ACCEPTED
JUL 11 1968
UNDER THE FEDERAL
INSECTICIDE AND RODENTICIDE
ACT FOR ECONOMIC POISON
ED UNDER NO. 49-247

GRAPE: Downy Mildew, Black Rot — Use 8 tsp. just before bloom, just after bloom, and 7 to 10 days after bloom. Make an additional application 2 to 3 weeks later for downy mildew. Dead Arm (current season infections) — Use 8 tbsp. per gallon when shoots are 1 to 2 inches long. Repeat application when shoots are 4 to 6 inches long.

STRAWBERRY: Botrytis Fruit Rot, Leaf Spot — Use 5 to 8 tbsp. Use sufficient water for thorough coverage. Begin applications when new growth starts and repeat at 7 day intervals. Under conditions favorable to fruit rot, continue applications through harvest, treating immediately after each picking.

TOMATO: Anthracnose, Early Blight, Late Blight, Gray Leaf Spot, Septoria Leaf Spot — Use 5 to 8 tbsp. Use sufficient water for thorough coverage. Under severe infection conditions use 8 to 11 tbsp. Use sufficient water for thorough coverage. Apply at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control.

ROSE: Black Spot — **CARNATION:** Rust, Leaf Spot; **CHRYSANTHEMUM:** Botrytis Flower Blight — Use 8 tsp. and apply thoroughly at first signs of disease. Repeat at 7 to 10 day intervals. Shorten interval during periods of frequent rain and heavy dew.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

NOTICE: Do not reuse empty container. Destroy by perforating and burying in a safe place.
EPA Reg. No. 449-250
Rev. No. 1

Manufactured For
WOODBURY CHEMICAL COMPANY
Division of Techne Corp.
St. Joseph, Mo. 64502; Denver, Colo.