

ACCEPTED

FEB 1957

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR THE CONTROL OF PESTS
 EN 449-251 SUBJECT
 TO COMMENTS

449-251

CAUTION

May cause irritation of nose, throat and skin. Do not breathe dust or spray mist. In case of prolonged or frequent use, wear clean, protective clothing and bathe after each exposure.

Keep containers closed. Store in a cool place. Keep away from fire and

This product will kill fish. Keep out of any body of water. Apply this product only as specified on this label.

ACTIVE INGREDIENTS:

*Zineb	75%
INERT INGREDIENTS:	25%
TOTAL	100%

*Zinc ethylene bisdithiocarbamate (metallic zinc equivalent 17.7%)

CONTENTS 8 OZ.

Woodbury's

WOODBURY
 CHEMICAL COMPANY

ZINEB GARDEN FUNGICIDE

For Black Spot on Roses, and Other Flower Fungi

Manufactured by

WOODBURY CHEMICAL CO.

St. Joseph, Mo.; Denver, Colo.; Orlando, Fla.
 SEE SIDE PANEL FOR ADDITIONAL
 PRECAUTIONARY LABELING

GENERAL INSTRUCTIONS

Use ZINEB at 1 level tablespoonful per gallon. Spray plants thoroughly.

Azaleas and Camellias — Petal blight: Spray at least 2 to 3 times a week while flowers are opening. Direct spray into center of flower.

Gladiolus — Botrytis blight, Curvularia, Stemphylium: Spray at weekly intervals or oftener under severe disease conditions, from early stage until spikes begin to open.

Chrysanthemums, Carnations, Soap dragons, Hollyhocks — Leaf spot, Rust, and Anthracnose: Apply at weekly to 10 day intervals as soon as disease conditions threaten.

Roses — Black spot: Apply at weekly intervals.

USDA-449-251

CAUTION

KEEP OUT OF REACH OF CHILDREN