

**DIRECTIONS**

**\*BEANS:** Mexican Bean Beetle: 1 lb. covers 4300 sq. ft. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles, Japanese Beetle, Velvet Bean Caterpillars: 1 lb. covers 2200 sq. ft. Corn Earworm, Armyworms, Stink Bug, Tarnished Plant Bug: 1 to 1 1/2 lb. covers 2200 sq. feet.

**\*CORN:** Corn Earworm, Corn Rootworm adults, European Corn Borer, Fall Armyworms, Saw Beetles, Japanese Beetle, Flea Beetles, Leafhoppers: 1 to 2 lb. covers 2200 sq. ft.

**\*POTATOES, TOMATOES, EGGPLANT, PEPPERS:** For Colorado Potato Beetle, Flea Beetles, Leafhoppers: 1 to 2 lb. covers 4300 sq. ft. For Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Stink Bugs, Lace Bugs and Lygus Bugs: 1 to 2 lb. covers 2200 sq. ft.

**\*OKRA:** For Corn Earworm, Stink Bugs: 1 to 2 lb. covers 2200 sq. ft.


**\*CUCUMBERS, SQUASH, PUMPKIN, MELONS (Cantaloupe, Casaba, Crenshaw, Honeydew, Honey Ball, Muskmelon, Watermelon):** For Pickworm, Melonworm: 1 to 2 lb. covers 4300 sq. ft. For Cucumber Beetles, Squash Bug, Flea Beetles, Leafhoppers: 1 lb. covers 2200 sq. ft.

**HEAD LETTUCE, CARROTS:** For Flea Beetles, Leafhoppers, Harlequin Bug: 1 to 2 lb. covers 4300 sq. ft. For Six-spotted Leafhopper (Aster Yellows Vector): 1 to 1 1/2 lb. covers 2200 sq. ft. For Corn Earworm, Armyworms, Spittlebugs, Lygus, and Stink Bugs: 1 to 2 lb. covers 2200 sq. ft. Do not apply within 3 days before harvest.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI:** For Flea Beetle, Harlequin Bug: 1 to 2 lb. covers 4300 sq. ft. For Corn Earworm, Armyworms, Cabbage Caterpillars: 1 to 2 lb. covers 2200 sq. ft. (Where Cabbage Looper is the principal problem use an alternative pesticide. For small Loopers after heads form, use 5% Sevin Dust every 5 to 7 days.)

Do not apply within 3 days before harvest.

Net Contents Two Pounds



ACCEPTED

1/14/1967

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. \_\_\_\_\_ SUBJECT TO ATTACHED COMMENTS.

449-307

**5% SEVIN® DUST**

FOR USE ON VEGETABLES, DOGS and CATS


ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) .....	5%
INERT INGREDIENTS .....	95%
TOTAL .....	100%

\*SEVIN—Trademark of Union Carbide Corp.

**CAUTION!**  
KEEP OUT OF REACH OF CHILDREN

Net Contents Two Pounds



ACCEPTED

14/1967

UNDER THE FEDERAL INSECTICIDE ACT  
ECONOMIC POISON REGISTERED  
UNDER NO. SUBJECT  
ATTACHED COMMENTS.

449-307

# 5% SEVIN® DUST

## FOR USE ON VEGETABLES, DOGS and CATS

### ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) .....	5%
INERT INGREDIENTS .....	95%
TOTAL .....	100%

\*SEVIN—Trademark of Union Carbide Corp.

### CAUTION! KEEP OUT OF REACH OF CHILDREN

### CAUTION!

AVOID BREATHING OF DUST.  
AVOID SKIN CONTACT.  
AVOID CONTAMINATION OF FEED AND FOODSTUFFS.  
WASH WITH SOAP AND WATER AFTER USING.

HARMFUL IF SWALLOWED OR INHALED  
OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe the use limitations given on the label for specific crops. Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter. Contamination of productive waters may destroy essential fish foods. Do not apply when weather conditions favor drift of dust from crop areas treated. To protect bees, do not apply when crops are in bloom. Remove or cover hives if dusting is necessary. Apply in early morning or evening when there is little or no wind.

### ANIMALS

**POULTRY:** (Chickens, Partridges, Pheasants, Pigeons): On Birds, for Northern Fowl Mite, Chicken Mite, Fleas, Lice: Use 1 lb. per 100 birds. **IN PREMISES:** For Chicken Mite, Fleas, Bedbugs: Use 1 lb. per 40 sq. ft. floor litter. Do not treat birds within 7 days of slaughter. Do not apply directly to eggs or nest litter. Do not contaminate feed or drinking water. Do not house birds in treated houses within 7 days of slaughter. Do not repeat applications to poultry within 4 weeks.

**DOGS AND CATS:** For Brown Dog Tick, Fleas — rub in skin and apply in sleeping quarters weekly. Do not treat kittens under 4 weeks old. Do not treat puppies under 4 weeks of age.

**NOTE:** All items bearing (\*) may be treated up to and including day of harvest. Items not so marked do not make application within 3 days of harvest.

*PET SLEEPING QUARTERS*

**NOTICE:** Buyer assumes all risk of use, storage, or handling of this material not in strict accordance with directions given herewith.

USDA Reg. No. 449-307



Net Contents Two Pounds



# SEVIN<sup>®</sup> DUST

FOR USE ON VEGETABLES, DOGS and CATS

INGREDIENTS:  
(95% gamma-benzene hexachlorocyclopentadiene  
(5% Diethyl N-methylcarbamate)

5%  
95%  
100%

## CAUTION!

KEEP OUT OF REACH OF CHILDREN

### CAUTION!

AVOID BREATHING OF DUST.  
AVOID SKIN CONTACT.  
AVOID CONTAMINATION OF FEED AND FOODSTUFFS.  
WASH WITH SOAP AND WATER AFTER USING.

HARMFUL IF SWALLOWED OR INHALED  
OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe the use limitations given on the label for specific crops.  
Do not use this product on feed or forage to be fed to dairy animals  
or to livestock being finished for slaughter.  
Contamination of productive waters may destroy essential fish foods.  
Do not apply when weather conditions favor drift of dust from crop  
areas treated.  
To protect bees, do not apply when crops are in bloom. Remove  
or cover hives if dusting is necessary.  
Apply in early morning or evening when there is little or no wind.

### ANIMALS

**POULTRY:** (Chickens, Partridges, Pheasants, Pigeons): On Birds, for  
Northern Fowl Mite, Chicken Mite, Fleas, Lice: Use 1 lb. per  
100 birds. **IN PREMISES:** For Chicken Mite, Fleas, Bedbugs:  
Use 1 lb. per 40 sq. ft. floor litter. Do not treat birds within  
7 days of slaughter. Do not apply directly to eggs or nest  
litter. Do not contaminate feed or drinking water. Do not  
house birds in treated houses within 7 days of slaughter. Do  
not repeat applications to poultry within 4 weeks.

**DOGS AND CATS:** For Brown Dog Tick, Fleas — rub in skin and  
apply in sleeping quarters weekly. Do not treat kittens under  
4 weeks old. Do not treat puppies under 4 weeks of age.

**NOTE:** All items bearing (\*) may be treated up to and including  
day of harvest. Items not so marked do not make application  
within 3 days of harvest.

*PLT SLEEPING QUARTERS*

NOTICE: