



CONTROLS

Control of brush, including Prickly Ash, Black Birch and Scrub Oak on brush and similar non-crop areas. Some susceptible species may require 2 or more applications.

CAUTION

HARMFUL IF SWALLOWED

Avoid contact with skin, eyes or clothing.

Avoid spray drifts to susceptible plants as this may injure cotton, beans, peas, ornamentals, etc. (Coarse sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes (or do not use same spray equipment for other purposes). Do not store near fertilizers, seeds, insecticides or fungicides. Vapors from this product may cause injury to susceptible plants growing nearby.

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

Apply this product only as specified on this label.

Do not reuse container. Destroy by perforating or crushing. Bury or discard in a safe place.

Do not use around the home, recreation areas or similar areas.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

2,4,5-T 4 lb. butyl ester

4 POUNDS 2,4,5-T ACID PER GALLON

For Control of Many Woody Plants

DO NOT USE, POUR, SPILL OR STORE NEAR
HEAT OR OPEN FLAME

ACTIVE INGREDIENTS:

Butyl Ester of 2,4,5-trichlorophenoxyacetic Acid.....	56.4%
INERT INGREDIENTS.....	43.6%
	100.0%

*Equivalent to 4 pounds 2,4,5-trichlorophenoxyacetic acid per gallon.

*Equivalent to 46.2% 2,4,5-trichlorophenoxyacetic acid

CONTENTS GALLON

CAUTION
KEEP OUT OF REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL
CAUTION LABELING

DIRECTIONS

For control of mixed species of brush, mix this product with 2,4-D at the following ratios (to 100 gallons water):

	2,4-D	2,4,5-T
2.64 lbs (40%) Ester.....	4 pts	2½ pts
3.34 lbs Ester.....	3 pts	2½ pts
4.00 lbs Ester.....	2½ pts	2½ pts

and apply to foliage, covering thoroughly (50-100 gallons of solution per acre, depending on density of brush). Apply at full leaf.

As a basal spray (at any time of year), mix this product with 2,4-D at the following rates (to 100 gallons oil):

	2,4-D	2,4,5-T
2.64 lbs (40%) Ester.....	4 gals	2½ gals
3.34 lbs Ester.....	3 gals	2½ gals
4.00 lbs Ester.....	2½ gals	2½ gals

and apply all around plants from ground line up 18 inches, wetting thoroughly. To make a mixture of only 3 gallons, multiply above rates in gallons by 4 to get amounts necessary in ounces.

Do not graze dairy animals on treated areas within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

Stump treatment: Same as basal rates. Best applied to freshly cut stumps.

Observe all cautions and limitations on labeling of all products used in mixtures.

2,4,5-T may be used alone for Brambles, Hawthorn, Osage Orange, Blackjack or Post Oak, Persimmon or Wild Rose at the following dosages: Foliage Spray: 1½ gal per 100 gals water. Do not use more than 1.0 gal of finished mixture per acre on post oak and Basal Spray or Stump Treatment: 3.4 gals per 100 gals oil or 15 lbs to 3 gals oil.

Manufactured by
TECHNE CORPORATION
St. Joseph, MO 64502 Denver, CO 80216
EPA Reg. No. 449-317

ACCEPTED WITH COMMENTS

449-317 - Comments